


Pd^I Dimer CatalysisInternational Edition: DOI: 10.1002/anie.201906063
German Edition: DOI: 10.1002/ange.201906063Arylation of Axially Chiral Phosphorothioate Salts by Dinuclear Pd^I Catalysis

Xiang-Yu Chen, Maoping Pu, Hong-Gang Cheng, Theresa Sperger, and Franziska Schoenebeck*

Abstract: *S*-aryl phosphorothioates are privileged motifs in pharmaceuticals, agrochemicals, and catalysts; yet, the challenge of devising a straightforward synthetic route to enantioenriched *S*-aryl phosphorothioates has remained unsolved to date. We demonstrate herein the first direct C–SP(=O)–(OR')(OR'') coupling of diverse and chiral phosphorothioate salts with aryl iodides, enabled by an air- and moisture-stable Pd^I dimer. Our mechanistic and computational data suggest distinct dinuclear Pd^I catalysis to be operative, which allows for operationally simple couplings with broad scope and full retention of stereochemistry.

While phosphorothioates display privileged structure/activity features and find applications as important agrochemicals, top-selling pharmaceuticals, and powerful chiral catalysts (see Scheme 1 A), the control of their inherent chirality remained a daunting synthetic challenge for decades. Indeed, indirect multistep strategies via P^{III} species have historically been utilized to accomplish this feat,^[1] until Baran's recent announcement of accessing chiral phosphorothioate oligonucleotides by an indirect strategy from limonene derived P^V reagents under auxiliary control (Scheme 1 C).^[2] On the other hand, the synthesis of chiral phosphorothioate salts has been realized through a straightforward three step sequence.^[3] As such, if it would be possible to further derivatize these salts at sulfur, without compromising their stereochemical integrity, then access to the wider phosphorothioate class of compounds would in principle be unleashed.

In this context, the synthesis of chiral aryl phosphorothioates, which are commonly found in drugs, pesticides, and bioactive compounds,^[4] would require the direct coupling of the salt with a C_{sp2} center (Scheme 1 D). If realizable, this strategy would be highly powerful, as it would allow for the introduction of the phosphorothioate moiety in any desired chirality and at any stage in a synthesis. However, although thiolations of C_{sp2} sites, especially aryl halides, are well precedented and can be accomplished by Pd⁰-catalyzed cross coupling reactions,^[5] to date, no general, safe, robust and



Scheme 1. Motivation and synthetic strategy.

direct C_{sp2}–SP(=O)(OR²)(OR³) coupling to access aryl phosphorothioates exists.^[6] Instead, the vast majority of synthetic approaches rely on a bond formation at the phosphorus center directly (S–P coupling, Scheme 1 B),^[7] which has so far not been accomplished in a stereoselective manner.

We therefore set out to develop a method to access diverse and chiral phosphorothioates in a direct coupling approach with widely available aryl halides.

Owing to the general mildness and high functional group tolerance, which are key requirements for late-stage synthetic applications, a Pd⁰/Pd^{II} catalyzed cross-coupling approach appears ideal at first sight to accomplish this goal. However, when we subjected various Pd⁰ catalysts to iodobenzene and NMe₄[SP(=O)(OPh)₂],^[8] such as Pd(PPh₃)₄ or Pd₂(dba)₃/dppf,

[*] X.-Y. Chen, M. Pu, H.-G. Cheng, T. Sperger, F. Schoenebeck
Institute of Organic Chemistry, RWTH Aachen University
Landoltweg 1, 52074 Aachen (Germany)
E-mail: franziska.schoenebeck@rwth-aachen.de

Supporting information and the ORCID identification number(s) for the author(s) of this article can be found under:
<https://doi.org/10.1002/anie.201906063>.

© 2019 The Authors. Published by Wiley-VCH Verlag GmbH & Co. KGaA. This is an open access article under the terms of the Creative Commons Attribution Non-Commercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited, and is not used for commercial purposes.



A Experimental results with Pd⁰/Pd^{II} catalystsB Pd⁰-Pd^{II} catalysis

Figure 1. Free energy profile of Pd^I-catalyzed coupling of PhI with [SP(O)(OPh)₂][−]. Gibbs free energies (in kcal mol^{−1}) shown, calculated at the SMD (toluene) M06L/def2TZVP//B3LYP-D3/6-31G(d, p) (SDD for Pd, I) level of theory.

no coupling was observed (see Figure 1A). Moreover, when we subjected the phosphorothioate salt to a Pd^{II} complex and heated the corresponding mixture, we did not observe the corresponding product **2**.^[10] Our computational studies support this observation,^[9] indicating that the iodide/phosphorothioate exchange at [L_nPd^{II}(Ph)(I)], followed by reductive elimination of Ar–SP(=O)(OPh)₂ has a prohibitively high activation free energy barrier of ΔG[‡] ≥ 30 kcal mol^{−1} and is endergonic. For comparison, the corresponding thiolation, that is, Ar–SPh reductive elimination is characterized by about 10 kcal mol^{−1} lower activation free energy barrier and significantly increased driving force (see Supporting Information for details). As such, and in agreement with the lack of literature precedence, Pd⁰/Pd^{II} catalysis appears to not be an optimal strategy to accomplish C–SP(=O)(OR²)(OR³) coupling.


We previously established that dinuclear Pd^I catalysis is a viable strategy to circumvent potentially poisonous Pd^{II} intermediates as well as alter the driving force of the transformation.^[11] Our extensive mechanistic studies had indicated that the bridging iodides of the dimer are initially exchanged by the nucleophilic coupling partner, and the newly formed Pd^I complex then reacts with the aryl halide, therefore formally reversing the sequence of elementary steps (Figure 1B). As such, we envisioned that a dinuclear Pd^I coupling protocol^[11,12] might offer a solution to the phosphorothioate problem.

To assess whether Pd^I-dimer catalyzed phosphorothioation would be feasible, we undertook computational studies of the coupling with iodobenzene, which clearly indicated that there is a pronounced driving force to convert iodobenzene to the C–SP(=O)(OPh)₂ coupled product (Figure 1). The acti-

vation barrier was found to be in the range of our previous findings of dinuclear Pd^I catalysis,^[11,12] and as such appeared to be feasible.

A key requirement of the Pd^I dimer catalysis concept is that the coupling partner will also function as a stabilizing bridging unit in the Pd^I dimer. To examine this, we subjected 4 equiv. of tetramethyl ammonium phosphorothioate salt to a solution of the air-stable, iodide bridged Pd^I dimer **1**. To our delight, within 2 hours at room temperature we saw the appearance of a signal at 97.9 ppm upon ³¹P-NMR spectroscopic analysis, which is very reminiscent of a functionalized

Table 1: Scope of the Pd^I-catalyzed phosphorothioation of aryl iodides.^[a]




[a] Yields of isolated products **2–27** after chromatography. [b] The reaction mixture was stirred for 36 hours.

Pd^{I} dimer, suggesting that the iodide in **1** can be exchanged for a phosphorothioate.^[13]

We next set out to explore whether Pd^{I} catalyzed phosphorothioation of aryl iodides is also feasible. To our delight, the desired O,O-diphenyl phosphorothioate **2**^[14] was obtained in 83 % yield in the presence of 5 mol % of $[\text{Pd}(\mu\text{-I})(\text{P}t\text{Bu}_3)_2]$ dimer **1** (Table 1). Encouraged by these findings, we subsequently turned to explore the wider scope of the transformation (Table 1). A series of aryl iodides bearing electron-donating or electron-withdrawing substituents ($\text{R} = 4\text{-Me}$, $4\text{-}n\text{Bu}$, $4\text{-}t\text{Bu}$, 4-F , 4-Cl and 4-Ph) reacted smoothly and gave the corresponding products **3–8** in good yields. The reactions of sterically hindered *ortho*-substituted (2-MeO , 2-Me , and 2-Ph) aryl iodides also proceeded well, with slightly lower yield observed with *meta* substitution (**9–12**). For **10**, prolonging the reaction time increased the product yield. The reactions of disubstituted substrates were also successful without any loss of yield (**13** and **14**). This was also true for the polyaromatic iodides giving rise to products **15** and **16**.

Table 2: Applications of the coupling strategy to the efficient synthesis of axially chiral S-aryl phosphorothioates.^[a]



[a] Yields of isolated products **28–39** after chromatography and the *ee* determined by HPLC analysis of the purified product on a chiral stationary phase.

Moreover, the method tolerated a variety of functional groups, such as esters and morpholines (**17–19**).

The effect of O,O-dialkyl phosphorothioate salt was examined as well. Various substituted aryl iodides were readily coupled under the standard reaction conditions and the corresponding products **20–24** were obtained in excellent yields. Notably, the chirality in the substituents of the salt was retained (**25–27**).

The P^{V} -series of TADDOL derived organophosphorus compounds are usually found in various valuable chiral catalysts and ligands.^[15] To further expand the synthetic utility of the present catalytic strategy, the synthesis of axially chiral S-aryl phosphorothioates was tested (Table 2). Gratifyingly, uniformly excellent enantioselectivities of up to >99 % *ee* were obtained using C2-symmetric TADDOL-derived phosphorothioate salts with a variety of aryl iodides, resulting in the desired products **28–39** in very good yields. The absolute configuration of the product **30** was determined by X-ray crystallographic analysis (Table 2), unambiguously confirming the complete retention of chirality.^[16]

In conclusion, while the examined $\text{Pd}^0/\text{Pd}^{\text{II}}$ catalysis failed to deliver the phosphorothioation of aryl iodides as a consequence of disfavored exchange at Pd^{II} and a high reductive elimination barrier, the mechanistically distinct dinuclear Pd^{I} catalysis operates with an orthogonal driving force that allows for the first catalytic S-arylation of phosphorothioate salts with aryl iodides. The method is characterized by generality, broad scope, and operational simplicity, making use of an easily accessible phosphorothioate salt and an air-stable Pd^{I} dimer. The stereochemical integrity of chiral salts was not compromised in the coupling process, allowing the efficient assembly of axially chiral S-aryl phosphorothioates. In light of the promising properties of S-aryl phosphorothioate compounds as well as the privileged Pd^{I} catalysis reactivity, we anticipate that the presented methodology will facilitate numerous applications in the agrochemical, pharmaceutical, and stereoselective synthesis arenas.

Acknowledgements

We thank the RWTH Aachen and the European Research Council (ERC-637993) for funding. Calculations were performed with computing resources granted by JARA-HPC from RWTH Aachen University under project “jara0091”.

Conflict of interest

The authors declare no conflict of interest.

Keywords: arylation · axial chirality · computation · dinuclear catalysis · S-aryl phosphorothioate

How to cite: *Angew. Chem. Int. Ed.* **2019**, *58*, 11395–11399
Angew. Chem. **2019**, *131*, 11517–11521

[1] a) W. J. Stec, A. Grajkowski, A. Kobylanska, B. Karwowski, M. Koziolkiewicz, K. Misiura, A. Okruszek, A. Wilk, P. Guga, M.

- Boczkowska, *J. Am. Chem. Soc.* **1995**, *117*, 12019–12029; b) W. J. Stec, B. Karwowski, M. Boczkowska, P. Guga, M. Koziolkiewicz, M. Sochacki, M. W. Wiecek, J. Błaszczyk, *J. Am. Chem. Soc.* **1998**, *120*, 7156–7167; c) N. Oka, M. Yamamoto, T. Sato, T. Wada, *J. Am. Chem. Soc.* **2008**, *130*, 16031–16037; d) N. Oka, T. Kondo, S. Fujiwara, Y. Maizuru, T. Wada, *Org. Lett.* **2009**, *11*, 967–970; e) Y. Nukaga, K. Yamada, T. Ogata, N. Oka, T. Wada, *J. Org. Chem.* **2012**, *77*, 7913–7922; f) N. Iwamoto, D. C. D. Butler, N. Svrikapa, S. Mohapatra, I. Zlatev, D. W. Y. Sah, Meena, S. M. Standley, G. Lu, L. H. Apponi, M. Frank-Kamenetsky, J. J. Zhang, C. Vargeese, G. L. Verdine, *Nat. Biotechnol.* **2017**, *35*, 845.
- [2] K. W. Knouse, J. N. deGruyter, M. A. Schmidt, B. Zheng, J. C. Vantourout, C. Kingston, S. E. Mercer, I. M. McDonald, R. E. Olson, Y. Zhu, C. Hang, J. Zhu, C. Yuan, Q. Wang, P. Park, M. D. Eastgate, P. S. Baran, *Science* **2018**, *361*, 1234–1238.
- [3] C. R. Hall, T. D. Inch, *J. Chem. Soc. Perkin Trans. 1* **1979**, 1104–1111.
- [4] a) L. L. Murdock, T. L. Hopkins, *J. Agric. Food Chem.* **1968**, *16*, 954–958; b) A.-h. Lee, R. L. Metcalf, *Pestic. Biochem. Physiol.* **1973**, *2*, 408–417; c) A.-H. Lee, R. L. Metcalf, G. M. Booth, *Ann. Entomol. Soc. Am.* **1973**, *66*, 333–343; d) M. Sekine, T. Hata, *J. Am. Chem. Soc.* **1986**, *108*, 4581–4586; e) H. Hotoda, M. Koizumi, R. Koga, M. Kaneko, K. Momota, T. Ohmine, H. Furukawa, T. Agatsuma, T. Nishigaki, J. Sone, S. Tsutsumi, T. Kosaka, K. Abe, S. Kimura, K. Shimada, *J. Med. Chem.* **1998**, *41*, 3655–3663; f) Y. Qiu, J. C. Worch, D. N. Chirdon, A. Kaur, A. B. Maurer, S. Amsterdam, C. R. Collins, T. Pintauer, D. Yaron, S. Bernhard, K. J. T. Noonan, *Chem. Eur. J.* **2014**, *20*, 7746–7751.
- [5] a) *Handbook of organopalladium chemistry for organic synthesis* (Eds.: E.-i. Negishi, A. de Meijere), Wiley, New York, **2002**; b) *Metal-Catalyzed Cross-Coupling Reactions*, 2nd ed. (Eds.: F. Diederich, A. de Meijere), Wiley-VCH, Weinheim, **2004**; c) *Organotransition metal chemistry: from bonding to catalysis* (Eds.: J. F. Hartwig, J. P. Collman), University Science Books, Sausalito, **2010**; d) C. C. C. Johansson Seechurn, M. O. Kitching, T. J. Colacot, V. Snieckus, *Angew. Chem. Int. Ed.* **2012**, *51*, 5062–5085; *Angew. Chem.* **2012**, *124*, 5150–5174; e) *New trends in cross-coupling: theory and applications* (Eds.: T. J. Colacot), Royal Society of Chemistry, London, **2014**; f) T. Kondo, T.-a. Mitsudo, *Chem. Rev.* **2000**, *100*, 3205–3220; g) J. F. Hartwig, *Acc. Chem. Res.* **2008**, *41*, 1534–1544; h) C. Valente, M. Pompeo, M. Sayah, M. G. Organ, *Org. Process Res. Dev.* **2014**, *18*, 180–190; i) M. A. Fernández-Rodríguez, Q. Shen, J. F. Hartwig, *J. Am. Chem. Soc.* **2006**, *128*, 2180–2181; j) E. Alvaro, J. F. Hartwig, *J. Am. Chem. Soc.* **2009**, *131*, 7858–7868; k) M. Sayah, M. G. Organ, *Chem. Eur. J.* **2011**, *17*, 11719–11722; l) G. Teverovskiy, D. S. Surry, S. L. Buchwald, *Angew. Chem. Int. Ed.* **2011**, *50*, 7312–7314; *Angew. Chem.* **2011**, *123*, 7450–7452; m) J. L. Farmer, M. Pompeo, A. J. Lough, M. G. Organ, *Chem. Eur. J.* **2014**, *20*, 15790–15798; n) C. Uyeda, Y. Tan, G. C. Fu, J. C. Peters, *J. Am. Chem. Soc.* **2013**, *135*, 9548–9552; o) M. Jiang, H. Li, H. Yang, H. Fu, *Angew. Chem. Int. Ed.* **2017**, *56*, 874–879; *Angew. Chem.* **2017**, *129*, 892–897; p) B. Liu, C.-H. Lim, G. M. Miyake, *J. Am. Chem. Soc.* **2017**, *139*, 13616–13619; q) D. Liu, H.-X. Ma, P. Fang, T.-S. Mei, *Angew. Chem. Int. Ed.* **2019**, *58*, 5033–5037; *Angew. Chem.* **2019**, *131*, 5087–5091; r) Y. Wang, L. Deng, X. Wang, Z. Wu, Y. Wang, Y. Pan, *ACS Catal.* **2019**, *9*, 1630–1634.
- [6] For a single kind of phosphorothioate, that is, a non-chiral bisdimethoxy substituted example, the coupling of aryl diazonium salts under Cu-catalysis has been demonstrated and suggested to proceed via organic free radicals, which could pose challenges for the conservation of stereochemical information as well as late-stage synthetic applicability. For details, see: a) S. Kovács, B. Bayarmagnai, A. Aillerie, L. J. Gooben, *Adv. Synth. Catal.* **2018**, *360*, 1913–1918; For catalytic methods with elemental sulfur, which bears inherent challenges for retention of stereochemistry at phosphorus, see: b) L. Zhang, P. Zhang, X. Li, J. Xu, G. Tang, Y. Zhao, *J. Org. Chem.* **2016**, *81*, 5588–5594; c) J. Xu, L. Zhang, X. Li, Y. Gao, G. Tang, Y. Zhao, *Org. Lett.* **2016**, *18*, 1266–1269.
- [7] a) D. C. Morrison, *J. Am. Chem. Soc.* **1955**, *77*, 181–182; b) F. W. Hoffmann, T. R. Moore, B. Kagan, *J. Am. Chem. Soc.* **1956**, *78*, 6413–6414; c) J. Michalski, T. Modro, J. Wiecekowsky, *J. Chem. Soc.* **1960**, 1665–1670; d) J. Michalski, J. Wasiak, *J. Chem. Soc.* **1962**, 5056–5061; e) R. G. Harvey, H. I. Jacobson, E. V. Jensen, *J. Am. Chem. Soc.* **1963**, *85*, 1618–1623; f) C. E. Dreef, C. M. Dreef-Tromp, G. A. van der Marel, J. H. van Boom, *Synlett* **1990**, 481–483; g) D.-D. Liu, D.-w. Chen, Z.-C. Chen, *Synth. Commun.* **1992**, *22*, 2903–2908; h) L. N. Pribytkova, M. S. Erzhanova, *Chem. Nat. Compd.* **1994**, *30*, 193–196; i) C. M. Timperley, S. A. Saunders, J. Szpalek, M. J. Waters, *J. Fluorine Chem.* **2003**, *119*, 161–171; j) M. Arisawa, T. Ono, M. Yamaguchi, *Tetrahedron Lett.* **2005**, *46*, 5669–5671; k) B. Kaboudin, F. Farjadian, *Beilstein J. Org. Chem.* **2006**, *2*, 4; l) Y.-X. Gao, G. Tang, Y. Cao, Y.-F. Zhao, *Synthesis* **2009**, 1081–1086; m) B. Kaboudin, Y. Abedi, J.-y. Kato, T. Yokomatsu, *Synthesis* **2013**, *45*, 2323–2327; n) Y.-J. Ouyang, Y.-Y. Li, N.-B. Li, X.-H. Xu, *Chin. Chem. Lett.* **2013**, *24*, 1103–1105; o) J. Bai, X. Cui, H. Wang, Y. Wu, *Chem. Commun.* **2014**, *50*, 8860–8863; p) G. Kumaraswamy, R. Raju, *Adv. Synth. Catal.* **2014**, *356*, 2591–2598; q) Y.-C. Liu, C.-F. Lee, *Green Chem.* **2014**, *16*, 357–364; r) S. Mitra, S. Mukherjee, S. K. Sen, A. Hajra, *Bioorg. Med. Chem. Lett.* **2014**, *24*, 2198–2201; s) D. S. Panmand, A. D. Tiwari, S. S. Panda, J.-C. M. Monbaliu, L. K. Beagle, A. M. Asiri, C. V. Stevens, P. J. Steel, C. D. Hall, A. R. Katritzky, *Tetrahedron Lett.* **2014**, *55*, 5898–5901; t) J. Wang, X. Huang, Z. Ni, S. Wang, Y. Pan, J. Wu, *Tetrahedron* **2015**, *71*, 7853–7859; u) Y. Zhu, T. Chen, S. Li, S. Shimada, L.-B. Han, *J. Am. Chem. Soc.* **2016**, *138*, 5825–5828; v) Y. Moon, Y. Moon, H. Choi, S. Hong, *Green Chem.* **2017**, *19*, 1005–1013; w) S. Song, Y. Zhang, A. Yeerlan, B. Zhu, J. Liu, N. Jiao, *Angew. Chem. Int. Ed.* **2017**, *56*, 2487–2491; *Angew. Chem.* **2017**, *129*, 2527–2531; x) H. Huang, J. Ash, J. Y. Kang, *Org. Biomol. Chem.* **2018**, *16*, 4236–4242; y) D. J. Jones, E. M. O'Leary, T. P. O'Sullivan, *Tetrahedron Lett.* **2018**, *59*, 4279–4292.
- [8] G. Hilgetag, H. Teichmann, *J. Prakt. Chem.* **1959**, *8*, 104–111.
- [9] Calculations were performed at the SMD (toluene) M06L/def2TZVP//B3LYP-D3/6-31G(d, p) (SDD for Pd, I) level of theory using Gaussian09 (Revision D.01), M. J. Frisch et al. (see Supporting Information for full reference). For appropriateness of applied computational model, see: T. Sperger, I. A. Sanhueza, I. Kalvet, F. Schoenebeck, *Chem. Rev.* **2015**, *115*, 9532–9586.
- [10] A previous attempt to react $(\text{PPh}_3)_2\text{Pd}(\text{Ph})(\text{Br})$ with KSP(=O)(OEt)₂ also showed no reaction, see: Okamoto, J. B. Housekeeper, C. K. Luscombe, *Appl. Organomet. Chem.* **2013**, *27*, 639–643.
- [11] a) G. Yin, I. Kalvet, F. Schoenebeck, *Angew. Chem. Int. Ed.* **2015**, *54*, 6809–6813; *Angew. Chem.* **2015**, *127*, 6913–6917; b) M. Aufiero, T. Sperger, A. S.-K. Tsang, F. Schoenebeck, *Angew. Chem. Int. Ed.* **2015**, *54*, 10322–10326; *Angew. Chem.* **2015**, *127*, 10462–10466; c) T. Scattolin, E. Senol, G. Yin, Q. Guo, F. Schoenebeck, *Angew. Chem. Int. Ed.* **2018**, *57*, 12425–12429; *Angew. Chem.* **2018**, *130*, 12605–12609.
- [12] a) K. J. Bonney, F. Proutiere, F. Schoenebeck, *Chem. Sci.* **2013**, *4*, 4434–4439; b) I. Kalvet, K. J. Bonney, F. Schoenebeck, *J. Org. Chem.* **2014**, *79*, 12041–12046; c) I. Kalvet, G. Magnin, F. Schoenebeck, *Angew. Chem. Int. Ed.* **2017**, *56*, 1581–1585; *Angew. Chem.* **2017**, *129*, 1603–1607; d) I. Kalvet, T. Sperger, T. Scattolin, G. Magnin, F. Schoenebeck, *Angew. Chem. Int. Ed.* **2017**, *56*, 7078–7082; *Angew. Chem.* **2017**, *129*, 7184–7188; e) T. Sperger, C. K. Stirner, F. Schoenebeck, *Synthesis* **2017**, *49*, 115–120; f) S. T. Keaveney, G. Kundu, F. Schoenebeck, *Angew. Chem.*

Int. Ed. **2018**, 57, 12573–12577; *Angew. Chem.* **2018**, 130, 12753–12577.

- [13] See Supporting Information for details.
- [14] It remains challenging to access O,O-diphenyl phosphorothioates by traditional methods, see ref. [7m].
- [15] a) R. Hilgraf, A. Pfaltz, *Adv. Synth. Catal.* **2005**, 347, 61–77; b) S. A. Moteki, D. Wu, K. L. Chandra, D. S. Reddy, J. M. Takacs, *Org. Lett.* **2006**, 8, 3097–3100; c) T. Akiyama, *Chem. Rev.* **2007**, 107, 5744–5758; d) *Syntheses of simple TADDOL-like phosphite, phosphonite and phosphonate chiral ligands* (Eds.: J. P. Perotti, R. M. Cravero, L. E. Luna, R. J. Grau, S. E. Vaillard), University of Michigan Library and Michigan Publishing, Ann Arbor, **2011**.
- [16] CCDC 1911726 (**30**) contains the supplementary crystallographic data for this paper. These data can be obtained free of charge from The Cambridge Crystallographic Data Centre.

Manuscript received: May 15, 2019

Accepted manuscript online: June 13, 2019

Version of record online: July 9, 2019