



RESEARCH PAPER

Inducible biosynthesis and immune function of the systemic acquired resistance inducer *N*-hydroxypipecolic acid in monocotyledonous and dicotyledonous plants

Anika Schnake^{1,*}, Michael Hartmann^{1,*}, Stefan Schreiber¹, Jana Malik¹, Lisa Brahmman¹, Ipek Yildiz¹, Janina von Dahlen², Laura E. Rose^{2,4}, Ulrich Schaffrath³, and Jürgen Zeier^{1,4,†} 

¹ Institute for Molecular Ecophysiology of Plants, Department of Biology, Heinrich Heine University, Universitätsstraße 1, D-40225 Düsseldorf, Germany

² Institute for Population Genetics, Department of Biology, Heinrich Heine University, Universitätsstraße 1, D-40225 Düsseldorf, Germany

³ Department of Plant Physiology, RWTH Aachen University, D-52056 Aachen, Germany

⁴ Cluster of Excellence on Plant Sciences (CEPLAS), Heinrich Heine University, Universitätsstraße 1, D-40225 Düsseldorf, Germany

* These authors contributed equally to this work.

† Correspondence: juergen.zeier@hhu.de

Received 7 April 2020; Editorial decision 29 June 2020; Accepted 2 July 2020

Editor: Steven Spoel, University of Edinburgh, UK

Abstract

Recent work has provided evidence for the occurrence of *N*-hydroxypipecolic acid (NHP) in *Arabidopsis thaliana*, characterized its pathogen-inducible biosynthesis by a three-step metabolic sequence from L-lysine, and established a central role for NHP in the regulation of systemic acquired resistance. Here, we show that NHP is biosynthesized in several other plant species in response to microbial attack, generally together with its direct metabolic precursor pipecolic acid and the phenolic immune signal salicylic acid. For example, NHP accumulates locally in inoculated leaves and systemically in distant leaves of cucumber in response to *Pseudomonas syringae* attack, in *Pseudomonas*-challenged tobacco and soybean leaves, in tomato inoculated with the oomycete *Phytophthora infestans*, in leaves of the monocot *Brachypodium distachyon* infected with bacterial (*Xanthomonas translucens*) and fungal (*Magnaporthe oryzae*) pathogens, and in *M. oryzae*-inoculated barley. Notably, resistance assays indicate that NHP acts as a potent inducer of acquired resistance to bacterial and fungal infection in distinct monocotyledonous and dicotyledonous species. Pronounced systemic accumulation of NHP in leaf phloem sap of locally inoculated cucumber supports a function for NHP as a phloem-mobile immune signal. Our study thus generalizes the existence and function of an NHP resistance pathway in plant systemic acquired resistance.

Keywords: *Brachypodium*, cucumber, *Magnaporthe*, *N*-hydroxypipecolic acid, pipecolic acid, plant immunity, *Pseudomonas*, salicylic acid, systemic acquired resistance, tobacco.

Introduction

Plant metabolites play diverse basic roles in plant defense, since they can act as antimicrobial weapons to directly harm attacking pathogens, contribute to the fortification of plant cell walls to impede the entrance of pathogens into plant

tissue, and coordinate the inducible expression of components of plant immune responses (Ahuja *et al.*, 2012; Berens *et al.*, 2017; Chezem *et al.*, 2017). Salicylic acid (SA) is a central metabolic activator of plant immunity that guarantees the

effective induction of resistance responses in the defense of plants against bacterial, fungal, oomycete, and viral infections. In many species, the levels of SA and SA derivatives such as SA- β -glucoside (SAG) and SA glucose ester in the leaf tissue of unstressed plants are low but significantly increase in response to pathogen attack (Klessig *et al.*, 2018). In the dicot model *Arabidopsis thaliana*, genetic and biochemical evidence indicates that this stress-inducible biosynthesis of SA proceeds via conversion of the shikimate pathway product chorismate to isochorismate (Wildermuth *et al.*, 2001).

Systemic acquired resistance (SAR) is an inducible defense response of plants that is triggered by a localized microbial attack in leaves and within a few days renders the whole foliage immune to subsequent pathogen infection (Shah and Zeier, 2013). It is well established that SA is an important resistance-enhancing metabolite in the systemic immunity of dicotyledonous plants such as *Arabidopsis* or tobacco, because plants that are unable to accumulate SA show strongly attenuated SAR (Gaffney *et al.*, 1993; Nawrath and Métraux, 1999; Bernsdorff *et al.*, 2016). However, previous grafting experiments with tobacco and time-course studies with *Pseudomonas syringae*-inoculated cucumber have suggested that SA is not the primary SAR inducer that transmits resistance signaling from inoculated leaves to distantly located systemic leaves. Instead, the existence of another SAR signal has been proposed (Rasmussen *et al.*, 1991; Vernooij *et al.*, 1994). More recent research has provided evidence that the lysine-derived metabolite N-hydroxypipecolic acid (NHP) is a critical activator of systemic immunity in *Arabidopsis* and that its endogenous biosynthesis is indispensable for pathogen-induced SAR (Hartmann *et al.*, 2018). In addition, a function for NHP as a mobile inducer of SAR has been suggested (Chen *et al.*, 2018).

NHP is not synthesized in unstressed *Arabidopsis* plants but markedly accumulates systemically in the leaves of pathogen-inoculated plants (Hartmann *et al.*, 2018). Three pathogen-inducible genes are involved in the biosynthesis of NHP in *Arabidopsis*: *AGD2-LIKE DEFENSE RESPONSE PROTEIN1* (*ALD1*), *SAR-DEFICIENT4* (*SARD4*), and *FLAVIN-DEPENDENT-MONOOXYGENASE1* (*FMO1*) (Hartmann and Zeier, 2019). *ALD1* functions as an L-Lys- α -aminotransferase and thus deaminates L-Lys to ϵ -amino- α -ketocaproic acid, which spontaneously cyclizes to dehydropipecolic acid intermediates. These are reduced by *SARD4* to pipecolic acid (Pip) (Ding *et al.*, 2016; Hartmann *et al.*, 2017). The non-protein amino acid Pip (homoproline) accumulates to high levels in locally inoculated and systemic leaf tissue of *Arabidopsis*, and this accumulation is necessary for SAR (Návarová *et al.*, 2012). Pip is also synthesized by an *ALD1*-/*SARD4*-mediated L-Lys catabolic pathway in the fir clubmoss *Huperzia serrata* (Xu *et al.*, 2018). Moreover, Pip has been shown to accumulate in several angiosperms other than *Arabidopsis* upon contact with a pathogen—for example, in soybean inoculated with the fungi *Rhizoctonia solani* and *Fusarium virguliforme* (Aliferis *et al.*, 2014; Abeyssekara *et al.*, 2016), in tobacco mosaic virus- or *P. syringae*-inoculated tobacco (*Nicotiana tabacum*) (Pálfi and Dézsi, 1968; Vogel-Adzhoghue *et al.*, 2013; Ádám *et al.*, 2018), in rice infected with

the blast fungus *Magnaporthe oryzae* (Pálfi and Dézsi, 1968), and in *P. syringae*-infected barley (Lenk *et al.*, 2019).

In *Arabidopsis*, elevated Pip was found to trigger SAR in an *FMO1*-dependent manner (Návarová *et al.*, 2012; Bernsdorff *et al.*, 2016). The monooxygenase *FMO1*, which was first identified as an essential player in pathogen-induced SAR in *Arabidopsis* (Mishina and Zeier, 2006), was more recently shown to catalyze the N-hydroxylation of Pip to NHP (Hartmann *et al.*, 2018; Chen *et al.*, 2018). When exogenously supplied to *Arabidopsis*, NHP potently activated SAR to bacterial (*P. syringae*) and oomycete (*Hyaloperonospora arabidopsidis*) infections and overrode the SAR incompetence of *fmo1* mutant plants (Hartmann *et al.*, 2018). These findings demonstrated that NHP, as the end product of the described three-step L-Lys catabolic pathway, functions as the active, SAR-inducing metabolite of this immune pathway. When locally applied to lower leaves, NHP also induced resistance in upper leaves, suggesting a function in leaf-to-leaf signal transmission in SAR (Chen *et al.*, 2018).

The natural occurrence of NHP in living organisms was described for the first time in *Arabidopsis* (Hartmann *et al.*, 2018). Since the NHP biosynthetic precursor Pip is naturally present in many different angiosperms (Broquist, 1991; Zeier, 2013), and genes homologous to *Arabidopsis FMO1* exist in monocots and dicots (Hartmann *et al.*, 2018; Chen *et al.*, 2018), it is conceivable that NHP biosynthesis and its function in SAR are conserved across angiosperms. In *Arabidopsis*, the induction of an effective SAR response by NHP requires its interplay with SA (Bernsdorff *et al.*, 2016; Hartmann *et al.*, 2018; Hartmann and Zeier, 2019). However, it is unclear so far whether the molecular mechanisms of SAR elucidated for *Arabidopsis* and other dicots, as well as the function of SA in systemic immunity, hold true for monocot species (Balmer *et al.*, 2013; Dey *et al.*, 2014).

In this study, we show that the pathogen-inducible biosynthesis of NHP and its ability to induce plant SAR is present in distinct monocotyledonous and dicotyledonous plant species. Our results generalize the existence of an NHP-associated immune pathway in plants and highlight mechanistic similarities in the SAR of dicot and monocot plants.

Materials and methods

Plant material and growth conditions

Arabidopsis thaliana accession Col-0 plants were grown as outlined previously (Hartmann *et al.*, 2018). Plants of six species (*Cucumis sativus* cv. Wisconsin SMR58, *Nicotiana tabacum* cv. Xanthi, *Solanum lycopersicum* cv. Heinz1706, *Glycine max* cv. Maple Arrow, *Brachypodium distachyon* accession BD21, and *Hordeum vulgare* cv. Ingrid) were grown in environmentally controlled growth chambers on a mixture of soil, vermiculite, and sand with a 16 h/8 h light/dark cycle (see Supplementary Table S1 at JXB online for specific growth conditions). Plants of *S. lycopersicum* cv. M82 infected with *Phytophthora infestans* were cultivated as described previously (de Vries *et al.*, 2017).

Cultivation of pathogens

Pseudomonas syringae pv. *lachrymans* strain 814/98 (Psl) (Slomnicka *et al.*, 2018), *P. syringae* pv. *tomato* (Pst) DC3000, *P. syringae* pv. *maculicola* ES4326 (Psm), Psm carrying the *Photorhabdus luminescens* luxCDABE operon (Psm

lux) (Hartmann et al., 2018), and *Pseudomonas savastanoi* pv. *glycinea* (DSM 50267) (*Psg*) were grown at 28 °C in King's B medium containing the appropriate antibiotics (*Pst* DC3000 and *Psm*, 50 µg l⁻¹ rifampicin; *Psm lux*, 50 µg l⁻¹ of rifampicin and 25 µg l⁻¹ kanamycin; *Psl* and *Psg*, no antibiotics). *Pseudomonas syringae* pv. *tabaci* (DSM 1856) (*Pstb*) was grown at 18 °C in King's B medium without antibiotics (Vogel-Adghough et al., 2013). *Xanthomonas translucens* (DSM 18974) was grown at 28 °C on peptone-sucrose agar (20 g l⁻¹ sucrose, 5 g l⁻¹ peptone, 0.5 g l⁻¹ K₂HPO₄, 0.25 g l⁻¹ MgSO₄·7H₂O, 15 g l⁻¹ agar). *Phytophthora infestans* strain IPO-C was cultured in the dark at 18 °C on rye sucrose agar as described previously (de Vries et al., 2017). *Magnaporthe oryzae* strain Guy11 was cultivated on cornmeal agar supplemented with a nitrate salt solution at 18 °C (9 h light) as reported by Parker et al. (2008). The *M. oryzae* isolate TH6772 was cultured on oatmeal agar plates (20 g l⁻¹ agar, 2 g l⁻¹ yeast extract, 10 g l⁻¹ starch, 30 g l⁻¹ oat flakes) at 23 °C in the dark (Delventhal et al., 2017).

Plant inoculation experiments

For bacterial inoculation experiments, log-phase cultures of the respective bacteria were grown overnight under constant shaking (240 rpm). The cultures were washed three times with 10 mM MgCl₂ and diluted to different final optical density levels at 600 nm (OD₆₀₀) for leaf inoculations. Mock-control plants were inoculated likewise with 10 mM MgCl₂ solution, unless stated otherwise.

Arabidopsis thaliana

Leaves of 5-week-old *Arabidopsis* Col-0 plants were infiltrated with a suspension of *Psm lux* (OD₆₀₀=0.001) as described previously (Hartmann et al., 2018).

Cucumis sativus

Three-week-old *C. sativus* plants were spray-inoculated with a suspension of *Psl* of OD₆₀₀=0.2 (supplemented with 0.005% of the surfactant Silwet® L-77) applied to the abaxial side of the first true leaf. For the collection of *C. sativus* phloem sap, locally inoculated (first true) or distal (second true) leaves were cut and the escaping sap was collected at the site of the petiole by using filter paper. Filter paper soaked with ~10 µl of phloem sap was transferred into 100 µl MeOH/H₂O (80:20, v/v) for metabolite extraction. Leaf discs (0.5 cm²) and phloem sap samples for subsequent metabolite analyses were collected at 1, 3, and 5 days post-inoculation (dpi), frozen in liquid nitrogen, and stored at -80 °C (n=5). For the resistance assays, discs of inoculated *C. sativus* leaves were harvested at 3 dpi (n=4) and 6 dpi (n=4).

Nicotiana tabacum

Nicotiana tabacum plants 4 to 5 weeks old were carefully pressure-infiltrated from the abaxial side of the leaves with a *Pstb* or *Psm* suspension, using a needleless syringe. For metabolite analyses, discs (1.0 cm²) of *N. tabacum* leaves inoculated with *Pstb* (OD₆₀₀=0.005) were harvested from 1 to 3 dpi, frozen in liquid nitrogen, and stored at -80 °C (n=3). For resistance assays, leaf discs of *Pstb* [OD₆₀₀=0.001 (n=12)] and *Psm* [OD₆₀₀=0.001 (n=25)] inoculated plants were sampled at 3 dpi.

Glycine max

Three-week-old *G. max* plants at growth stage 12–13 (BBCH scale) were pressure-infiltrated as described above with a *Psg* suspension (OD₆₀₀=0.1). Samples of inoculated *G. max* leaves were harvested at 9 dpi as described above.

Solanum lycopersicum

For bacterial inoculations, 4-week-old *S. lycopersicum* cv. Heinz1706 plants were pressure-infiltrated with a suspension of *Pst* DC3000 (OD₆₀₀=0.005) from the abaxial side of fully expanded leaves, using a needleless syringe. Sampling for metabolite analyses was performed at

1 dpi and 2 dpi (n=3) as described above. For oomycete inoculations of *S. lycopersicum* cv. M82, a suspension of *P. infestans* containing 50 spores µl⁻¹ was used for droplet inoculation (10 µl droplet applied to each leaf) of 4-week-old plants grown in closed vessels (de Vries et al., 2017). Mock controls were treated likewise with sterile H₂O. For the metabolite analyses of *P. infestans*- and mock-inoculated *S. lycopersicum* cv. M82 leaves, samples were harvested at 1 dpi and 2 dpi (n≥4) as described above.

Brachypodium distachyon

For metabolite analyses, 5-week-old *B. distachyon* plants were inoculated with a suspension of *X. translucens* (OD₆₀₀=0.5). Sampling (one plant per sample) was carried out at 1, 3, and 5 dpi (n=3). Leaf samples were frozen in liquid nitrogen and stored at -80 °C. For resistance assays, 3-week-old *B. distachyon* plants were inoculated with a bacterial suspension of OD₆₀₀=0.005. The assessment of resistance was performed at 3 dpi (n=9). In both experimental setups, the bacterial suspension was infiltrated from the adaxial site of the leaves, using a needleless syringe.

For the inoculation of *B. distachyon* with *M. oryzae*, a suspension of conidia (50 spores µl⁻¹) was applied to 3- to 4-week-old plants by airbrush spray-inoculation. To prepare the spore suspension, spores were isolated within 15 days of growth on plates by washing the mycelium with sterile H₂O and scraping off the spores using a coverslip. The resulting solution was filtered through Miracloth (Merck Millipore, Billerica, MA, USA) and centrifuged for 4 min at 2200 g. After resuspending the pellet in 3 ml sterile H₂O, the concentration of spores was determined using a Neubauer chamber (NanoEnTek, Inc., Seoul, Korea). Plants were sprayed until they were covered evenly with fine droplets. After inoculation, the plants were maintained hermetically sealed in autoclave bags as described by Parker et al. (2008). For metabolite analyses, one plant per sample was harvested at 2, 3, and 4 dpi (n≥4). The assessment of resistance against *M. oryzae* was performed at 4 dpi.

Hordeum vulgare

Magnaporthe oryzae conidia were harvested from oatmeal agar plates after incubation for 2 weeks under UV-A light and the suspension was adjusted to 250 spores µl⁻¹. Primary leaves of barley plants were spray-inoculated with a spore suspension containing 1 g l⁻¹ gelatin and 0.5 ml l⁻¹ Tween. After spray-inoculation, plants were kept for 24 h at 24 °C and 100% relative humidity in the dark and then transferred to growth chamber conditions. Mock-control plants were spray-inoculated with H₂O containing 1 g l⁻¹ gelatin and 0.5 ml l⁻¹ Tween. Inoculated *H. vulgare* leaves were sampled at 5 dpi (n=3).

Plant treatments with N-hydroxyproline

Arabidopsis thaliana

NHP treatment of *Arabidopsis* via the root was performed by pipetting 10 ml of aqueous NHP solutions of different concentrations on to the soil of individually cultivated 5-week-old plants (Hartmann et al., 2018). For spray treatments of *Arabidopsis* leaves, the whole rosette of 5-week-old plants was sprayed with a 1 mM aqueous solution of NHP, supplemented with 0.005% Silwet® L-77, until small droplets formed evenly on the leaf surfaces (~0.5 ml solution per plant). For treatments by leaf infiltration, three rosette leaves per plant were infiltrated with 1 mM NHP solution. Application of H₂O by the respective method served as a control treatment for all treatment types. Leaves were inoculated with *Psm lux* 1 day after the NHP treatments.

Nicotiana tabacum

Treatments of tobacco plants with NHP were performed following the procedure described by Vogel-Adghough et al. (2013) for Pip solutions. Individually cultivated plants were watered with 10 ml of a 1 mM aqueous solution of NHP by pipetting. As a control treatment, 10 ml H₂O was applied in the same manner. Inoculation of leaves with *Pstb* or *Psm* was performed 24 h after the application of the NHP treatment.

Cucumis sativus

Approximately 1 ml of 1 mM NHP solution was directly injected into the plant stem using a syringe with a needle. An injection of sterile H₂O served as a control. One day later, the leaves were spray-inoculated with a *Pst* suspension.

Brachypodium distachyon

1 mM NHP solution was syringe-infiltrated thoroughly from the adaxial site into the leaves of *B. distachyon* until the entire leaf area appeared to be soaked. Plants were inoculated with *X. translucens* or *M. oryzae* 1 day later. Infiltration with H₂O was used as a control treatment.

Assessment of resistance

The resistance of *Arabidopsis* against *Psm lux* infection was determined as described previously (Hartmann *et al.*, 2018). For other bacterial growth assays, leaf discs were cut out of inoculated leaf material and homogenized in 10 mM MgCl₂. Each sample was diluted in 10 mM MgCl₂ (1:40 000 for *Pst*/*C. sativus*; 1:20 000 for *Pstb*/*N. tabacum*; 1:5000 for *Psm*/*N. tabacum*; 1:400 000 for *X. translucens*/*B. distachyon*), plated on agar plates (1.5% agar) containing the above-described media, and incubated for 2–4 days at 28 °C before the number of bacterial colonies was determined.

To assay the resistance of *B. distachyon* to *M. oryzae*, the youngest fully grown leaves at the time of inoculation were removed and bleached using an ethanol–chloroform solution (4:1) containing 1.5% trichloroacetic acid. After a minimum of 2 days, the bleaching solution was removed with H₂O and the leaves were photographed on a light table. The proportion of the necrotic leaf area was measured using ImageJ (<https://imagej.nih.gov/ij/>).

Analysis of plant metabolites by gas chromatography-mass spectrometry

To determine the levels of Pip, NHP, SA, SAG, NHP-hexose, and lysine in the leaves of the different species, the gas chromatography-mass spectrometry-based analytical procedure previously established for *Arabidopsis* was used (detailed in Hartmann *et al.*, 2018; Stahl *et al.*, 2019). For metabolic analyses of cucumber phloem sap, the above-described filter paper extracts were supplemented with 900 µl of MeOH/H₂O (80:20, v/v) and analysed.

Analysis of gene expression

The expression of the *C. sativus* *ALD1* (XM_011651615.1), *FMO1* (XM_004144232.1), and *Actin* (XM_004147305.2) genes was assessed by semi-quantitative RT-PCR analysis. RNA extraction and cDNA synthesis were performed as described by Wang *et al.* (2018). Primers and PCR conditions are listed in Supplementary Table S2. PCRs were performed using GoTaq Green Master Mix (Promega) following the manufacturer's instructions.

Statistical analyses and reproducibility

The presented results are derived from experimental datasets consisting of the number of biological replicates indicated in the figure legends and/or in the above sections. For the statistical analyses of bacterial growth data, log₁₀-transformed colony-forming units or relative light units values were used. ANOVA with type II sum of squares was applied for multiple comparisons (Bernsdorff *et al.*, 2016). For pairwise comparisons of two distinct treatments, two-tailed Student's *t*-tests were performed. The presented results were generally confirmed in at least two other independent experiments, except for the barley infection experiment (Fig. 5C), which was conducted once.

Results

We previously reported the identification of the hitherto unknown plant natural product NHP, its FMO1-mediated

biosynthesis from Pip, its pathogen-inducible accumulation, and its biological function as a SAR inducer in the model plant *A. thaliana* (Hartmann *et al.*, 2018). On this basis, we were interested to investigate whether the inducible biosynthesis of NHP and its function in acquired resistance were a common feature in angiosperms. To generalize the occurrence and function of the NHP pathway in plant immunity, we implemented different naturally relevant plant–pathogen interaction systems in the laboratory. We further conducted defense-related metabolite analyses and resistance assays to test whether exogenously applied NHP would induce acquired resistance in species other than *Arabidopsis*.

NHP induces SAR in *Arabidopsis* at low doses and by different treatment modes

In our previous study on the role of NHP in SAR, we observed that exogenous NHP applied via the soil to individual *Arabidopsis* plants in doses of 10 µmol (i.e. 10 ml of a 1 mM NHP solution) induced a strong SAR effect in the foliage (Hartmann *et al.*, 2018). To get more information about the dose dependency of this NHP-triggered resistance response, we supplied NHP solutions at concentrations ranging from 0.1 to 10 mM to individual *Arabidopsis* plants by the same method. We then inoculated leaves of the plants with the compatible bacterial strain *Psm lux* 1 day later and scored bacterial numbers at 2.5 dpi. Watering with a concentration of 0.1 mM NHP attenuated bacterial growth by ~3-fold, indicating that doses of NHP as low as 1 µmol applied via the roots induce a detectable SAR effect in *Arabidopsis* (Fig. 1A). Increases in the concentration of NHP up to 1 mM gradually enhanced this SAR response to more than one log difference compared with the control treatment, and increases to 10 mM resulted in further modest improvements of the SAR effect (Fig. 1A). Watering the soil with 1 mM NHP solution generally resulted in a stable SAR effect that reduced bacterial growth in the leaves by 10–50-fold (Fig. 1A, B; Hartmann *et al.*, 2018).

Next, to test whether treating leaves directly with NHP would yield similar resistance effects, we supplied NHP to the foliage by spraying and by infiltration. Spraying the whole leaf rosette with 1 mM NHP solution induced strong resistance and attenuated bacterial growth in infected leaves by a factor of ~50-fold compared with control plants (Fig. 1B). Because the overall volume applied to a single plant was ~0.5 ml for the spray application, the effective dose of NHP to induce a strong SAR was ~0.5 µmol per plant for leaf treatments. Moreover, when individual leaves were pre-infiltrated with 1 mM NHP solution, the same leaves acquired resistance to subsequent *Psm lux* infection, and bacterial growth at 2.5 dpi was again reduced by more than one order of magnitude (Fig. 1C). Notably, the untreated distant leaves of plants that had been pre-infiltrated with NHP in other leaves showed induction of a very strong SAR (Fig. 1C), corroborating the previously established critical role of NHP in SAR long-distance communication (Chen *et al.*, 2018; Hartmann *et al.*, 2018). Together, these results show that NHP applied by different modes of treatment is able to induce SAR and exerts a strong biological activity in doses lower than 1 µmol.




Fig. 1. Exogenous NHP induces SAR in Arabidopsis by different treatment modes. (A) Dose dependency of SAR in leaves triggered by treatment of roots with NHP. The cultivation soil of individually grown 5-week-old Arabidopsis Col-0 plants was watered with 10 ml of an aqueous NHP solution at the indicated concentration. Treatment with H₂O served as the control. One day later, three leaves of each plant were inoculated with the compatible bioluminescent *Pseudomonas syringae* pv. *maculicola* (*Psm lux*) strain (OD₆₀₀=0.001). Bacterial numbers in leaves were assessed at 2.5 dpi by measurement of luminescence and are expressed as relative light units (rlu) per cm² leaf area (Hartmann et al., 2018). Data represent the mean ±SD of at least 21 leaf replicates from 7 or more plants.

NHP accumulates systemically in the foliage and in phloem sap of cucumber, and induces SAR in this species

Arabidopsis has developed into the primary model plant for SAR research in the past two to three decades (e.g. Fu and Dong, 2013; Shah and Zeier, 2013). In the most studied Arabidopsis accession, Col-0, the basal levels of the SAR-associated metabolites SA, Pip, and NHP are generally low (SA, Pip) or undetectable (NHP) in unstressed plants, and rise strongly during the course of infection with *P. syringae* in both locally inoculated and distal systemic leaf tissue (Hartmann and Zeier, 2019). In addition, cucumber was used in a series of studies as a model to investigate the molecular mechanisms and physiology of the SAR response (e.g. Métraux et al., 1990; Rasmussen et al., 1991). Cucumber plants inoculated with compatible lesion-inducing bacterial, fungal, or viral phytopathogens in one leaf show induction of robust SAR and systemic defense responses in distal leaves. Cucumber and other species of Cucurbitaceae also release phloem sap from cut petiole or stem tissues that can readily be analyzed for its molecular content, including potentially phloem-mobile SAR-active metabolites (Métraux et al., 1990; Smith-Becker et al., 1998).

We inoculated *C. sativus* cv. Wisconsin SMR58 plants at the two-leaf stage in their first true leaf with the causal agent of angular leaf spot, *PsI* (Ślommnicka et al., 2018), a pathogen that elicits systemic defense responses in cucumber (Mölders et al., 1994). SA and its β-glucoside SAG accumulated significantly in the *PsI*-inoculated first leaves at 3 dpi and 5 dpi, whereas the systemic accumulation of SA in the distal leaves was modest, and that of SAG was not significant (Fig. 2A, B; Supplementary Fig. S1A). The basal levels of Pip were between 10 and 25 μg g⁻¹ fresh weight (FW) in uninfected cucumber leaves (Fig. 2A, B), markedly higher than the basal levels of Pip in leaves of Arabidopsis Col-0 (~0.5 μg g⁻¹ FW; Návarová et al., 2012). In contrast to the response in Arabidopsis, pathogen inoculation did not result in increases of Pip either locally or systemically (Fig. 2A, B). Consistent with this constitutively strong accumulation of Pip, transcripts of the *C. sativus* *ALD1* gene were readily detectable in the leaves of control plants (Supplementary Fig. S2). However, in contrast to the amount of Pip in uninfected

Different letters denote significant differences ($P < 0.05$; ANOVA and post-hoc Tukey's honestly significant difference test). (B) Comparison of soil application and foliar spray treatment of Col-0 plants with NHP solutions. Left panel: Col-0 plants were watered with 10 ml of a 1 mM NHP solution or H₂O. Right panel: Whole leaf rosettes were sprayed with ~0.5 ml of a 1 mM NHP solution or H₂O (both supplemented with 0.005% Silwet® L-77) until the adaxial side of the leaves was evenly covered in fine droplets of the solution. One day later, the leaves were inoculated with *Psm lux* and bacterial numbers were quantified at 2.5 dpi. Statistically significant differences between NHP-pretreated and control plants are indicated with asterisks: *** $P < 0.001$ (two-tailed *t* test). (C) Infiltration of individual Col-0 rosette leaves with NHP solution induces SAR in the infiltrated leaves and in distant leaves. Three leaves of a given Col-0 plant were syringe-infiltrated with 1 mM NHP solution or H₂O. One day later, either the pretreated (1°) leaves or three distant (2°) leaves were inoculated with *Psm lux* and bacterial numbers were quantified at 2.5 dpi (*** $P < 0.001$; two-tailed *t* test).




Fig. 2. NHP systemically accumulates in the leaves and petiole sap of *Cucumis sativus* upon inoculation with *Pseudomonas syringae* pv. *lachrymans*. (A) Levels of SA, Pip, and NHP in *Cucumis sativus* cv. Wisconsin SMR58 leaves inoculated with *P. syringae* pv. *lachrymans* (*Psl*) and in mock-treated control leaves. The first grown true leaves of plants at the two-leaf stage were spray-inoculated with a suspension of *Psl* ($\text{OD}_{600}=0.2$, containing 0.005% Silwet® L-77) or with 0.005% Silwet® L-77 dissolved in 10 mM MgCl_2 (mock treatment). Samples of treated leaves were collected at 1, 3, or 5 dpi. Metabolite levels are given in $\mu\text{g g}^{-1}$ leaf FW and represent the mean \pm SD of four or five biological replicates ($n=4$ or 5). (B) Levels of SA, Pip, and NHP in the second, distal leaves of cucumber plants inoculated on the first leaves with *Psl* or infiltrated with 10 mM MgCl_2 as mock treatment ($n=4$ or 5). (C) Concentrations of SA, Pip, and NHP in petiole sap collected from *Psl*-inoculated or mock-treated *C. sativus* leaves. Metabolite concentrations are given in $\mu\text{g ml}^{-1}$ collected petiole exudate ($n=4$ or 5). (D) Concentrations of SA, Pip, and NHP in petiole sap collected from the second, distal leaves of cucumber plants in which the lower, first leaves had been *Psl*-inoculated or mock-treated ($n=4$ or 5). Statistically significant differences between samples from pathogen-inoculated and mock-treated leaves at a given time point are indicated with asterisks: *** $P<0.001$, ** $P<0.01$, * $P<0.05$ (two-tailed t -test).

cucumber plants, the levels of NHP were low ($<0.05 \mu\text{g g}^{-1}$ FW). Notably, NHP accumulated markedly in both the inoculated ($2.2 \mu\text{g g}^{-1}$ FW) and distant ($1.4 \mu\text{g g}^{-1}$ FW) leaves in response to *Psl* at 3 dpi and 5 dpi (Fig. 2A, B). The pathogen-induced accumulation of NHP was paralleled by a further increase of *ALD1* transcripts in *Psl*-inoculated leaves at 3 dpi (Supplementary Fig. S2). Moreover, transcripts of the *C. sativus* *FMO1* gene, which were not detected in the leaves of control plants, accumulated at 3 dpi and 5 dpi (Supplementary Fig. S2). As previously reported for Arabidopsis (Hartmann and Zeier, 2018), we also detected a putative NHP-hexose conjugate in extracts from *Psl*-inoculated cucumber leaves (Supplementary Fig. S1B). Together, these results show that cucumber possesses an intact and pathogen-inducible NHP biosynthetic pathway.

SAR long-distance signal propagation from one leaf to another is supposedly mediated by signal-active, phloem-mobile metabolites (Shah and Zeier, 2013). We collected phloem sap exuded from the petioles of cut local (first true) or distant (second true) leaves following *Psl* inoculation or mock treatment and analyzed the contents of SAR-relevant metabolites (Fig. 2C, D). Both SA ($\sim 2 \mu\text{g ml}^{-1}$) and SAG ($\sim 8 \mu\text{g ml}^{-1}$) were significantly enriched in the phloem sap of *Psl*-inoculated local leaves, and a pathogen-induced increase of SAG content was also detected in phloem sap collected from distant leaves (Fig. 2C, D; Supplementary Fig. S1A). Notably, the constitutive levels of Pip in phloem sap were in the range $1\text{--}4 \text{ mg ml}^{-1}$ ($\sim 8\text{--}30 \text{ mM}$) and therefore about three orders of magnitude higher than the contents of SA/SAG in the petiole sap from inoculated leaves. As was observed for the leaf contents, *Psl* inoculation did not result in further increases in these prominent levels of Pip in phloem sap from local or distant leaves (Fig. 2C, D). By contrast, the levels of NHP were barely detectable in phloem sap from mock-treated control plants. Notably, NHP levels increased largely in the sap of both local ($\sim 80 \mu\text{g ml}^{-1}$) and distant ($\sim 8 \mu\text{g ml}^{-1}$) leaves upon *Psl* inoculation (Fig. 2C, D). Together, these results show that localized inoculation of a pathogen triggers a strong local and systemic accumulation of NHP in both the leaves and the leaf phloem sap of cucumber. This accumulation is greater than the increases of SA and SAG. This observation supports a function of NHP in SAR-associated phloem-based long-distance signaling. The NHP biosynthetic precursor Pip is constitutively present in high amounts in both leaves and phloem sap of cucumber.

To test whether elevated levels of NHP trigger acquired resistance to *Psl* infection in cucumber, we injected an aqueous solution of NHP (1 mM) into the epicotylar *C. sativus* stem as an inducing pretreatment and challenge-inoculated the first true leaves with *Psl* 1 day later. Plants pretreated with NHP showed strongly enhanced resistance to *Psl* leaf infection compared with H_2O -pretreated control plants (Fig. 3A, B). NHP-pretreated plants hosted 50- and 100-fold lower amounts of bacteria in the *Psl*-inoculated leaves at 3 dpi and 6 dpi, respectively (Fig. 3A). In addition, the yellowish and chlorotic disease symptoms observed on leaves of control cucumber plants were greatly diminished as a consequence of NHP pretreatment (Fig. 3B). Together, these results indicate that NHP is able to induce SAR in cucumber.

Biosynthesis and resistance function of NHP in other dicotyledonous plant species

Pip, the direct biosynthetic precursor of NHP, occurs as a natural product in a wide array of angiosperms (Broquist, 1991; Hartmann and Zeier, 2018). Several studies have demonstrated that Pip also accumulates in the foliage of various dicotyledonous species other than Arabidopsis in response to pathogen inoculation. For example, biosynthesis of Pip is induced in *N. tabacum* leaves in response to infection by the causal agent of tobacco wildfire disease, *Pstb*, and by tobacco mosaic virus (Pálfi and Dézsi, 1968; Vogel-Adzhough et al., 2013; Ádám et al., 2018). We examined whether NHP is detectable and accumulates, alongside Pip and SA, in *Pstb*-inoculated tobacco leaves. The basal levels of SA and Pip were low in the leaves of control (mock-treated) *N. tabacum* cv. Xanthi plants, but their biosynthesis was significantly increased in response to *Pstb* inoculation. In addition, NHP, which was not detected in the leaves of mock-treated tobacco plants, accumulated substantially in leaves of *Pstb*-inoculated plants, to amounts of ~ 5 and $8 \mu\text{g g}^{-1}$ FW at 48 and 72 h post inoculation (hpi), respectively (Fig. 4A). NHP also accumulated in *S. lycopersicum* cv. Heinz1706 leaves from non-detectable basal levels to $1.8 \mu\text{g g}^{-1}$ FW at 48 h after inoculation with the bacterial speck pathogen *Pst* DC3000 (Fig. 4B). Similarly, inoculation of *S. lycopersicum* cv. M82 leaves with the oomycete *P. infestans*, the causal agent of tomato late blight, resulted in the accumulation of NHP to $\sim 5 \mu\text{g g}^{-1}$ FW at 48 hpi (Fig. 4C). In both interactions, tomato leaves started to synthesize NHP from ~ 24 hpi onwards and showed markedly induced accumulation of Pip and SA alongside the biosynthesis of NHP (Fig. 4B, C).

Previous studies also indicated the inducible accumulation of Pip in soybean plants inoculated with the fungal pathogens *Rhizoctonia solani* and *Fusarium virguliforme* (Aliferis et al., 2014; Abeysekara et al., 2016). Consequently, we determined metabolite levels in *G. max* cv. Maple Arrow leaves inoculated with the bacterial blight pathogen *Psg*. Leaves of uninoculated soybean plants contained constitutive levels of both SA and Pip in the range of $1 \mu\text{g g}^{-1}$ FW, whereas NHP was not detectable. *Psg* inoculation resulted in a significant elevation of the levels of SA and Pip, as well as a noticeable accumulation of NHP to $\sim 1.3 \mu\text{g g}^{-1}$ FW (Fig. 4D).

Previous findings indicate that feeding of Pip to *N. tabacum* plants enhances resistance to leaf infection by *Pstb* and tobacco mosaic virus (Vogel-Adzhough et al., 2013; Ádám et al., 2018). We therefore tested whether exogenous NHP would induce acquired resistance of tobacco to bacterial infection by employing the soil-watering method of NHP pretreatment (Hartmann et al., 2018), which was used previously for supplementing *N. tabacum* plants with Pip (Vogel-Adzhough et al., 2013). NHP-pretreated plants showed a significant restriction of *Pstb* growth after challenge inoculation and also showed fewer disease symptoms compared with naive plants, indicating that NHP significantly induces resistance to this bacterial strain (Fig. 3C, E). In experiments in which the same doses of Pip and NHP were supplied to different plants, NHP pretreatment reduced disease severity even more markedly than Pip feeding (Supplementary Fig. S3). We also inoculated *N. tabacum* cv.




Fig. 3. *Cucumis sativus* and *Nicotiana tabacum* plants pretreated with a micromolar dose of NHP acquire resistance to bacterial infection. (A) Growth of *Pseudomonas syringae* pv. *lachrymans* (*Psl*) in *C. sativus* leaves. Three-week-old cucumber plants were pretreated with NHP by injecting 1 ml of 1 mM aqueous NHP solution into the stem. Injection of H₂O served as a control. One day later, leaves were spray-inoculated with *Psl* (OD₆₀₀=0.2). Bacterial numbers in leaves, expressed as colony-forming units (cfu) per cm² leaf surface, were scored at 3 dpi and 6 dpi. Data represent the mean ±SD of five biological replicates (*n*=5). Statistically significant differences between bacterial growth in the leaves of NHP- and H₂O-pretreated plants are indicated with asterisks: ****P*<0.001, ***P*<0.01 (two-tailed *t*-test). (B) Representative disease symptoms shown by *C. sativus* plants pretreated with H₂O or 1 mM NHP at 6 days post spray-inoculation with *Psl*. Arrows indicate the inoculated first leaves. (C) Growth of *P. syringae* pv. *tabaci* (*Pstb*) in *N. tabacum* cv. Xanthi leaves. Plants were pretreated with NHP by the addition of 10 ml of 1 mM NHP to the cultivation soil. Control plants were watered with 10 ml H₂O. Leaves were infiltrated with a suspension of *Pstb* (OD_{600nm}=0.001) 1 day later; bacterial growth was assessed at 3 dpi. Data represent the mean ±SD of at least 11 biological replicates (*n*=11). (D) Growth of *P. syringae* pv. *maculicola* (*Psm*) in leaves of *N. tabacum* cv. Xanthi after infiltration with a suspension of OD₆₀₀=0.001. The experimental setup was as described in C (*n*=11). (E) Representative disease symptoms shown by *Pstb*-inoculated *N. tabacum* leaves from plants pretreated with 10 ml H₂O or 1 mM NHP at 6 dpi.

Xanthi plants with *Psm*, which infects *Arabidopsis* and causes yellowish and chlorotic symptoms on leaves (Mishina and Zeier, 2007). The *Psm* strain was able to grow in tobacco leaf tissue, but its multiplication in naive plants was approximately one order of magnitude lower than that of the compatible *Pstb* strain at 3 dpi, and the interaction remained essentially symptomless (Fig. 3C, D). Nevertheless, NHP pretreatment strongly attenuated the growth of *Psm* in *N. tabacum* cv. Xanthi leaves (Fig. 3D). Together, these data show that NHP effectively induces resistance of tobacco to bacterial pathogen infection.

In sum, our results show that NHP biosynthesis is undetectable in unstressed plants of the dicotyledonous species we have investigated in this and previous studies. However, NHP significantly accumulates in leaves of members of

the Brassicaceae (*Arabidopsis*; Hartmann *et al.*, 2018), Cucurbitaceae (cucumber), Solanaceae (tobacco, tomato), and Fabaceae (soybean) in response to bacterial and/or oomycete infection (Figs 2, 4). In addition, our findings indicate that low micromolar doses of NHP activate acquired resistance in *Arabidopsis* (Fig. 1; Hartmann *et al.*, 2018), cucumber, and tobacco (Fig. 3) to bacterial or oomycete infection.

NHP is biosynthesized and induces acquired resistance in monocotyledonous plants

The Poaceae species *B. distachyon* has been adopted and developed into a useful model for investigating the biology of monocotyledonous plants and cereal crops, including the study




Fig. 4. NHP accumulates in leaves of *Nicotiana tabacum*, *Solanum lycopersicum*, and *Glycine max* inoculated with bacterial and/or oomycete pathogens. (A) Levels of SA, Pip, and NHP in leaves of *N. tabacum* cv. Xanthi inoculated with *Pseudomonas syringae* pv. *tabaci* (*Pstb*) and in leaves of mock-treated control plants. Leaves of 4-week-old plants were infiltrated with a bacterial suspension at OD=0.005 and samples were collected at 48 and 72 h post inoculation (hpi). Leaf metabolite levels are given in µg g⁻¹ FW (*n*=3). nd, Not detected. Other details are as described in Fig. 2A. (B) Levels of SA, Pip, and NHP in leaves of *S. lycopersicum* cv. Heinz1706 inoculated with *P. syringae* pv. *tomato* (*Pst*) DC3000 and in leaves of mock-treated control plants. Leaves of 3-week-old plants were infiltrated with a bacterial suspension of OD=0.005 and samples were collected at 24 and 48 hpi (*n*=3). (C) Levels of SA, Pip, and NHP in leaves of *S. lycopersicum* cv. M82 inoculated with *P. infestans* IPO-C and in leaves of mock-treated control plants. Leaves of 3-week-old plants were inoculated with droplets containing 50 spores µl⁻¹. Samples were collected at 24 and 48 hpi (*n*≥4). (D) Levels of SA, Pip, and NHP in leaves of *G. max* cv. Maple Arrow inoculated with *Pseudomonas savastanoi* pv. *glycinea* (*Psg*) and in leaves of mock-treated control plants. Leaves of 3-week-old plants were infiltrated with a bacterial suspension of OD=0.1 and samples were collected at 9 dpi (*n*=4). Data are presented as mean ±SD. Statistically significant differences between samples from pathogen-inoculated and mock-treated leaves at a given time point are indicated with asterisks: ****P*<0.001, ***P*<0.01, **P*<0.05 (two-tailed *t*-test).




Fig. 5. Pathogen-inducible activation of the pipecolate metabolic pathway in response to bacterial and fungal infection in leaves of the monocots *Brachypodium distachyon* and *Hordeum vulgare*. (A) Levels of SA, Pip, and NHP in leaves of *B. distachyon* BD21 inoculated with *Xanthomonas translucens* and in leaves of mock-treated plants. Leaves of 3-week-old plants were infiltrated with a bacterial suspension of OD₆₀₀=0.5 and samples were collected at 1, 3, and 5 dpi ($n=3$). Other details are as described in Fig. 2A. (B) Levels of SA, Pip, and NHP in leaves of *B. distachyon* BD21 inoculated with *Magnaporthe oryzae* strain Guy11 and in leaves of mock-treated plants. Leaves were spray-inoculated with a suspension of fungal spores (50 spores μl^{-1}) and samples were collected at 2, 3, and 4 dpi ($n=3$). nd, Not detected. (C) Levels of SA, Pip, and NHP in leaves of *H. vulgare* cv. Ingrid inoculated with *M. oryzae* isolate TH6772 and in leaves of mock-treated plants. Leaves were spray-inoculated with a suspension of 250 conidia μl^{-1} and samples were collected at 5 dpi ($n=3$). Data are presented as mean \pm SD. Statistically significant differences between samples from pathogen-inoculated and mock-treated leaves at a given time point are indicated with asterisks: *** $P<0.001$, ** $P<0.01$, * $P<0.05$ (two-tailed t -test).

of plant immune responses (Fitzgerald *et al.*, 2015; Scholthoff *et al.*, 2018). To investigate the occurrence and possible immune function of the NHP pathway in *B. distachyon* BD21, we inoculated leaves of 3-week-old plants with the bacterial leaf streak pathogen *X. translucens* (Fitzgerald *et al.*, 2015). SA started to accumulate from relatively low basal levels in the *B. distachyon* leaves at 1 dpi and reached values of 3–4 $\mu\text{g g}^{-1}$ FW during later stages of infection (3 dpi and 5 dpi; Fig. 5A).

These marked increases in the levels of free SA were accompanied by a massive accumulation of SAG and an elevation of the level of SA glucose ester from 3 dpi onwards (Supplementary Fig. S4). The basal levels of Pip in *B. distachyon* leaves, at 1–2 $\mu\text{g g}^{-1}$ FW, were higher than those generally observed for *Arabidopsis* Col-0 leaves (Návarová *et al.*, 2012), and Pip levels started to accumulate significantly only at later stages of infection (5 dpi), to $\sim 4 \mu\text{g g}^{-1}$ FW (Fig. 5A). By contrast,

NHP levels rose markedly from very low basal levels ($<0.03 \mu\text{g g}^{-1}$ FW) to moderate levels ($\sim 0.2 \mu\text{g g}^{-1}$ FW) at 3 dpi (Fig. 5A).

We also inoculated leaves of *B. distachyon* BD21 plants with the rice blast fungus *M. oryzae*, which has previously been shown to establish compatibility with *B. distachyon* (Parker et al., 2008, 2009). Accumulation of SA in *B. distachyon* leaves was observable in this interaction, but was less pronounced than that in response to *X. translucens* (Fig. 5B). By contrast, the

accumulation of Pip was quantitatively higher for the interaction with *M. oryzae* than for that with *X. translucens* at 5 dpi (Fig. 5B). Similarly, NHP accumulated to somewhat greater amounts in the course of *M. oryzae* infection and reached values of $\sim 0.6 \mu\text{g g}^{-1}$ FW at 5 dpi (Fig. 5B).

In addition, we analyzed the levels of defense-related metabolites in *Hordeum vulgare* cv. Ingrid inoculated with the compatible *M. oryzae* isolate TH6772 and at a progressed stage of




Fig. 6. NHP induces acquired resistance to bacterial and fungal infection in *Brachypodium distachyon*. (A) Growth of *Xanthomonas translucens* in leaves of *B. distachyon* BD21. Plants were pretreated by infiltrating 1 mM NHP solution or H₂O (control) into the leaves. One day later, leaves were pressure-inoculated with a suspension of *X. translucens* of OD₆₀₀=0.5. Bacterial numbers in leaves were assessed at 3 dpi ($n=10$). Other details are as described in Fig. 3A. (B) Representative disease symptoms of *X. translucens*-inoculated *B. distachyon* leaves from plants pretreated with H₂O or 1 mM NHP at 3 dpi and 8 dpi. (C) Representative stained *B. distachyon* BD21 leaves at 4 dpi after inoculation with *M. oryzae*. Leaves of plants were pretreated with NHP or H₂O as described in A. One day later, the leaves were spray-inoculated with *M. oryzae* conidia. Brownish areas represent necrotic disease symptoms as a consequence of *M. oryzae* infection. (D) Disease scoring of *B. distachyon* BD21 leaves inoculated with *M. oryzae* after pretreatment with H₂O or NHP. At 4 dpi, leaves were harvested and stained, and the percentage of necrotic area per leaf surface was determined as a measure of disease severity. Statistically significant differences in bacterial growth (A) or leaf necrotic areas (D) between NHP- and H₂O-pretreated plants are indicated with asterisks: *** $P<0.001$, * $P<0.05$ (two-tailed *t*-test).

disease (5 dpi; Delventhal *et al.*, 2017). The basal levels of the three metabolites under investigation were low or undetectable in barley leaves, and SA accumulation was not observed after fungal inoculation. However, we detected a strong accumulation of Pip to $\sim 30 \mu\text{g g}^{-1}$ FW at 5 dpi, and NHP accumulated to moderate but clearly detectable levels ($0.1 \mu\text{g g}^{-1}$ FW) in *M. oryzae*-infected barley leaves (Fig. 5C).

In summary, our metabolite analyses indicate that monocot plants are able to activate the pipelate pathway to synthesize both Pip and NHP upon bacterial and fungal inoculation. The SA biosynthetic pathway is induced in parallel, but the degree of induction depended on the particular plant–pathogen interaction.

To investigate whether NHP also acts as an immune-active substance in monocots, we pretreated the leaves of *B. distachyon* BD21 plants with 1 mM NHP solution or water, inoculated them with *X. translucens* or *M. oryzae* 1 day later, and scored bacterial growth and fungal disease progression in NHP-treated and control plants. With *X. translucens*, the leaves of the control plants hosted ~ 10 -fold higher bacterial numbers than the leaves of NHP-pretreated plants at 3 dpi (Fig. 6A). Moreover, the pronounced brownish necrotic disease symptoms that were evident on the infected leaves of control plants were rarely observed on the leaves of NHP-pretreated plants (Fig. 6B). Following inoculation with suspensions of *M. oryzae* conidia, leaves of *B. distachyon* control plants developed numerous dark brown necrotic lesions at infection sites (Parker *et al.*, 2008). As a measure of disease severity, we assessed the necrotic areas that emerged at 4 dpi on leaf surfaces by quantifying the extent of brownish areas of chlorophyll–destained leaves (Fig. 6C). We observed that pretreatment with NHP significantly attenuated the otherwise widely developing necrotic lesions of *M. oryzae*-inoculated leaves (Fig. 6C, D). Together, these resistance assays indicate that NHP can act as a potent inducer of acquired resistance against leaf infection by both the bacterium *X. translucens* and the fungus *M. oryzae* in the monocot *B. distachyon*.

Discussion

NHP was identified as a naturally occurring metabolite for the first time in the dicot model species *A. thaliana*. In this plant, NHP accumulated markedly in response to leaf infection by *P. syringae* in both locally inoculated and distal uninfected leaves (Hartmann *et al.*, 2018). Phylogenetic analyses of publicly available genomic and proteomic databases on the basis of *A. thaliana* sequence information suggest a widespread occurrence of homologs of the core NHP biosynthetic pathway genes across the plant kingdom, encompassing monocotyledonous and dicotyledonous plant species alike, many of which are crop species with high commercial or nutritional value (Hartmann *et al.*, 2018; Holmes *et al.*, 2019). In the present study, we show that NHP accumulates in the local and systemic leaves of cucumber upon bacterial inoculation (Fig. 2). Further, NHP biosynthesis and accumulation were observed in the leaves of *P. syringae*-challenged tobacco and tomato, *P. infestans*-infected tomato, *B. distachyon* infected with *X. translucens* and *M. oryzae*, and *M. oryzae*-inoculated

barley (Figs 4, 5). In addition to our work, a recent study detected NHP accumulation in tobacco, tomato, and mustard plants when treated with *Pst* DC3000 bacteria, although absolute NHP levels were not determined (Holmes *et al.*, 2019). Therefore, both monocot and dicot plants have the capacity to induce NHP biosynthesis in response to microbial attack. Moreover, leaf infections by diverse bacterial, oomycete, and fungal pathogens can trigger plant NHP biosynthesis.

A common feature of all the species analyzed so far is that NHP biosynthesis is inducible and that the levels of NHP in leaves of unstressed naive plants are very low or, in most cases, not detectable (Hartmann *et al.*, 2018; Figs 2, 4, 5). Thus, in the absence of biotic challenge, plants apparently keep NHP at very low basal levels. Two recent studies have reported that Arabidopsis lines or mutant plants with constitutively elevated levels of NHP show permanently activated defenses and dwarfed phenotypes: specifically, lines overexpressing the calcium-dependent protein kinase CPK5 and mutants in which *CAMTA3*, a negative immune-regulatory gene, is defective (Guerra *et al.*, 2020; Sun *et al.*, 2020). When crossed with the *fmo1* mutant, both lines lose their dwarfism and their ability to synthesize NHP, indicating that the growth retardation is caused by constitutively elevated levels of NHP. This observation suggests that the usually observed low level of the resistance activator NHP in naive plants is a prerequisite for maximal growth performance. The inducible character of NHP biosynthesis would ensure that acquired resistance is switched on only when plants encounter a (biotic) stress stimulus and in this way limit constitutive growth impairment. A time-course analysis in Arabidopsis revealed that the levels of NHP reach a maximum over the course of *P. syringae*-triggered accumulation and later decrease, as would be expected for a signaling molecule (Hartmann *et al.* 2018; Hartmann and Zeier, 2019). Biochemical modification, such as through glycosylation, very likely contributes to the lower levels of free NHP after its stress-induced accumulation (Chen *et al.*, 2018; Hartmann and Zeier, 2018). Accordingly, we detected a stress-inducible NHP–hexose conjugate in Arabidopsis and in cucumber (Hartmann and Zeier, 2018; Supplementary Fig. S1B). However, the levels of this conjugate were below the limit of detection in the other species under investigation.

Pip functions as the direct metabolic precursor of NHP (Hartmann *et al.*, 2018). Previous studies have reported that inoculation with a pathogen markedly induces the accumulation of Pip in different angiosperm species, including Arabidopsis, tobacco, potato, soybean, rice, and barley (Pálfi and Dézsi, 1968; Návarová *et al.*, 2012; Vogel-Adzhough *et al.*, 2013; Aliferis *et al.*, 2014; Abeysekara *et al.*, 2016; Adam *et al.*, 2018; Lenk *et al.*, 2019). Consistently, the results of the present study show that the levels of Pip increase strongly in leaves of tobacco, soybean, and barley inoculated with compatible *Pseudomonas* bacteria (Figs 4, 5). In addition, Pip accumulated to high levels in tomato leaves infected with bacterial and oomycete pathogens (Fig. 4B, C). While most of the species investigated have low basal levels of Pip and show early and strong pathogen-induced accumulation, the situation is different for *C. sativus* and *B. distachyon*. In cucumber, the constitutive levels of Pip were remarkably high (between ~ 10 and $25 \mu\text{g g}^{-1}$ FW) and

did not increase further after *P. syringae* inoculation (Fig. 2A, B). Further, in *B. distachyon*, relatively high basal levels of Pip were detected ($1\text{--}2\ \mu\text{g}^{-1}\ \text{FW}$), and the increases in Pip upon pathogen inoculation were slower than the induced accumulation of its derivative NHP (Fig. 5A, B).

Therefore, different mechanisms for the induction of NHP biosynthesis seem to exist in different species. In Arabidopsis, tobacco, tomato, soybean, and barley, NHP accumulates in response to pathogen attack in parallel with its direct metabolic precursor Pip. In addition, the levels of the pathway precursor amino acid lysine increase in these plants (Supplementary Fig. S5; Návarová et al., 2012). The accumulation of Pip in leaf tissue of inoculated Arabidopsis is closely linked to a concomitant increase in the expression of the biosynthetic genes *ALD1* and *SARD4*, and the direct Pip precursor dehydropipecolic acid accumulates to detectable levels (Návarová et al., 2012; Hartmann et al., 2017; Hartmann and Zeier, 2019). The immune-regulatory genes *EDS1* and *PAD4*, as well as several transcription factors, have been shown to regulate inducible *ALD1* gene expression and the biosynthesis of Pip and NHP (Návarová et al., 2012; Hartmann et al., 2018; Sun et al., 2018, 2020; Wang et al., 2018; Kim et al., 2020). Similarly, the pathogen-induced accumulation of Pip in soybean is closely linked to a concomitant transcriptional induction of the soybean *ALD1* homolog *GmALD1* (Abeysekara et al., 2016).

Cucumber belongs to a second category. In this species, high constitutive levels of Pip, associated with constitutive expression of the *C. sativus* *ALD1* gene, were observed, and the levels of lysine remained unchanged after pathogen inoculation (Fig. 2; Supplementary Figs S2, S5). Although induction of the pathway after *P. syringae* inoculation was associated with further increases in the level of *ALD1* transcripts, the *de novo* expression of *FMO1* after pathogen infection appears to be the major pathogen-inducible trigger that switches on the generation of NHP (Supplementary Fig. S2). In this way, endogenous Pip could be used directly as a substrate as *FMO1* accumulates to rapidly induce NHP biosynthesis. The metabolic situation in cucumber corroborates our previous findings that NHP, but not its metabolic precursor Pip, is the actual mediator of resistance, because otherwise, the elevated constitutive levels of Pip would result in constitutive immunity in *C. sativus*. Consistently, Arabidopsis plants defective in the Pip *N*-hydroxylase gene *FMO1* were unable to induce SAR in response to exogenous Pip, but acquired resistance when supplied with exogenous NHP (Hartmann et al., 2018). The situation in *B. distachyon* appears to be intermediate: although the whole metabolic pathway is induced upon pathogen attack, an early accumulation of NHP might be favored by the relatively high constitutive levels of the Pip precursor (Fig. 5A, B; Supplementary Fig. S2).

Our present and previous results show that NHP is able to induce acquired resistance in distinct angiosperm species. On the one hand, genetic and biochemical evidence demonstrated that the endogenous accumulation of NHP is required for pathogen-induced SAR in Arabidopsis. On the other hand, exogenously applied NHP induced SAR to *P. syringae* and *H. arabidopsidis* infection in Arabidopsis and restored the defects in acquired resistance of the NHP-deficient *ald1* and *fmo1* mutants (Hartmann et al., 2018; Chen et al., 2018). Although

the cellular compartment(s) in which NHP acts to trigger SAR and what concentration of NHP is necessary for biological activity in target cells remain elusive, our feeding experiments indicate that effective doses of exogenously applied NHP per plant below $1\ \mu\text{mol}$ are sufficient to activate SAR (Fig. 1). In addition to our results, Holmes et al. (2019) recently reported that exogenously applied NHP induces SAR in tomato and pepper (*Capsicum annuum*) to *Xanthomonas euvesicatoria* and *Pst* DC3000 infection, respectively. Here, we show that *C. sativus* and tobacco plants exogenously treated with NHP acquire resistance to *P. syringae* infection (Fig. 3). In addition to these dicots, the monocot *B. distachyon* acquires resistance to both *X. translucens* and *M. oryzae* infection when supplied with NHP (Fig. 6). Together, these findings indicate that NHP functions as a general and conserved regulator of SAR in angiosperms. NHP mediates the resistance of both mono- and dicotyledonous plant species against infection with bacterial, oomycete, and fungal pathogens. Such an activation of broad-spectrum plant resistance has long been considered a hallmark of SAR (Sticher et al., 1997).

Plants with activated SAR are primed for a fortified induction of defenses when attacked by pathogens (Jung et al., 2009; Návarová et al., 2012). Defense priming in Arabidopsis is completely absent in the NHP-deficient *ald1* and *fmo1* mutants, indicating a central role for NHP in this process (Návarová et al., 2012; Bernsdorff et al., 2016). NHP-induced resistance and SAR-related defense priming are substantially diminished in the SA-deficient *sid2* mutant (Bernsdorff et al., 2016; Hartmann et al., 2018). Moreover, Pip primed plants for enhanced SA-induced expression of *PR* genes in an *FMO1*-dependent manner (Bernsdorff et al., 2016). These observations suggest a close, coordinated interplay between NHP and SA for the effective activation of SAR and defense priming (Hartmann and Zeier, 2019). Our metabolite analyses show that in the leaves of pathogen-inoculated Arabidopsis, cucumber, tobacco, tomato, soybean, and *B. distachyon*, SA (and, commonly, SA derivatives such as SAG) accumulates in parallel with NHP, so that, in principle, such a positive interplay for effective SAR induction is possible in these species.

Exceptionally, although barley leaves accumulated Pip and NHP upon *M. oryzae* infection, SA levels were low and did not change upon inoculation (Fig. 5C). Similarly, SA did not accumulate in *H. vulgare* infected with the powdery mildew pathogen *Blumeria graminis* f. sp. *hordei* (Hückelhoven et al., 1999). Therefore, SA might not function as a signaling molecule for SAR in barley. In line with this assumption, a previous study has found that *H. vulgare* does not show enhanced pathogen resistance in response to exogenous SA treatments (Dey et al., 2014), which is a common feature in many other species. However, barley plants can acquire resistance upon treatment with the synthetic resistance activator 2,6-dichloroisonicotinic acid (Kogel et al., 1994). Our metabolite analyses suggest that an endogenous activation of SAR in barley might proceed via NHP signaling. This assumption is compatible with the finding that *H. vulgare* plants fed with exogenous Pip acquired resistance to *X. translucens* infection (Lenk et al., 2019).

The molecular mechanisms for SAR in monocots are not as well understood as those in dicots (Shah and Zeier, 2013;

Balmer *et al.*, 2013). Our findings that *B. distachyon* shows accumulation of Pip and NHP upon inoculation with both *X. translucens* and *M. oryzae*, and that NHP treatment induces acquired resistance in *B. distachyon* to infection by the same pathogens, strongly suggest that NHP functions as a central player in SAR in this species (Figs 5, 6). The capacities established for *B. distachyon* as a model monocot hold the promise of further investigations of this aspect at the genetic level (Fitzgerald *et al.*, 2015; Scholthoff *et al.*, 2018). Overexpression of the first NHP pathway gene, *ALD1*, in rice also enhanced resistance to infection by *M. oryzae* (Jung *et al.*, 2016), which might indicate a function of NHP for SAR in rice as well. Together, the existing data on *B. distachyon*, barley, and rice suggest that activation of SAR by NHP might be a conserved feature of monocotyledonous plants. Our metabolic data further show that SA strongly accumulates in *B. distachyon* leaves upon *X. translucens* infection, while SA levels were only moderately increased in response to *M. oryzae* inoculation (Fig. 5). Therefore, SA might well contribute to SAR in *B. distachyon*. In this light, a general separation of the molecular bases of SAR into “monocot” and “dicot” mechanisms might not be helpful. Rather, the degree of induction of different SAR-activating pathways appears to vary within distinct species and depends on the nature of the attacking pathogen.

An interesting—and controversial—mechanistic aspect of SAR is how plants achieve long-distance communication between inoculated and distant leaves (Shah and Zeier, 2013). When NHP was locally applied to individual rosette leaves of Arabidopsis by infiltration, it induced defense-related gene expression and a strong SAR response in distant leaves (Fig. 1; Chen *et al.*, 2018). Similarly, local application of NHP to leaflets proximal to the stem induced systemic resistance in distal leaflets of the same tomato leaf (Holmes *et al.*, 2019). Moreover, transient *Agrobacterium*-mediated expression of the Arabidopsis NHP biosynthetic genes *ALD1*, *SARD4*, and *FMO1* in *Nicotiana benthamiana* led to an accumulation of NHP in the absence of pathogen inoculation. When the same set of genes was transiently expressed in proximal leaflets of tomato leaves, systemic immunity was induced in distal leaflets (Holmes *et al.*, 2019). These results support a possible function of NHP as a mobile SAR signal, but the experimental setups do not clarify whether the NHP that accumulates biologically during a SAR-inducing plant–pathogen interaction in inoculated leaves ultimately travels to distant leaves and induces SAR. On the basis of these results, it is also possible that NHP triggers a long-distance communication process in the inoculated tissue that is based on cell-to-cell signal propagation.

Previous efforts to identify SAR-related mobile signaling compounds investigated the changes in the phloem sap of cucumber plants following SAR induction by pathogen infection. Upon inoculation of the first leaf of cucumber with necrotizing pathogens, Métraux *et al.* (1990) observed increases in the levels of SA in collected phloem sap that preceded the induction of resistance in the second (systemic) leaf. Studies using putative radiolabeled biosynthetic precursors of SA concluded that SA is transportable from lower inoculated to upper cucumber leaves, but explicitly pointed out the potential existence of a different primary signal that

induces *de novo* synthesis of SA in upper leaves (Meuwly *et al.*, 1995). In line with the latter suggestion, systemic defense responses in distal cucumber leaves were observed before increases in SA levels were detectable in the phloem sap of inoculated leaves (Rasmussen *et al.*, 1991). In addition to SA, its structural isomer, 4-hydroxybenzoic acid, was found to accumulate in the phloem sap from inoculated leaves of cucumber, but its relation to SAR remained unclear (Smith-Becker *et al.*, 1998). Using *Psl* inoculation of the first leaf of cucumber, we confirmed that SA and its derivative, SAG, strongly accumulate in the inoculated leaves and the phloem sap of these leaves, but only SAG accumulated substantially in the exudates of systemic leaves (Fig. 2; Supplementary Fig. S1). By contrast, the levels of NHP increased strongly in the inoculated leaves and their phloem sap, in the phloem sap of distant leaves and, finally, in the distant leaf tissue itself (Fig. 2). These findings support the hypothesis that NHP might function as a mobile primary inducer of SAR within a biological plant–pathogen interaction. Its biosynthetic precursor Pip, which was previously identified as a constituent of the phloem in cucumber (Hu *et al.*, 2016), occurs at very high constitutive levels in both cucumber leaves and exudates and does not increase in response to the pathogen stimulus (Fig. 2). This observation argues against a function of Pip as a mobile SAR signal, but indicates that Pip could be a ready-to-use precursor in cucumber leaf and phloem tissue that is converted into the primary, and presumably phloem-mobile, SAR inducer NHP in response to pathogen attack.

Conclusions

The current study shows that the biosynthesis of NHP is induced in diverse monocot and dicot plants species upon inoculation with compatible pathogens. In addition, NHP functions as a potent activator of acquired resistance in both mono- and dicotyledonous plants. The systemic accumulation of NHP in leaves and phloem sap of inoculated cucumber plants is consistent with a possible function of NHP as a primary, mobile SAR-inducer.

Supplementary data

Supplementary data are available at *JXB* online.

Fig. S1. Levels of glycosylated salicylic acid and N-hydroxyphenylpicolinic acid conjugates in leaves and phloem sap of *Cucumis sativus*.

Fig. S2. Transcript levels of *Cucumis sativus* *ALD1* and *FMO1* in *P. syringae* pv. *lachrymans*-inoculated and mock-treated leaves.

Fig. S3. Exogenous application of Pip and NHP elevate resistance of tobacco to *P. syringae* pv. *tabaci* infection.

Fig. S4. Levels of SA- β -glucoside and SA glucose ester in leaves of *Brachypodium distachyon*.

Fig. S5. Levels of lysine in pathogen-inoculated leaves of different plant species.

Table S1. Plant growth conditions.

Table S2. Primers and conditions used for RT-PCR analysis.

Acknowledgements

We thank Grzegorz Bartoszewski and Nick Talbot for kindly providing *Pseudomonas syringae* pv. *lachrymans* strain 814/98 and *Magnaporthe oryzae* strain Guy11, respectively. This work was funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) via a grant to JZ (ZE467/6-2) and Germany's Excellence Strategy – EXC-2048/1 (project ID 390686111).

References

- Abeysekera NS, Swaminathan S, Desai N, Guo L, Bhattacharyya MK.** 2016. The plant immunity inducer pipecolic acid accumulates in the xylem sap and leaves of soybean seedlings following *Fusarium virguliforme* infection. *Plant Science* **243**, 105–114.
- Ádám AL, Nagy ZÁ, Kátay G, Mergenthaler E, Viczián O.** 2018. Signals of systemic immunity in plants: progress and open questions. *International Journal of Molecular Sciences* **19**, E1146.
- Ahuja I, Kissen R, Bones AM.** 2012. Phytoalexins in defense against pathogens. *Trends in Plant Science* **17**, 73–90.
- Aliferis KA, Faubert D, Jabaji S.** 2014. A metabolic profiling strategy for the dissection of plant defense against fungal pathogens. *PLoS One* **9**, e111930.
- Balmer D, Planchamp C, Mauch-Mani B.** 2013. On the move: induced resistance in monocots. *Journal of Experimental Botany* **64**, 1249–1261.
- Berens ML, Berry HM, Mine A, Argueso CT, Tsuda K.** 2017. Evolution of hormone signaling networks in plant defense. *Annual Review of Phytopathology* **55**, 401–425.
- Bernsdorff F, Döring AC, Gruner K, Schuck S, Bräutigam A, Zeier J.** 2016. Pipecolic acid orchestrates plant systemic acquired resistance and defense priming via salicylic acid-dependent and -independent pathways. *The Plant Cell* **28**, 102–129.
- Broquist HP.** 1991. Lysine-pipecolic acid metabolic relationships in microbes and mammals. *Annual Review of Nutrition* **11**, 435–448.
- Chen YC, Holmes EC, Rajniak J, Kim JG, Tang S, Fischer CR, Mudgett MB, Sattely ES.** 2018. *N*-hydroxy-pipecolic acid is a mobile metabolite that induces systemic disease resistance in *Arabidopsis*. *Proceedings of the National Academy of Sciences, USA* **115**, E4920–E4929.
- Chezem WR, Memon A, Li FS, Weng JK, Clay NK.** 2017. SG2-type R2R3-MYB transcription factor MYB15 controls defense-induced lignification and basal immunity in *Arabidopsis*. *The Plant Cell* **29**, 1907–1926.
- Delventhal R, Rajaraman J, Stefanato FL, et al.** 2017. A comparative analysis of nonhost resistance across the two Triticaceae crop species wheat and barley. *BMC Plant Biology* **17**, 232.
- de Vries S, von Dahlen JK, Uhlmann C, Schnake A, Kloesges T, Rose LE.** 2017. Signatures of selection and host-adapted gene expression of the *Phytophthora infestans* RNA silencing suppressor PSR2. *Molecular Plant Pathology* **18**, 110–124.
- Dey S, Wenig M, Langen G, et al.** 2014. Bacteria-triggered systemic immunity in barley is associated with WRKY and ETHYLENE RESPONSIVE FACTORS but not with salicylic acid. *Plant Physiology* **166**, 2133–2151.
- Ding P, Rekhter D, Ding Y, et al.** 2016. Characterization of a pipecolic acid biosynthesis pathway required for systemic acquired resistance. *The Plant Cell* **28**, 2603–2615.
- Fitzgerald TL, Powell JJ, Schneebeil K, et al.** 2015. *Brachypodium* as an emerging model for cereal–pathogen interactions. *Annals of Botany* **115**, 717–731.
- Fu ZQ, Dong X.** 2013. Systemic acquired resistance: turning local infection into global defense. *Annual Review of Plant Biology* **64**, 839–863.
- Gaffney T, Friedrich L, Vernooij B, Negrotto D, Nye G, Uknes S, Ward E, Kessmann H, Ryals J.** 1993. Requirement of salicylic acid for the induction of systemic acquired resistance. *Science* **261**, 754–756.
- Guerra T, Schilling S, Hake K, Gorzolk K, Sylvester FP, Conrads B, Westermann B, Romeis T.** 2020. Calcium-dependent protein kinase 5 links calcium signaling with *N*-hydroxy-L-pipecolic acid- and *SARD1*-dependent immune memory in systemic acquired resistance. *New Phytologist* **225**, 310–325.
- Hartmann M, Kim D, Bernsdorff F, Ajami-Rashidi Z, Scholten N, Schreiber S, Zeier T, Schuck S, Reichel-Deland V, Zeier J.** 2017. Biochemical principles and functional aspects of pipecolic acid biosynthesis in plant immunity. *Plant Physiology* **174**, 124–153.
- Hartmann M, Zeier J.** 2018. L-lysine metabolism to *N*-hydroxypipecolic acid: an integral immune-activating pathway in plants. *The Plant Journal* **96**, 5–21.
- Hartmann M, Zeier J.** 2019. *N*-hydroxypipecolic acid and salicylic acid: a metabolic duo for systemic acquired resistance. *Current Opinion in Plant Biology* **50**, 44–57.
- Hartmann M, Zeier T, Bernsdorff F, et al.** 2018. Flavin monooxygenase-generated *N*-hydroxypipecolic acid is a critical element of plant systemic immunity. *Cell* **173**, 456–469.e16.
- Holmes EC, Chen Y-C, Sattely ES, Mudgett MB.** 2019. An engineered pathway for *N*-hydroxy-pipecolic acid synthesis enhances systemic acquired resistance in tomato. *Science Signaling* **12**, eaay3066.
- Hu C, Ham BK, El-Shabrawi HM, Alexander D, Zhang D, Ryals J, Lucas WJ.** 2016. Proteomics and metabolomics analyses reveal the cucurbit sieve tube system as a complex metabolic space. *The Plant Journal* **87**, 442–454.
- Hückelhoven R, Fodor J, Preis C, Kogel KH.** 1999. Hypersensitive cell death and papilla formation in barley attacked by the powdery mildew fungus are associated with hydrogen peroxide but not with salicylic acid accumulation. *Plant Physiology* **119**, 1251–1260.
- Jung GY, Park JY, Choi HJ, Yoo SJ, Park JK, Jung HW.** 2016. A rice gene homologous to *Arabidopsis* AGD2-LIKE DEFENSE1 participates in disease resistance response against infection with *Magnaporthe oryzae*. *The Plant Pathology Journal* **32**, 357–362.
- Jung HW, Tschaplinski TJ, Wang L, Glazebrook J, Greenberg JT.** 2009. Priming in systemic plant immunity. *Science* **324**, 89–91.
- Kim Y, Gilmour SJ, Chao L, Park S, Thomashow MF.** 2020. *Arabidopsis* CAMTA transcription factors regulate pipecolic acid biosynthesis and priming of immunity genes. *Molecular Plant* **13**, 157–168.
- Klessig DF, Choi HW, Dempsey DA.** 2018. Systemic acquired resistance and salicylic acid: past, present, and future. *Molecular Plant-Microbe Interactions* **31**, 871–888.
- Kogel KH, Beckhove U, Dreschers J, Munch S, Romme Y.** 1994. Acquired resistance in barley (The resistance mechanism induced by 2,6-dichloroisonicotinic acid is a phenocopy of a genetically based mechanism governing race-specific powdery mildew resistance). *Plant Physiology* **106**, 1269–1277.
- Lenk M, Wenig M, Bauer K, et al.** 2019. Pipecolic acid is induced in barley upon infection and triggers immune responses associated with elevated nitric oxide accumulation. *Molecular Plant-Microbe Interactions* **32**, 1303–1313.
- Métraux JP, Signer H, Ryals J, Ward E, Wyss-Benz M, Gaudin J, Raschdorf K, Schmid E, Blum W, Inverardi B.** 1990. Increase in salicylic acid at the onset of systemic acquired resistance in cucumber. *Science* **250**, 1004–1006.
- Meuwly P, Molders W, Buchala A, Métraux JP.** 1995. Local and systemic biosynthesis of salicylic acid in infected cucumber plants. *Plant Physiology* **109**, 1107–1114.
- Mishina TE, Zeier J.** 2006. The *Arabidopsis* flavin-dependent monooxygenase FMO1 is an essential component of biologically induced systemic acquired resistance. *Plant Physiology* **141**, 1666–1675.
- Mishina TE, Zeier J.** 2007. Pathogen-associated molecular pattern recognition rather than development of tissue necrosis contributes to bacterial induction of systemic acquired resistance in *Arabidopsis*. *The Plant Journal* **50**, 500–513.
- Mölders W, Meuwly P, Summermatter C, Métraux J-P.** 1994. Salicylic acid content in cucumber plants infected with *Pseudomonas lachrymans* and tobacco necrosis virus. *Acta Horticulturae* **381**, 375–378.
- Návarová H, Bernsdorff F, Döring AC, Zeier J.** 2012. Pipecolic acid, an endogenous mediator of defense amplification and priming, is a critical regulator of inducible plant immunity. *The Plant Cell* **24**, 5123–5141.
- Nawrath C, Métraux JP.** 1999. Salicylic acid induction-deficient mutants of *Arabidopsis* express *PR-2* and *PR-5* and accumulate high levels of camalexin after pathogen inoculation. *The Plant Cell* **11**, 1393–1404.
- Pálfi G, Dészi L.** 1968. Pipecolic acid as an indicator of abnormal protein metabolism in diseased plants. *Plant and Soil* **29**, 285–291.

- Parker D, Beckmann M, Enot DP, Overy DP, Rios ZC, Gilbert M, Talbot N, Draper J.** 2008. Rice blast infection of *Brachypodium distachyon* as a model system to study dynamic host/pathogen interactions. *Nature Protocols* **3**, 435–445.
- Parker D, Beckmann M, Zubair H, Enot DP, Caracuel-Rios Z, Overy DP, Snowdon S, Talbot NJ, Draper J.** 2009. Metabolomic analysis reveals a common pattern of metabolic re-programming during invasion of three host plant species by *Magnaporthe oryzae*. *The Plant Journal* **59**, 723–737.
- Rasmussen JB, Hammerschmidt R, Zook MN.** 1991. Systemic induction of salicylic acid accumulation in cucumber after inoculation with *Pseudomonas syringae* pv *syringae*. *Plant Physiology* **97**, 1342–1347.
- Scholthof KBG, Irigoyen S, Catalan P, Mandadi KK.** 2018. *Brachypodium*: a monocot grass model genus for plant biology. *The Plant Cell* **30**, 1673–1694.
- Shah J, Zeier J.** 2013. Long-distance communication and signal amplification in systemic acquired resistance. *Frontiers in Plant Science* **4**, 30.
- Smith-Becker J, Marois E, Huguet EJ, Midland SL, Sims JJ, Keen NT.** 1998. Accumulation of salicylic acid and 4-hydroxybenzoic acid in phloem fluids of cucumber during systemic acquired resistance is preceded by a transient increase in phenylalanine ammonia-lyase activity in petioles and stems. *Plant Physiology* **116**, 231–238.
- Ślomska R, Olczak-Woltman H, Korzeniewska A, Gozdowski D, Niemirowicz-Szczytt K, Bartoszewski G.** 2018. Genetic mapping of *psl* locus and quantitative trait loci for angular leaf spot resistance in cucumber (*Cucumis sativus* L.). *Molecular Breeding* **38**, 111.
- Stahl E, Hartmann M, Scholten N, Zeier J.** 2019. A role for tocopherol biosynthesis in Arabidopsis basal immunity to bacterial infection. *Plant Physiology* **181**, 1008–1028.
- Sticher L, Mauch-Mani B, Métraux JP.** 1997. Systemic acquired resistance. *Annual Review of Phytopathology* **35**, 235–270.
- Sun T, Busta L, Zhang Q, Ding P, Jetter R, Zhang Y.** 2018. TGACG-BINDING FACTOR 1 (TGA1) and TGA4 regulate salicylic acid and pipecolic acid biosynthesis by modulating the expression of *SYSTEMIC ACQUIRED RESISTANCE DEFICIENT 1* (*SARD1*) and *CALMODULIN-BINDING PROTEIN 60g* (*CBP60g*). *New Phytologist* **217**, 344–354.
- Sun T, Huang J, Xu Y, et al.** 2020. Redundant CAMTA transcription factors negatively regulate the biosynthesis of salicylic acid and N-hydroxypipecolic acid by modulating the expression of *SARD1* and *CBP60g*. *Molecular Plant* **13**, 144–156.
- Vernooij B, Friedrich L, Morse A, Reist R, Kolditz-Jawhar R, Ward E, Uknes S, Kessmann H, Ryals J.** 1994. Salicylic acid is not the translocated signal responsible for inducing systemic acquired resistance but is required in signal transduction. *The Plant Cell* **6**, 959–965.
- Vogel-Adzhoghov D, Stahl E, Návarová H, Zeier J.** 2013. Pipecolic acid enhances resistance to bacterial infection and primes salicylic acid and nicotine accumulation in tobacco. *Plant Signaling & Behavior* **8**, e26366.
- Wang Y, Schuck S, Wu J, Yang P, Döring AC, Zeier J, Tsuda K.** 2018. A MPK3/6-WRKY33-ALD1-pipecolic acid regulatory loop contributes to systemic acquired resistance. *The Plant Cell* **30**, 2480–2494.
- Wildermuth MC, Dewdney J, Wu G, Ausubel FM.** 2001. Isochorismate synthase is required to synthesize salicylic acid for plant defence. *Nature* **414**, 562–565.
- Xu B, Fan Z, Lei Y, Ping Y, Jaisi A, Xiao Y.** 2018. Insights into pipecolic acid biosynthesis in *Huperzia serrata*. *Organic Letters* **20**, 2195–2198.
- Zeier J.** 2013. New insights into the regulation of plant immunity by amino acid metabolic pathways. *Plant, Cell & Environment* **36**, 2085–2103.