

BRIEF REPORT

Open Access



Intracranial hemorrhage in COVID-19 patients during extracorporeal membrane oxygenation for acute respiratory failure: a nationwide register study report

Saskia von Stillfried¹, Roman David Bülow¹, Rainer Röhrig², Patrick Meybohm^{3*†}, Peter Boor^{1*†} and for the German Registry of COVID-19 Autopsies (DeRegCOVID), DeRegCOVID Collaborators#

Abstract

Background: In severe cases, SARS-CoV-2 infection leads to acute respiratory distress syndrome (ARDS), often treated by extracorporeal membrane oxygenation (ECMO). During ECMO therapy, anticoagulation is crucial to prevent device-associated thrombosis and device failure, however, it is associated with bleeding complications. In COVID-19, additional pathologies, such as endotheliitis, may further increase the risk of bleeding complications. To assess the frequency of bleeding events, we analyzed data from the German COVID-19 autopsy registry (DeRegCOVID).

Methods: The electronic registry uses a web-based electronic case report form. In November 2021, the registry included $N = 1129$ confirmed COVID-19 autopsy cases, with data on 63 ECMO autopsy cases and 1066 non-ECMO autopsy cases, contributed from 29 German sites.

Findings: The registry data showed that ECMO was used in younger male patients and bleeding events occurred much more frequently in ECMO cases compared to non-ECMO cases (56% and 9%, respectively). Similarly, intracranial bleeding (ICB) was documented in 21% of ECMO cases and 3% of non-ECMO cases and was classified as the immediate or underlying cause of death in 78% of ECMO cases and 37% of non-ECMO cases. In ECMO cases, the three most common immediate causes of death were multi-organ failure, ARDS and ICB, and in non-ECMO cases ARDS, multi-organ failure and pulmonary bacterial $\pm$ fungal superinfection, ordered by descending frequency.

Interpretation: Our study suggests the potential value of autopsies and a joint interdisciplinary multicenter (national) approach in addressing fatal complications in COVID-19.

Keywords: Autopsy, Registry, COVID-19, ECMO, Intracranial bleeding, Bleeding events

Introduction

Venovenous extracorporeal membrane oxygenation (VV-ECMO) is used for refractory severe acute respiratory failure with a survival rate of about 50% [1]. The interface between blood and non-biological ECMO circuit elements requires therapeutic anticoagulation, predisposing patients to an increased risk of bleeding.

Based on a recent analysis from the Extracorporeal Life Support Organization (ELSO) registry including

*Correspondence: meybohm_p@ukw.de; pboor@ukaachen.de

[†]Patrick Meybohm and Peter Boor have contributed equally to this work

¹ Institute of Pathology, University Hospital RWTH Aachen, Pauwelsstraße 30, 52074 Aachen, Germany

³ Department of Anaesthesiology, Intensive Care, Emergency and Pain Medicine, University Hospital Würzburg, Oberdürrbacher Str. 6, 97080 Würzburg, Germany

Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

7579 patients from 2007 to 2018, 37% experienced any bleeding event, and 21.2% experienced bleeding combined with a thrombotic event. While the most common bleeding events with cannulation (15.5%) and surgical site (9.6%) bleeding are easy to handle, intracranial hemorrhage occurred in only 4.5% and has been consistently associated with poor survival [2]. Other retrospective studies reported intracranial hemorrhage in 6–35.4% of patients during ECMO therapy [3–5].

Causes of intracranial hemorrhage in ECMO patients may be associated with heparin overdose, circuit-associated defibrination, thrombocytopenia, disseminated intravascular coagulation, acquired von Willebrand syndrome, and also COVID-19-associated endotheliitis. Importantly, intracranial hemorrhage was observed in many patients receiving VV-ECMO without coagulopathy or anticoagulant use [6].

In this respect, we analyzed preliminary data from the German COVID-19 Autopsy Registry [7], involving 63 deceased COVID-19 patients who received ECMO for acute respiratory failure. Specifically, we address the following questions, comparing COVID-19 autopsy cases that received ECMO support with cases that did not:

1. What differences in demographical characteristics exist?
2. What are the prevalences of bleeding events, specifically intracranial bleeding events found at autopsy?

Patient inclusion, data acquisition and management, and cohort stratification were performed as previously described [8] and are provided in the Additional file 1.

For analyses, $N=1129$ autopsy cases with positive COVID-19 test (preclinical, clinical, or post-mortem, point of care antigen test from nasopharyngeal or pulmonary swabs or PCR test from nasopharyngeal or pulmonary swabs or tissue samples) were eligible in the German COVID-19 Autopsy Registry (DeRegCOVID). 20–22% of the COVID-19 autopsy cases were located in the East and West, respectively and 28–30% in the North

and South of Germany, respectively by patient residential region (Fig. 1a) or by the autopsy center region (Fig. 1b), respectively.

The percentage of COVID-19 ECMO autopsy cases of all autopsied cases increased with each pandemic wave, from 2% in the first pandemic wave to 6% in the second and 12% in the third wave (Fig. 1c).

$N=63$ patients underwent ECMO therapy. The male to female ratio in this cohort was 5:1 with a homogeneous distribution of females over the different age range of <30–80 years, and a male peak in the age range of 51–70 years (Fig. 1d). Of the remaining $N=1065$ COVID-19 non-ECMO autopsy cases (one value missing), the male to female ratio was 2:1, with male peaks at 61–90 years and female predominance at >90 years of age (Fig. 1e). The percentage of COVID-19 ECMO autopsy cases in the respective sex/age group was higher for males compared to females, however, in the younger age groups, total numbers were relatively low, resulting in large effects of single cases on the percentage (Fig. 1f).

Any kind of a bleeding event was found in 56% of ECMO cases ($N=35$ cases) and 9% of non-ECMO cases ($N=100$ cases, p value < 0.0001). Intracranial bleeding (ICB) was documented in $N=13$ ECMO cases (21%) and in $N=30$ non-ECMO cases (3%, p value < 0.0001, Fig. 1g). In ECMO patients with ICB, in three cases ($N=2$ soft tissue bleeding due to cannulation and $N=1$ epistaxis) and in non-ECMO patients with ICB, in three cases extracranial bleeding events were documented, respectively ($N=2$ acute posthemorrhagic anemia not otherwise specified and $N=1$ recurrent bleeding of the lower gastrointestinal tract, a detailed specification of bleeding events is provided in Fig. 1h). In 78% of ECMO cases and 37% of non-ECMO cases with ICB, the intracranial bleeding was classified as the immediate or underlying cause of death. The five most common immediate causes of death were multi-organ failure, DAD/ARDS, ICB, pulmonary bacterial $\pm$ fungal superinfection and extracranial bleeding events in $N=63$ ECMO cases and DAD/ARDS, multi-organ failure, pulmonary bacterial $\pm$ fungal

(See figure on next page.)

Fig. 1 **a** Number of COVID-19 autopsy cases and percentage of COVID-19 autopsies after ECMO therapy by postal code of the deceased person (1 value missing of ECMO cases, 17 values missing of non-ECMO cases). **b** Number of COVID-19 autopsy cases and percentage of COVID-19 autopsies after ECMO therapy by postal code of the contributing center. **c** Individual disease duration (orange bars) or death date (black boxes, when no data on symptom onset/ first positive SARS-CoV-2 test was available) in $N=63$ ECMO COVID-19 autopsy cases. **d** Age and sex distribution in COVID-19 autopsies after ECMO therapy ($N=63$). **e** Age and sex distribution in COVID-19 autopsies without ECMO therapy ($N=1065$, 1 value missing). **f** Age and sex distribution in COVID-19 autopsies as a percentage of respective age group. **g** Intracranial bleeding (ICB) and other hemorrhages in ECMO and non-ECMO COVID-19 cases. The associations between the variables ECMO and ICB and ECMO and any bleeding event were significant (both p value < 0.0001 Fisher's exact test, two-tailed). Note that the number of bleeding events exceeds the number of patients, because in $N=3$ non-ECMO, and $N=3$ ECMO autopsies, both ICB and other bleeding events were present at the autopsy, respectively. **h** ECMO cases (violet) and non-ECMO cases (dark yellow) with any bleeding event. The number of extracranial bleeding events is higher compared to **h**, because, in $N=4$ ECMO cases, two different extracranial bleeding events were documented. ICB, intracranial bleeding




Fig. 1 (See legend on previous page.)

superinfection, pulmonary embolism, and ischemic heart disease in $N=1031$ non-ECMO cases with the complete cause of death data (ordered by descending frequency).

VV-ECMO used in refractory severe acute respiratory failure is associated with an increased risk of bleeding, of which intracranial hemorrhage has been consistently associated with very poor survival. In this report, we analyzed data from the German Registry of COVID-19 Autopsies (DeRegCOVID) to gain further insights into COVID-19 ECMO autopsy cases with bleeding events in comparison to non-ECMO cases.

ECMO being more often documented in younger and male patients, is in line with data from the Extracorporeal Life Support Organization (ELSO) Registry [3].

The prevalence of any bleeding event in more than 50% of COVID-19 ECMO autopsy cases is higher compared to a previous multicenter observational study of 152 consecutive non-autopsy patients with severe COVID-19 supported by ECMO in four UK commissioned centers during the first wave of the COVID-19 pandemic (30.9% major bleedings) [9]. This might be explained by our cohort consisting of fatal cases, which may lead to an overrepresentation of cases with bleeding events. Also, all macroscopically identified bleeding events are documented, irrespective of major bleeding criteria [10], as these data are usually not available at autopsy. Our findings regarding the prevalence of intracranial bleeding and associated mortality are consistent with a report from three tertiary care ECMO centers in Germany and Switzerland [11]. In an observational study from Northern Germany, the observation of intracranial bleeding in COVID-19 non-autopsy ECMO patients (35.4%) was higher compared to our findings. In a study from a single tertiary center on 25 non-COVID ECMO autopsy cases, 52% had intracranial macrohemorrhages [12]. However, it is possible, that due to concerns regarding occupational hazards, the omission of brain examination especially during the first pandemic wave led to an underreporting of intracranial hemorrhage at autopsy in our cohort.

Limitations

The registry only gathers data available to pathologists at the time of autopsy. The clinical information provided during autopsies is usually comprehensive, particularly regarding treatment approaches such as ECMO. Still, we cannot exclude missing data in the registry on potential ECMO therapy. Another limitation is the missing specific data on invasive ventilation therapy and missing reliable data on the mode and time of anticoagulant therapy in our cohort. Because we aimed at the broadest possible participation in the registry, the eCRF does not cover therapy and intensive care details and duration of

ventilation or ECMO therapy. Considering that in more than 50% of our cohort, the immediate cause of death at autopsy was COVID-19 DAD/ARDS, it is likely that some of these patients underwent invasive ventilation therapy before death in hospitalized cases [8].

In conclusion, our report showed autopsy-confirmed increased prevalence of bleeding events and intracranial hemorrhages as causes of death in COVID-19 patients with ECMO treatment, compared to those without ECMO treatment. This illustrates the value of autopsies and a joint interdisciplinary multicenter (national) approach in addressing fatal complications in intensive care.

Abbreviations

ARDS: Acute respiratory distress syndrome; COVID-19: Coronavirus disease 2019; DAD: Diffuse alveolar damage; DeRegCOVID: German COVID-19 Autopsy Registry; eCRF: Electronic case report form; ELSO: Extracorporeal Life Support Organization; ICB: Intracranial bleeding; PCR: Polymerase chain reaction; RWTH Aachen University: Rhenish-Westphalian Technical University Aachen; SARS-CoV-2: Severe acute respiratory syndrome coronavirus 2; VV-ECMO: Venovenous extracorporeal membrane oxygenation.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13054-022-03945-x>.

Additional file 1. Supplementary Methods and Discussion.

Acknowledgements

Not applicable.

#DeRegCOVID collaborators: Jana Böcker⁴, Jens Schmidt⁴, Pauline Tholen⁴, Raphael Majeed⁵, Jan Wienströer⁵, Joachim Weis⁶, Juliane Bremer⁶, Ruth Knüchel⁷, Anna Breitbach⁷, Claudio Cacchi⁷, Benita Freeborn⁷, Sophie Wucherpfennig⁷, Oliver Spring⁸, Georg Braun⁹, Christoph Römmele⁹, Bruno Märkl¹⁰, Rainer Claus¹⁰, Christine Dhillon¹⁰, Tina Schaller¹⁰, Eva Sipos¹⁰, Klaus Hirschbühl¹¹, Michael Wittmann¹¹, Elisabeth Kling¹², Thomas Kröncke¹³, Frank L. Heppner^{14,15,16}, Jenny Meinhardt¹⁴, Helena Radbruch¹⁴, Simon Streit¹⁴, David Horst¹⁷, Sefer Elezkurtaj¹⁷, Alexander Quaas¹⁸, Heike Göbel¹⁸, Torsten Hansen¹⁹, Ulf Titze¹⁹, Johann Lorenzen²⁰, Thomas Reuter²⁰, Jaroslav Woloszyn²⁰, Gustavo Baretton²¹, Julia Hilsenbeck²¹, Matthias Meinhardt²¹, Jessica Pablik²¹, Linna Sommer²¹, Olaf Holotiu²², Meike Meinel²², Nina Mahlke²³, Irene Esposito²⁴, Graziano Crudele²⁴, Maximilian Seidl²⁴, Kerstin U. Amann²⁵, Roland Coras²⁶, Arndt Hartmann²⁷, Philip Eichhorn²⁷, Florian Haller²⁷, Fabienne Lange²⁷, Kurt Werner Schmid²⁸, Marc Ingenwerth²⁸, Josefine Rawitzer²⁸, Dirk Theegarten²⁸, Christoph G. Birngruber²⁹, Peter Wild³⁰, Elise Gradhand³⁰, Kevin Smith³⁰, Martin Werner³¹, Oliver Schilling³¹, Till Acker³², Stefan Gattenlöhner³³, Christine Stadelmann³⁴, Imke Metz³⁴, Jonas Franz³⁴, Lidia Stork³⁴, Carolina Thomas³⁴, Sabrina Zechel³⁴, Philipp Ströbel³⁵, Claudia Wickenhauser³⁶, Christine Fathke³⁶, Anja Harder³⁶, Benjamin Ondruschka³⁷, Eric Dietz³⁷, Carolin Edler³⁷, Antonia Fitzek³⁷, Daniela Fröb³⁷, Axel Heinemann³⁷, Fabian Heinrich³⁷, Anke Klein³⁷, Inga Kniep³⁷, Larissa Lohner³⁷, Dustin Möbius³⁷, Klaus Püschel³⁷, Julia Schädler³⁷, Ann-Sophie Schröder³⁷, Jan-Peter Sperhake³⁷, Martin Aepfelbacher³⁸, Nicole Fischer³⁸, Marc Lütgehetmann³⁸, Susanne Pfefferle³⁸, Markus Glatzel³⁹, Susanne Krasemann³⁹, Jakob Matschke³⁹, Danny Jonigk⁴⁰, Christopher Werlein⁴⁰, Peter Schirmacher⁴¹, Lisa Maria Domke⁴¹, Laura Hartmann⁴¹, Isabel Madeleine Klein⁴¹, Constantin Schwab⁴¹, Christoph Röcken⁴², Johannes Friemann⁴³, Dorothea Langer⁴⁴, Wilfried Roth⁴⁵, Stephanie Strobl⁴⁵, Martina Rudelius⁴⁶, Konrad Friedrich Stock⁴⁷, Wilko Weichert⁴⁸, Claire Delbridge⁴⁸, Atsuko Kasajima⁴⁸, Peer-Hendrik Kuhn⁴⁸, Julia Slotta-Huspenina⁴⁸, Gregor Weirich⁴⁸, Peter Barth⁴⁹, Eva Wardelmann⁴⁹, Alexander Schnepfer⁴⁹, Katja Evert⁵⁰, Andreas Büttner⁵¹, Johannes Manhart⁵¹, Stefan Nigbur⁵¹, Iris

Bittmann⁵², Falko Fend⁵³, Hans Bösmüller⁵³, Massimo Granai⁵³, Karin Klingel⁵³, Verena Warm⁵³, Konrad Steinestel⁵⁴, Vincent Gottfried Umatham⁵⁴, Andreas Rosenwald⁵⁵, Florian Kurz⁵⁵, Niklas Vogt⁵⁵

⁴Center for Translational & Clinical Research (CTC-A), University Hospital RWTH Aachen, Aachen, Germany; ⁵Institute for Medical Informatics, University Hospital RWTH Aachen, Aachen, Germany; ⁶Institute of Neuropathology, University Hospital RWTH Aachen, Aachen, Germany; ⁷Institute of Pathology, University Hospital RWTH Aachen, Aachen, Germany; ⁸Anesthesiology and Intensive Care, University Hospital Augsburg, Augsburg, Germany; ⁹Gastroenterology, University Hospital Augsburg, Augsburg, Germany; ¹⁰General Pathology and Molecular Diagnostics, University Hospital Augsburg, Augsburg, Germany; ¹¹Hematology and Oncology, University Hospital Augsburg, Augsburg, Germany; ¹²Laboratory Medicine and Microbiology, University Hospital Augsburg, Augsburg, Germany; ¹³Radiology, University Hospital Augsburg, Augsburg, Germany; ¹⁴Department of Neuropathology, Charité - Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany; ¹⁵German Center for Neurodegenerative Diseases (DZNE) Berlin, Berlin, Germany; ¹⁶Cluster of Excellence, NeuroCure, Berlin, Germany; ¹⁷Institute of Pathology, Charité - Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany; ¹⁸Department of Pathology, University Hospital Cologne, Cologne, Germany; ¹⁹Institute of Pathology, University Hospital OWL of the Bielefeld University, Campus Lippe, Detmold, Germany; ²⁰Department of Pathology, Klinikum Dortmund, Dortmund, Germany; ²¹Institute of Pathology, University Hospital Dresden, Dresden, Germany; ²²Gemeinschaftspraxis für Pathologie, Dresden, Germany; ²³Institute of Forensic Medicine, University Hospital Düsseldorf, Düsseldorf, Germany; ²⁴Institute of Pathology, University Hospital Düsseldorf, Düsseldorf, Germany; ²⁵Department of Nephropathology, University Hospital Erlangen-Nürnberg, Erlangen, Germany; ²⁶Institute of Neuropathology, University Hospital Erlangen-Nürnberg, Erlangen, Germany; ²⁷Institute of Pathology, University Hospital Erlangen-Nürnberg, Erlangen, Germany; ²⁸Institute of Pathology, University Hospital Essen, Essen, Germany; ²⁹Institute of Forensic Medicine, University Hospital Frankfurt, Frankfurt, Germany; ³⁰Senckenberg Institute of Pathology, University Hospital Frankfurt, Frankfurt, Germany; ³¹Institute for Surgical Pathology, Medical Center-University of Freiburg, Freiburg, Germany; ³²Institute of Neuropathology, University Hospital Giessen and Marburg, Giessen, Germany; ³³Institute of Pathology, University Hospital Giessen and Marburg, Giessen, Germany; ³⁴Institute of Neuropathology, University Medical Center Göttingen, Göttingen, Germany; ³⁵Institute of Pathology, University of Göttingen, Göttingen, Germany; ³⁶Institute of Pathology, University Hospital Halle (Saale), Halle (Saale), Germany; ³⁷Institute of Legal Medicine, University Medical Center Hamburg- Eppendorf, Hamburg, Germany; ³⁸Institute of Medical Microbiology, Virology, and Hygiene, University Medical Center Hamburg- Eppendorf, Hamburg, Germany; ³⁹Institute of Neuropathology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany; ⁴⁰Institute of Pathology, Hannover Medical School, Hannover, Germany; ⁴¹Institute of Pathology, Heidelberg University Hospital, Heidelberg, Germany; ⁴²Department of Pathology, University Hospital Schleswig-Holstein, Kiel, Germany; ⁴³Department of Pathology, University Hospital Cologne, Lüdenscheid, Germany; ⁴⁴Institute of Pathology, Klinikum Magdeburg, Magdeburg, Germany; ⁴⁵Institute of Pathology, University Medical Center Mainz, Mainz, Germany; ⁴⁶Institute of Pathology, Ludwig-Maximilians-Universität München, Munich, Germany; ⁴⁷Department of Nephrology, TUM School of Medicine of Technical University of Munich, Munich, Germany; ⁴⁸Institute of Pathology, TUM School of Medicine of Technical University of Munich, Munich, Germany; ⁴⁹Gerhard Domagk Institute of Pathology, University Hospital Münster, Münster, Germany; ⁵⁰Institute of Pathology, University Hospital Regensburg, Regensburg, Germany; ⁵¹Rostock University Medical Center, University Hospital Rostock, Rostock, Germany; ⁵²Institute of Pathology, Agaplesion Diakonieklinikum Rotenburg, Rotenburg, Germany; ⁵³Institute of Pathology and Neuropathology, University Hospital Tübingen, Tübingen, Germany; ⁵⁴Department of Pathology, Bundeswehrkrankenhaus Ulm, Ulm, Germany; ⁵⁵Institute of Pathology, University of Würzburg, Würzburg, Germany

Authors' contributions

Drs SVS and PB had full access to all of the data in the study and take responsibility for the integrity of the data, which were provided by the study collaborators, and the accuracy of the data analysis. Concept and design: SVS, RDB, RR, PB. Acquisition, analysis, or interpretation of data: SVS, RDB, RR, PB. Drafting of the manuscript: SVS, RDB, RR, PB. Obtained funding: Boor. Administrative, technical, or material support: SVS, RDB, RR, PB. Supervision: PB. All authors

and collaborators approved the manuscript. All authors read and approved the final manuscript.

Funding

Open Access funding enabled and organized by Projekt DEAL. This work was supported by the German Registry of COVID-19 Autopsies (www.DeRegCOVID.ukaachen.de), funded Federal Ministry of Health (ZMVI1-2520COR201), by the Federal Ministry of Education and Research within the framework of the network of university medicine (DEFEAT PANDEMIcs, 01KX2021).

Availability of data and materials

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declarations

Ethics approval and consent to participate

Only cases with a consent given by the deceased or next of kin for autopsy or request for autopsy by the health authorities or by the prosecutor's office were included in the registry. The registry was approved by the ethical committee of the medical faculty of the RWTH Aachen University (EK 092/20). Additionally, each participating center had a local ethical approval. The registry was registered with the German Clinical Trials Registry [13].

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

⁴Center for Translational & Clinical Research (CTC-A), University Hospital RWTH Aachen, Aachen, Germany. ⁵Institute for Medical Informatics, University Hospital RWTH Aachen, Aachen, Germany. ⁶Institute of Neuropathology, University Hospital RWTH Aachen, Aachen, Germany. ⁷Institute of Pathology, University Hospital RWTH Aachen, Aachen, Germany. ⁸Anesthesiology and Intensive Care, University Hospital Augsburg, Augsburg, Germany. ⁹Department of Gastroenterology, University Hospital Augsburg, Augsburg, Germany. ¹⁰General Pathology and Molecular Diagnostics, University Hospital Augsburg, Augsburg, Germany. ¹¹Hematology and Oncology, University Hospital Augsburg, Augsburg, Germany. ¹²Laboratory Medicine and Microbiology, University Hospital Augsburg, Augsburg, Germany. ¹³Diagnostic and Interventional Radiology, University Hospital Augsburg, Augsburg, Germany. ¹⁴Department of Neuropathology, Charité - Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany. ¹⁵German Center for Neurodegenerative Diseases (DZNE) Berlin, Berlin, Germany. ¹⁶Cluster of Excellence, NeuroCure, Berlin, Germany. ¹⁷Institute of Pathology, Charité - Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany. ¹⁸Department of Pathology, University Hospital Cologne, Cologne, Germany. ¹⁹Institute of Pathology, University Hospital OWL of the Bielefeld University, Campus Lippe, Detmold, Germany. ²⁰Department of Pathology, Klinikum Dortmund, Dortmund, Germany. ²¹Institute of Pathology, University Hospital Dresden, Dresden, Germany. ²²Gemeinschaftspraxis für Pathologie, Dresden, Germany. ²³Institute of Forensic Medicine, University Hospital Düsseldorf, Düsseldorf, Germany. ²⁴Institute of Pathology, University Hospital Düsseldorf, Düsseldorf, Germany. ²⁵Department of Nephropathology, University Hospital Erlangen-Nürnberg, Erlangen, Germany. ²⁶Institute of Neuropathology, University Hospital Erlangen-Nürnberg, Erlangen, Germany. ²⁷Institute of Pathology, University Hospital Erlangen-Nürnberg, Erlangen, Germany. ²⁸Institute of Pathology, University Hospital Essen, Essen, Germany. ²⁹Institute of Forensic Medicine, University Hospital Frankfurt, Frankfurt, Germany. ³⁰Senckenberg Institute of Pathology, University Hospital Frankfurt, Frankfurt, Germany. ³¹Institute for Surgical Pathology, Medical Center-University of Freiburg, Freiburg, Germany. ³²Institute of Neuropathology, University Hospital Giessen and Marburg, Giessen, Germany. ³³Institute of Pathology, University Hospital Giessen and Marburg, Giessen, Germany. ³⁴Institute of Neuropathology, University Medical Center Göttingen, Göttingen, Germany. ³⁵Institute of Pathology, University of Göttingen, Göttingen, Germany. ³⁶Institute of Pathology, University Hospital Halle (Saale), Halle (Saale), Germany. ³⁷Institute of Legal

Medicine, University Medical Center Hamburg-Eppendorf, Hamburg, Germany. ³⁸Institute of Medical Microbiology, Virology, and Hygiene, University Medical Center Hamburg-Eppendorf, Hamburg, Germany. ³⁹Institute of Neuropathology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany. ⁴⁰Institute of Pathology, Hannover Medical School, Hannover, Germany. ⁴¹Institute of Pathology, Heidelberg University Hospital, Heidelberg, Germany. ⁴²Department of Pathology, University Hospital Schleswig-Holstein, Kiel, Germany. ⁴³Department of Pathology, University Hospital Cologne, Lüdenscheid, Germany. ⁴⁴Institute of Pathology, Klinikum Magdeburg, Magdeburg, Germany. ⁴⁵Institute of Pathology, University Medical Center Mainz, Mainz, Germany. ⁴⁶Institute of Pathology, Ludwig-Maximilians-Universität München, Munich, Germany. ⁴⁷Department of Nephrology, TUM School of Medicine of Technical University of Munich, Munich, Germany. ⁴⁸Institute of Pathology, TUM School of Medicine of Technical University of Munich, Munich, Germany. ⁴⁹Gerhard Domagk Institute of Pathology, University Hospital Münster, Münster, Germany. ⁵⁰Institute of Pathology, University Hospital Regensburg, Regensburg, Germany. ⁵¹Rostock University Medical Center, University Hospital Rostock, Rostock, Germany. ⁵²Institute of Pathology, Agaplesion Diakonieklinikum Rotenburg, Rotenburg, Germany. ⁵³Institute of Pathology and Neuropathology, University Hospital Tübingen, Tübingen, Germany. ⁵⁴Department of Pathology, Bundeswehrkrankenhaus Ulm, Ulm, Germany. ⁵⁵Institute of Pathology, University of Würzburg, Würzburg, Germany. ¹Institute of Pathology, University Hospital RWTH Aachen, Pauwelsstraße 30, 52074 Aachen, Germany. ²Institute for Medical Informatics, University Hospital RWTH Aachen, Pauwelsstraße 30, 52074 Aachen, Germany. ³Department of Anaesthesiology, Intensive Care, Emergency and Pain Medicine, University Hospital Würzburg, Oberdürrbacher Str. 6, 97080 Würzburg, Germany.

Received: 17 February 2022 Accepted: 4 March 2022
Published online: 28 March 2022

References

- Friedrichson B, Mutlak H, Zacharowski K, Piekarski F. Insight into ECMO, mortality and ARDS: a nationwide analysis of 45,647 ECMO runs. *Crit Care*. 2021;25(1):38.
- Nunez JJ, Gosling AF, O'Gara B, Kennedy KF, Rycus P, Abrams D, Brodie D, Shaefi S, Garan AR, Grandin EW. Bleeding and thrombotic events in adults supported with venovenous extracorporeal membrane oxygenation: an ELSO registry analysis. *Intensive Care Med*. 2022;48(2):213–24.
- Barbaro RP, MacLaren G, Boonstra PS, Iwashyna TJ, Slutsky AS, Fan E, Bartlett RH, Tonna JE, Hyslop R, Fanning JJ, et al. Extracorporeal membrane oxygenation support in COVID-19: an international cohort study of the Extracorporeal Life Support Organization registry. *Lancet*. 2020;396(10257):1071–8.
- Lebreton G, Schmidt M, Ponnaiah M, Folliquet T, Para M, Guihaire J, Lansac E, Sage E, Cholley B, Megarbane B, et al. Extracorporeal membrane oxygenation network organisation and clinical outcomes during the COVID-19 pandemic in Greater Paris, France: a multicentre cohort study. *Lancet Respir Med*. 2021;9(8):851–62.
- Pantel T, Roedl K, Jarczak D, Yu Y, Frings DP, Sensen B, Pinnschmidt H, Bernhardt A, Cheng B, Lettow I, et al. Association of COVID-19 with intracranial hemorrhage during extracorporeal membrane oxygenation for acute respiratory distress syndrome: a 10-year retrospective observational study. *J Clin Med*. 2021;11(1):28.
- Schmidt M, Chommeloux J, Frere C, Hekimian G, Combes A. Overcoming bleeding events related to extracorporeal membrane oxygenation in COVID-19 - Authors' reply. *Lancet Respir Med*. 2020;8(12):e89.
- von Stillfried S, Bulow RD, Rohrig R, Knuchel-Clarke R, Boor P. DeReg-Covid: autopsy registry can facilitate COVID-19 research. *EMBO Mol Med*. 2020;12(8):e12885.
- von Stillfried S, Bulow RD, Röhrig R, Boor P. First report from the German COVID-19 autopsy registry. *Lancet Reg Health - Europe*. 2022. <https://doi.org/10.1016/j.lanepe.2022.100330>.
- Arachchillage DJ, Rajakaruna I, Scott I, Gaspar M, Odho Z, Banya W, Vlachou A, Isgro G, Cagova L, Wade J, et al. Impact of major bleeding and thrombosis on 180-day survival in patients with severe COVID-19 supported with veno-venous extracorporeal membrane oxygenation in the United Kingdom: a multicentre observational study. *Br J Haematol*. 2022;196(3):566–76.
- Schulman S, Kearon C. Subcommittee on control of anticoagulation of the S, Standardization Committee of the International Society on Thrombosis and Haemostasis: definition of major bleeding in clinical investigations of antithrombotic medicinal products in non-surgical patients. *J Thromb Haemost*. 2005;3(4):692–4.
- Seeliger B, Doeblner M, Hofmaenner DA, Wendel-Garcia PD, Schuepbach RA, Schmidt JJ, Welte T, Hoepfer MM, Gillmann HJ, Kuehn C, et al. Intracranial hemorrhages on extracorporeal membrane oxygenation: differences between COVID-19 and other viral acute respiratory distress syndrome. *Crit Care Med*. 2022.
- Hwang J, Caturegli G, White B, Chen L, Cho SM. Cerebral microbleeds and intracranial hemorrhages in adult patients on extracorporeal membrane oxygenation-autopsy study. *Crit Care Explor*. 2021;3(3):e0358.
- German COVID-19 autopsy registry. German. https://www.drks.de/drks_web/navigate.do?navigationId=trial.HTML&TRIAL_ID=DRKS00025136.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

