

CERN-EP-2024-338  
2025/08/26

CMS-EXO-22-014

# Search for dark matter produced in association with one or two top quarks in proton-proton collisions at $\sqrt{s} = 13$ TeV

The CMS Collaboration\*

## Abstract

A search is performed for dark matter (DM) produced in association with a single top quark or a pair of top quarks using the data collected with the CMS detector at the LHC from proton-proton collisions at a center-of-mass energy of 13 TeV, corresponding to  $138 \text{ fb}^{-1}$  of integrated luminosity. An excess of events with a large imbalance of transverse momentum is searched for across 0, 1 and 2 lepton final states. Novel multivariate techniques are used to take advantage of the differences in kinematic properties between the two DM production mechanisms. No significant deviations with respect to the standard model predictions are observed. The results are interpreted considering a simplified model in which the mediator is either a scalar or pseudoscalar particle and couples to top quarks and to DM fermions. Axion-like particles that are coupled to top quarks and DM fermions are also considered. Expected exclusion limits of 410 and 380 GeV for scalar and pseudoscalar mediator masses, respectively, are set at the 95% confidence level. A DM particle mass of 1 GeV is assumed, with mediator couplings to fermions and DM particles set to unity. A small signal-like excess is observed in data, with the largest local significance observed to be 1.9 standard deviations for the 150 GeV pseudoscalar mediator hypothesis. Because of this excess, mediator masses are only excluded below 310 (320) GeV for the scalar (pseudoscalar) mediator. The results are also translated into model-independent 95% confidence level upper limits on the visible cross section of DM production in association with top quarks, ranging from 1 pb to 0.02 pb.

*Published in the Journal of High Energy Physics as doi:10.1007/JHEP08(2025)085.*



# 1 Introduction

Astrophysical observations point to the existence of dark matter (DM) [1]. Its existence has been inferred from gravitational effects on galaxies and other large-scale objects. While the nature of DM remains elusive, there are many models suggesting that DM could be explained in the context of particle physics [2]. In a large class of these new theories, DM is made of stable weakly interacting massive particles that interact with the standard model (SM) constituents through mediator particles, and hence may be produced at colliders, such as the CERN Large Hadron Collider (LHC). The DM particles are effectively invisible to the detectors located at the LHC interaction points, traversing these experiments undetected and leaving behind an unbalanced momentum in the plane transverse to the proton beams.

A convenient approach to model the coupling mechanisms between the SM and DM sectors is provided by simplified models, which can cover the main kinematic features exhibited in more complex ones. One possible simplification assumes the existence of either a new neutral scalar ( $\phi$ ) or pseudoscalar ( $a$ ) particle that can interact with both the SM fermion sector and a new fermionic Dirac DM particle ( $\chi$ ) [3, 4]. This construction involves a Yukawa-like coupling of the new mediator to fermions.

The Lagrangians of these models can be expressed as follows [4]:

$$\mathcal{L}_\phi = \mathcal{L}_{\text{SM}} + \frac{1}{2}(\partial_\mu \phi)^2 - \frac{1}{2}m_\phi^2 \phi^2 + i\bar{\chi}\not{\partial}\chi - m_\chi \bar{\chi}\chi - g_\chi \phi \bar{\chi}\chi - \frac{\phi}{\sqrt{2}} \sum_{q=u,d,s,c,b,t} g_q y_q q \bar{q}, \quad (1)$$

$$\mathcal{L}_a = \mathcal{L}_{\text{SM}} + \frac{1}{2}(\partial_\mu a)^2 - \frac{1}{2}m_a^2 a^2 + i\bar{\chi}\not{\partial}\chi - m_\chi \bar{\chi}\chi - ig_\chi a \bar{\chi}\gamma_5 \chi - \frac{ia}{\sqrt{2}} \sum_{q=u,d,s,c,b,t} g_q y_q \bar{q} \gamma_5 q, \quad (2)$$

where  $\mathcal{L}_{\text{SM}}$  is the SM Lagrangian, and  $y_q$  is the SM Yukawa coupling, normalized to the Higgs vacuum expectation value as  $y_q = \sqrt{2}m_q/v$ . There are only four additional parameters that control the kinematic properties or cross section: the mass of the DM particle ( $m_\chi$ ), the mass of the mediator ( $m_{\phi/a}$ ), the universal quark-mediator coupling ( $g_q$ ), and the DM-mediator coupling ( $g_\chi$ ). Under these conditions, the mediators would couple preferentially to heavy third-generation quarks, which motivates the focus on DM particles produced in association with a top quark pair ( $t\bar{t}$ +DM events) or a single top quark ( $t/\bar{t}$ +DM events) [5]. The main processes involving the  $t\bar{t}$ +DM and  $t/\bar{t}$ +DM production in the context of this simplified model are shown in Fig. 1.




Figure 1: Principal production diagrams in the context of the simplified model with a scalar/pseudoscalar ( $\phi/a$ ) mediator for the associated production of a pair of DM particles ( $\chi$ ) with a top quark pair (left) and a single top quark in both  $t$ -channel (center), and  $tW$ -channel (right) production modes. The additional quark  $q$  in the  $t$ -channel diagram is often produced at high pseudorapidity.

Complete models can also lead to enhanced couplings between the heavy quarks and the various DM mediators. One such case is the type II two-Higgs doublet model extended by an additional light pseudoscalar boson  $a$  (2HDM+a) [6, 7], featuring a differentiable coupling of scalar or pseudoscalar mass eigenstates to up-type and down-type fermions for a type II 2HDM configuration, and in which, for relatively low values of  $\tan \beta$ , the production of top quarks accompanied by DM particles is favored. This scenario is characterized by a higher number of parameters that control the kinematic properties with respect to the simplified model presented above and it is not considered in this paper. The simplified model considered in this analysis has a similar behavior to the 2HDM+a model in the limit of high additional Higgs boson masses when the charged Higgs bosons are completely decoupled.

Further models that could provide a mediator to the dark sector are those that propose axions or axion-like particles (ALPs, denoted as  $A$  in the following), which are spin-0 particles that do not interact with the SM gauge groups. The ALPs are a common feature in many extensions of the SM, where they emerge as pseudo-Nambu-Goldstone bosons because of an approximate axion shift symmetry. This symmetry allows for the masses of ALPs to be significantly smaller than the energy scale of the underlying ultraviolet model, making them promising candidates for studies at the LHC. Initially, axions were proposed as a solution to the strong-CP problem [8–11]. The ALPs are also promising candidates for mediators to the dark sector [12–15]. These models often feature a Dirac fermionic DM candidate  $\chi$  and a pseudoscalar mediator  $A$  that interact with SM fermions. Since the couplings of ALPs to fermions are typically proportional to the fermion masses, the ALP coupling to top quarks becomes particularly significant. In this paper, we also exploit our analysis to search for ALPs as DM mediators using a Lagrangian of the form:

$$\mathcal{L}_A = \mathcal{L}_{\text{SM}} + \frac{1}{2}(\partial_\mu A)(\partial^\mu A) - \frac{1}{2}m_A^2 A^2 + i\bar{\chi}\not{\partial}\chi - m_\chi\bar{\chi}\chi - ic_t \frac{m_t}{f_A}\bar{t}\gamma_5 t A - ic_\chi \frac{m_\chi}{f_A}\bar{\chi}\gamma_5\chi A, \quad (3)$$

where  $f_A$  denotes the ALP decay constant,  $c_t$  the ALP coupling to top quarks, and  $c_\chi$  its coupling to DM particles. The similarity of this Lagrangian to the one in Eq. (2) ensures that the DM signatures considered in Fig. 1 can also be interpreted in terms of models with an ALP mediator  $A$ .

Searches for DM in signatures containing a top quark pair or a single top quark have been previously carried out by the ATLAS and CMS Collaborations at a center-of-mass energy of 13 TeV. The CMS Collaboration has performed a search that specifically targets the  $t\bar{t}$ +DM production mode using the data set collected in 2016–2018 (Run 2) and covering all three commonly explored lepton categories (considering only muons and electrons) [16]. These lepton categories are named in some previous works as  $0\ell$ ,  $1\ell$ , and  $2\ell$  channels; here they will be referred to as all-hadronic (AH), single-lepton (SL), and dileptonic (DL) channels, respectively. The ATLAS Collaboration has delivered a combination across all channels using Run 2 data and considering both the  $t\bar{t}$ +DM and  $t/\bar{t}$ +DM production modes, optimizing for  $t\bar{t}$ +DM topologies [17], with a further combination with an update to the SL channel, only considering the  $t\bar{t}$ +DM signal [18]. Further, ATLAS analyses have searched for  $t/\bar{t}$ +DM final states, in boosted topologies predicted by 2HDM+a models [19, 20]. The work presented in this paper improves upon the constraints set by the CMS Collaboration with 2016 data from Run 2 on the combination of both the  $t\bar{t}$ +DM and  $t/\bar{t}$ +DM modes [21]. Various analysis improvements, as explained in later Sections, and the new dileptonic final state introduced lead to improved results of about 20% with respect to the increase in sensitivity from solely adding the data available from 2017 and 2018.

The analysis described in this paper is performed using the data collected with the CMS de-

tector in 2016–2018, corresponding to an integrated luminosity of  $138 \text{ fb}^{-1}$ . Although it probes similar signal topologies to the searches described above, the analysis includes considerable differences in strategy for some of the channels. In this work, a search optimized for both  $t\bar{t} + \text{DM}$  and  $t/\bar{t} + \text{DM}$  processes mediated by the presence of a neutral spin-0 particle is performed across all channels. The analysis makes use of dedicated search strategies for both DM production mechanisms to further enhance the separation power against the SM background. Novel multivariate techniques are employed using characteristic features of both signal production modes. An interpretation in the context of the spin-0 simplified DM model for both types of mediators is provided.

The paper is organized as follows: a brief introduction of the CMS detector and event reconstruction is presented in Section 2, followed by a description in Section 3 of the data and of the simulated samples used. The event selection is presented in Section 4, while Section 5 is dedicated to the treatment of experimental and theoretical systematic uncertainties in the analysis. The signal extraction methodology is discussed in Section 6, and the results obtained after this procedure are shown in Section 7. The paper is summarized in Section 8. Tabulated results are provided in the HEPData record for this analysis [22].

## 2 The CMS detector and event reconstruction

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass and scintillator hadron calorimeter (HCAL), each composed of a barrel and two endcap sections. Forward calorimeters extend the pseudorapidity ( $\eta$ ) coverage provided by the barrel and endcap detectors. Muons are detected in gas-ionization chambers embedded in the steel flux-return yoke outside the solenoid. A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, can be found in Ref. [23].

Events of interest are selected using a two-tiered trigger system [24]. The first level, composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a time interval of less than  $4 \mu\text{s}$ . The second level, known as the high-level trigger, consists of a farm of processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to around 1 kHz before data storage.

The particle-flow (PF) algorithm [25] aims to reconstruct and identify each individual particle in an event, with an optimized combination of information from the various elements of the CMS detector. The energy of photons is obtained directly from the ECAL measurement. The energy of electrons is obtained from a combination of the electron momentum at the primary interaction vertex as determined by the tracker, the energy of the corresponding ECAL cluster, and the energy sum of all bremsstrahlung photons spatially compatible with originating from the electron track. The muon track is obtained from the combination of central tracker and muon system information, and its curvature provides an estimate of the momentum. The energy of charged hadrons is determined from a combination of their momentum measured in the tracker and the matching ECAL and HCAL energy deposits, corrected for the response function of the calorimeters to hadronic showers. Finally, neutral hadrons are identified as HCAL energy clusters not linked to any charged-hadron trajectory, or as a combined ECAL and HCAL energy excess with respect to the expected charged-hadron energy deposit.

The primary vertex (PV) is taken to be the vertex corresponding to the hardest scattering in the event, evaluated using tracking information alone, as described in Section 9.4.1 of Ref. [26]. For each event, hadronic jets are clustered from the particles reconstructed with PF (PF candidates) using the infrared- and collinear-safe anti- $k_T$  algorithm [27, 28] with a distance parameter of 0.4. The jet momentum is determined as the vector sum of all particle momenta in the jet and is found from simulation to be within 5–10% of the parton’s generated momentum over the entire  $p_T$  spectrum and detector acceptance. Additional proton-proton (pp) interactions within the same or nearby bunch crossings (pileup) can contribute with additional tracks and calorimetric energy depositions to the jet momentum. To mitigate this effect, tracks identified as originating from pileup vertices are discarded and an offset correction is applied to correct for remaining contributions [29]. Jet energy corrections are derived from simulation and applied to calibrate the jet momentum. In situ measurements of the momentum balance in dijet, photon+jet, Z+jets, and multijet events are used to account for any residual differences in jet energy scale in data and simulation [30]. Additional selection criteria are applied to each jet to remove jets potentially dominated by anomalous contributions from various subdetector components or reconstruction failures [30].

The deep neural network (DNN)-based combined secondary vertex algorithm (DeepCSV) is used to identify jets originating from the hadronization of bottom quarks [31], denoted in the following as “b-tagged jets”. At the operating point of the tagging algorithm chosen for this analysis, the efficiency of identifying b quark jets in simulate  $t\bar{t}$  events is about 75%, averaged over all  $p_T$ , and the misidentification rate for light-flavor quark and gluon jets is about 1%. Scale factors are applied to the simulated samples to reproduce the b tagging efficiency measured in data.

The missing transverse momentum vector  $\vec{p}_T^{\text{miss}}$  is defined as the negative vector  $p_T$  sum of all PF particles originating from the primary vertex; its magnitude is defined as  $p_T^{\text{miss}}$ . Jet energy scale and resolution corrections are also propagated to the  $\vec{p}_T^{\text{miss}}$  calculation.

### 3 Data and simulated samples

The data used in this search were collected by the CMS detector in 2016–2018 and correspond to an integrated luminosity of  $138 \text{ fb}^{-1}$ . Several triggers were used to collect the data, either requiring large  $p_T^{\text{miss}}$  or the presence of one or two leptons (electrons or muons). The  $p_T^{\text{miss}}$ -based triggers are employed to select events that do not contain leptons (i.e., for the AH channel) if they have  $p_T^{\text{miss}}$  and missing hadronic activity  $H_T^{\text{miss}}$  [24] both above 120 GeV, where missing hadronic activity is defined as the negative vector sum of jet transverse momenta. These triggers are nearly 100% efficient for events with  $p_T^{\text{miss}}$  of at least 250 GeV. A second set of triggers, used in the SL channel, requires the presence of at least one isolated electron (muon) with  $p_T > 27$  (24) GeV. The corresponding trigger efficiencies are above 90% for leptons with  $p_T > 30$  GeV. The last set of triggers, for the DL channel, uses a combination of single- and dilepton triggers to maximize the efficiency, which is on average 98% for events with two leptons with  $p_T > 25$  (20) GeV.

Monte Carlo (MC) simulated samples of the main SM backgrounds and of the signal processes are used to optimize the event selection and to improve the background estimation. The SM processes with the largest contributions to the background in the various final states are  $t\bar{t}$ , W+jets, and Z+jets production, although the precise composition depends on the specific channel under consideration. Simulated events of  $t\bar{t}$  production and single top quark processes ( $t$ -,  $s$ -, and  $tW$ -channel) are generated at next-to-leading order (NLO) in quantum chromodynam-

ics (QCD) using the POWHEG v2 [32–37] event generators. For the  $t\bar{t}$  process, the top quark  $p_T$  distribution is reweighted to match next-to-NLO (NNLO) theoretical computations [38]. The samples for top quark pair associated production with a W boson ( $t\bar{t}+W$ ) and a Z boson ( $t\bar{t}+Z$ ) plus up to two additional partons are generated based on the NLO ME calculations implemented in MADGRAPH5\_aMC@NLO and the FxFx [39] prescription to merge multileg processes. Samples of Z+jets and W+jets are generated using leading order (LO) predictions with up to four partons in the final state using MADGRAPH5\_aMC@NLO [40] (v2.2.2 in 2016, v2.4.2 in 2017 and 2018) with the MLM prescription [41] for matching jets from the matrix element (ME) calculation to the parton shower description. Dedicated electroweak [42–47] and QCD (calculated with MADGRAPH5\_aMC@NLO) LO to NLO corrections ( $K$  factors), parameterized as functions of the generated boson  $p_T$ , are applied to Z+jets and W+jets events. Multijet events are simulated at LO either using PYTHIA or MADGRAPH5\_aMC@NLO event generators, making use of the MLM matching for the latter case. Other processes with a smaller contribution, such as diboson production, are generated at NLO using either MADGRAPH5\_aMC@NLO using FxFx matching or with POWHEG v2 and normalized to the most accurate cross section calculations available [48, 49].

The  $t\bar{t}+DM$  and  $t/\bar{t}+DM$  signal processes are simulated at LO with MADGRAPH5\_aMC@NLO v2.6.1 with one and zero additional partons, respectively, in the matrix element calculations using the simplified model referenced in Section 1. The benchmark values chosen for the coupling parameters are  $g_\chi = g_q = 1$ , following the recommendations of the LHC Dark Matter Working Group [50]. In addition, the mass of the DM candidate is set to  $m_\chi = 1$  GeV as a benchmark point, taking into account that for the on-shell production of the mediators, the impact of the DM mass in the  $p_T^{\text{miss}}$  distribution is irrelevant. For the off-shell production of  $\phi/a$ , there is typically a moderate change in the distribution. However, these scenarios are not considered in this study because of their highly suppressed cross sections. The parameter scan focuses on variations of the parameter  $m_{\phi/a}$ , chosen from the following list of mass points (for both type of mediators and both  $t\bar{t}+DM$  and  $t/\bar{t}+DM$  production modes): 50, 100, 150, 200, 250, 300, 350, 400, 450, and 500 GeV.

The initial-state partons are modeled with the NNPDF 3.0 [51] parton distribution function (PDF) set, of the same order in QCD as was used for the matrix element calculation for the samples corresponding to the 2016 period, while the NNPDF 3.1 NNLO [52] PDF set is used for the 2017 and 2018 samples. Parton showering and hadronization are handled by PYTHIA v8.226 (8.230) [53] using the CUETP8M1(2) underlying event tune [54] for samples in the 2016 period, and the CP5 tune [55] for most samples corresponding to the 2017 and 2018 periods. All signal and background samples are processed using GEANT4 [56] to provide a full simulation of the CMS detector, including a simulation of the previously mentioned triggers. The effects of additional pp interactions in the same or adjacent bunch crossings, referred to as pileup, are included in all simulation samples. To match the simulation distribution of genuine pileup interactions with the one observed in data, a reweighting procedure is implemented. Correction factors are derived and applied to the simulated samples to match the trigger efficiencies measured in data. Additional corrections are applied to cover remaining residual differences between data and simulation that arise from the lepton identification and reconstruction efficiencies, as well as from b-tagged jet identification efficiencies.

## 4 Event selection

This search defines several orthogonal signal regions (SRs) that are statistically combined in a simultaneous global fit of the  $p_T^{\text{miss}}$  spectrum of the events in those regions. Control regions

(CRs) enriched in the major background processes are included in the fit for each channel to improve the estimates of the SM contributions. Events are classified into three mutually exclusive “channels”, based on the number of leptons in the final state: the AH channel, which contains events with no electrons or muons with  $p_T > 10$  GeV; the SL channel, which contains events with one lepton with  $p_T > 35(30)$  GeV for electrons (muons) and with no other lepton with  $p_T > 10$  GeV; and the DL channel, which contains events with one lepton with  $p_T > 25$  GeV, a second lepton with  $p_T > 20$  GeV, and no other lepton with  $p_T > 10$  GeV.

In the AH CRs and SL channel, electrons are selected if they have  $p_T > 35$  GeV and  $|\eta| < 2.1$ , and muons are selected if they have  $p_T > 30$  GeV and  $|\eta| < 2.4$ . For the DL channel, electrons and muons are selected if they have  $p_T > 20$  GeV and  $|\eta| < 2.4$ . In all channels, events containing additional leptons with  $p_T > 10$  GeV and  $|\eta| < 2.4$  are vetoed. To ensure that candidate leptons are well measured, identification requirements based on hit information in the tracker and muon systems and on energy deposits in the calorimeters are imposed. Leptons are further required to be isolated from hadronic activity to reject leptons within jets that could arise, for example, from the decay of  $b$  quarks. A relative isolation quantity is defined as the scalar  $p_T$  sum of all PF candidates within a  $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$  cone of radius 0.3 (0.4) centered around the electron (muon) candidate, where  $\phi$  is the azimuthal angle in radians, divided by the lepton. The requirement put on this relative isolation depends on the  $p_T$  for electrons and is on average required to be less than 0.2, while is required to be less than 0.15 for muons [57, 58].

Each channel SR is further divided based on the number of  $b$ -tagged jets ( $n_b$ ) into a region with exactly one  $b$ -tagged jet (1  $b$ ), targeting the  $t/\bar{t}$ +DM process, and a region with two or more  $b$ -tagged jets (2  $b$ ), targeting the  $t\bar{t}$ +DM process. The regions with exactly one  $b$ -tagged jet in the AH and SL channels are further subdivided into exactly zero or  $\geq 1$  forward jet regions (0FJ and 1FJ) to increase the sensitivity to the  $t$ -channel  $t/\bar{t}$ +DM process, which tends to have a forward jet, as shown in Fig. 1. Jet candidates are required to have  $p_T > 30$  GeV in the AH and SL channels, or  $p_T > 20$  GeV in the DL channel, and are categorized as “central” if they lie within  $|\eta| < 2.4$  and as “forward” if they are within  $2.4 < |\eta| < 4.0$ . The  $b$ -tagged jets are required to have the same  $p_T$  threshold and to lie within  $|\eta| < 2.4$ .

In each channel, a set of variables is identified to discriminate between signal and background events. In the AH and SL channels, a selection on these variables is applied to increase the signal significance, which is optimized using as a figure of merit the ratio between the number of expected signal events and the square root of the expected SM background contributions. In the DL channel, the discriminating variables are employed to train a neural network.

#### 4.1 All-hadronic signal regions

Events categorized into the AH channel are events with no electrons or muons with  $p_T > 10$  GeV, at least one identified  $b$ -tagged jet, at least three jets with  $p_T > 30$  GeV, and  $p_T^{\text{miss}} > 250$  GeV. The dominant background after this selection consists of SL  $t\bar{t}$  ( $t\bar{t}(1\ell)$ ) events, where the lepton either falls out of detector acceptance or is not identified, providing an additional source of  $p_T^{\text{miss}}$ . To reduce this background, we require the transverse mass  $m_T^b$  of the  $\vec{p}_T^{\text{miss}}$  vector and of a  $b$ -tagged jet to be greater than 140 GeV. The  $m_T^b$  variable is defined as:

$$m_T^b = \sqrt{2p_T^{\text{miss}}p_T^b[1 - \cos(\Delta\phi)]}, \quad (4)$$

where  $p_T^b$  is the transverse momentum of the  $b$ -tagged jet and  $\Delta\phi$  is the opening angle between the  $b$ -tagged jet direction and the  $\vec{p}_T^{\text{miss}}$  in the transverse plane. For  $t\bar{t}$  background events,  $m_T^b$

tends to have values below or around the top quark mass in the case where the b-tagged jet belongs to the top quark decay chain and the lepton is not identified. For the calculation, we choose the b-tagged jet with the highest DeepCSV discriminant value if there is more than one candidate.

Studies have shown that to further reduce the  $t\bar{t}(1\ell)$  background, together with that from  $Z \rightarrow \nu\bar{\nu}$  events, an effective variable to use is  $p_T(j_1)/H_T$ , which is defined as the ratio of the leading  $p_T$  jet in the event to  $H_T$ , the scalar  $p_T$  sum of all reconstructed jets in the event with  $p_T > 30$  GeV within  $|\eta| < 2.4$ . The  $t\bar{t}$ +DM process, which has six jets at the ME level, tends to have lower values of this observable than the  $t\bar{t}(1\ell)$  and  $Z \rightarrow \nu\bar{\nu}$  backgrounds, which have fewer jets. We require  $p_T(j_1)/H_T < 0.5$  in the  $n_b \geq 2$  category. The  $t/\bar{t}$ +DM events, which have fewer central jets, tend to exhibit a distribution similar to that of the background, so no such requirement on the  $p_T(j_1)/H_T$  is applied in the  $n_b = 1$  category.

Another background in this channel is QCD multijet production. For this process, no intrinsic  $p_T^{\text{miss}}$  is expected, so the observed  $p_T^{\text{miss}}$  is mostly the result of jet mismeasurements. For these events,  $\vec{p}_T^{\text{miss}}$  is often aligned with one of the leading jets. As a result, the minimum opening angle between each of the two leading jets and the  $\vec{p}_T^{\text{miss}}$  is required to satisfy  $\min \Delta\phi(j_{1,2}, \vec{p}_T^{\text{miss}}) > 0.8$  radians to reduce this background.

The dominant backgrounds after the selection presented in Table 1 arise from  $t\bar{t}$ , W+jets, and  $Z \rightarrow \nu\bar{\nu}$  processes, with QCD multijet events, single top quark, Drell–Yan (DY), and diboson production giving smaller contributions.

## 4.2 Single-lepton signal regions

Events in the SL channel are selected by requiring the presence of one electron (muon) with  $p_T > 35(30)$  GeV,  $\geq 1$  identified b-tagged jets, at least two jets with  $p_T > 30$  GeV, and  $p_T^{\text{miss}} > 250$  GeV. After this selection, the dominant backgrounds are from  $t\bar{t}$  and W+jets processes.

To reduce the dominant backgrounds and improve the signal sensitivity, we impose a requirement on the transverse mass  $m_T$ , which is calculated as:

$$m_T = \sqrt{2p_T^{\text{miss}}p_T^\ell[1 - \cos(\Delta\phi)]}, \quad (5)$$

where  $p_T^\ell$  is the transverse momentum of the lepton and  $\Delta\phi$  is the opening angle between the lepton direction and the  $\vec{p}_T^{\text{miss}}$  in the transverse plane. The  $m_T$  variable is expected to be less than the W boson mass for leptonic on-shell W decays in  $t\bar{t}$  and W+jets events, while for signal, off-shell W decays, and dileptonic decays of  $t\bar{t}$ , the  $m_T$  variable is expected to exceed the W boson mass because of the additional  $p_T^{\text{miss}}$  in the event. A requirement of  $m_T > 140$  GeV reduces the background contributions from single-lepton events and enhances the analysis sensitivity to the DM signal.

After applying the  $m_T$  selection, the remaining  $t\bar{t}$  contributions are primarily from events with both top quarks decaying leptonically ( $t\bar{t}(2\ell)$ ) and with one lepton not identified. This background can be further reduced by making use of the  $m_{T2}^W$  variable [59], which is defined as the minimal value of the mass of a particle assumed to be pair produced that decays to a W boson and a b quark jet. The W bosons are assumed to be produced on-shell and to decay leptonically, where one of the two leptons is not detected. Based on the variable definition, in  $t\bar{t}(2\ell)$  events the  $m_{T2}^W$  distribution has a kinematic end point at the top quark mass for perfect detector response, while this is not the case for signal events, where two additional DM particles are present. The calculation of  $m_{T2}^W$  requires two b-tagged jets, where one comes from

the same decay chain as the reconstructed lepton. If only one b-tagged jet is present in the event, each of the first three (or two in three-jet events) leading non-b-tagged jets is assumed as the second b-tagged jet in the calculation. The  $m_{T2}^W$  is then evaluated for all possible jet-lepton combinations and the minimum  $m_{T2}^W$  value is considered to discriminate between signal and background events.

If two or more b-tagged jets are identified in the events,  $m_{T2}^W$  is calculated using all possible jet-lepton combinations. The smallest of all the  $m_{T2}^W$  values is taken as the event discriminant. A requirement of  $m_{T2}^W > 180$  GeV is applied in the analysis.

In addition, jets and the  $\vec{p}_T^{\text{miss}}$  vector tend to be more separated in the transverse plane in signal events than in  $t\bar{t}$  background processes. To improve the search sensitivity, the minimum opening angle  $\min \Delta\phi(j_{1,2}, \vec{p}_T^{\text{miss}})$  in the transverse plane between the direction of each of the first two leading- $p_T$  jets with  $|\eta| < 2.4$  and the  $\vec{p}_T^{\text{miss}}$  vector is required to be greater than 0.8 radians.

To reduce the  $t\bar{t}$  background further, the transverse mass  $m_T^b$  is required to be greater than 140 GeV.

To reject the  $t\bar{t}(2\ell)$  background, the modified topness variable  $t$  [60] is introduced. Unlike  $m_{T2}^W$ , which minimizes the transverse mass of the invisible particles, the modified topness minimizes the reconstructed center-of-mass energy of the event, subject to the conditions that both top quarks and W bosons are on shell. It is defined as:

$$t = \ln(\min S) \quad (6)$$

with

$$S = \frac{(m_W^2 - (p_\nu + p_\ell)^2)^2}{a_W^4} + \frac{(m_t^2 - (p_b + p_W)^2)^2}{a_t^4}, \quad (7)$$

where  $p$  and  $m$  denote the four-momentum and mass of the particles involved, respectively. The minimization of the variable  $S$  is done with respect to all components of the three momentum  $\vec{p}_W$  and the component of the three momentum  $\vec{p}_\nu$  along the beam line. The sum is performed over all five assumed final-state particles from the top quark and W boson decays. The  $a_W = 5$  GeV and  $a_t = 15$  GeV parameters determine the relative weighting of the mass shell conditions. The inputs to the calculation of  $S$  are two jets, a lepton, and the  $p_T^{\text{miss}}$ . The value of  $t$  quantifies how well an event is compatible with the  $t\bar{t}(2\ell)$  hypothesis. Rather than defining a single selection criterion on this variable, we split all the SRs in the SL channel into two further categories  $t \leq 0$  (T1) and  $t > 0$  (T2).

Though significantly reduced by these selections,  $t\bar{t}$  production, in particular  $t\bar{t}(2\ell)$ , remains the dominant background in this channel, followed by W+jets, single top quark, diboson, and DY production.

### 4.3 Dileptonic signal regions

Events in the dileptonic channel are selected by requiring at least two leptons with  $p_T > 25$  and 20 GeV, and at least one b-tagged jet with  $p_T > 30$  GeV. The invariant mass  $m_{\ell\ell}$  of the two-lepton system is required to satisfy  $m_{\ell\ell} > 20$  GeV to suppress light resonances, and for the case of two electrons or two muons, the region  $76 < m_{\ell\ell} < 106$  GeV is also excluded to suppress DY production. This selection gives a region that is dominated by  $t\bar{t}$  processes, which, because of the presence of two neutrinos in the final state, has a fairly hard  $p_T^{\text{miss}}$  spectrum that cannot

be simply removed with a  $p_T^{\text{miss}}$  requirement. A selection is therefore imposed on  $m_{T2}^{\ell\ell}$  [61, 62], which is defined as:

$$m_{T2}^{\ell\ell} = \min_{\vec{p}_{T,v} + \vec{p}_{T,\bar{v}} = \vec{p}_T^{\text{miss}}} [\max\{M_T(m_\ell, m_{\bar{v}}, \vec{p}_{T,\ell}, \vec{p}_{T,\bar{v}}), M_T(m_{\bar{\ell}}, m_v, \vec{p}_{T,\bar{\ell}}, \vec{p}_{T,v})\}], \quad (8)$$

with

$$M_T(m_1, m_2, \vec{p}_{T,1}, \vec{p}_{T,2}) = \sqrt{m_1^2 + m_2^2 + 2(E_{T,1}E_{T,2} - \vec{p}_{T,1} \cdot \vec{p}_{T,2})}, \quad (9)$$

where  $m_i$ ,  $\vec{p}_{T,i}$ , and  $E_{T,i}$  correspond to the mass, transverse momentum vector, and transverse energy of the particle  $i$  respectively, while  $\vec{p}_T^{\text{miss}}$  is the measured missing transverse momentum vector. This variable offers information on the mass of pair-produced particles in situations where both particles decay to a final state with undetected particles, as in the case of two leptonically decaying  $W$  bosons produced from top quark pairs. If the visible components in the decay chain are measured correctly, this variable has a kinematic endpoint at the  $W$  boson mass for the  $t\bar{t}$  processes but not for signal, because of the additional  $p_T^{\text{miss}}$  from DM.

After requiring  $m_{T2}^{\ell\ell} > 80 \text{ GeV}$ , the main remaining backgrounds are from the  $t\bar{t}$  production because of  $p_T^{\text{miss}}$  mismeasurements,  $tW$  production,  $DY$ , and  $t\bar{t}Z$  events, where the  $Z$  boson decays to neutrinos ( $t\bar{t}Z \rightarrow t\bar{t}\nu\bar{\nu}$ ), giving a very similar signature to the signal process. Most of the backgrounds in the DL channel are well modeled by the MC simulation. However, the  $DY$  process is difficult to model as it normally only enters the selection if  $p_T^{\text{miss}}$ , and hence the  $m_{T2}^{\ell\ell}$ , are significantly mismodeled. The DL signal regions are therefore split by lepton flavor into same-flavor (SF) regions with two electrons or two muons, which will be relatively enriched in the  $DY$  process, and different flavor (DF) regions with one electron and one muon, which will be depleted in  $DY$  events. A method is then used to estimate the  $DY$  in the more enriched  $t/\bar{t} + \text{DM}$  SF region (as discussed in Section 4.4.3), based on control samples in data.

#### 4.3.1 Neural network optimization

As previously mentioned, for the DL channel the  $p_T^{\text{miss}}$  distribution alone does not provide a strong discrimination of the signal from the background. The signal sensitivity in the DL channel can be improved by employing a neural network (NN) trained on a number of discriminating variables. A separate NN is trained in each of the 1 b tag and 2 b tag regions (the SF and DF lepton regions are combined to maximize the number of events used for the training) and for each of the mediator hypotheses (scalar or pseudoscalar). Regarding the  $t/\bar{t} + \text{DM}$  and  $t\bar{t} + \text{DM}$  processes, both types of signals are considered in the 1 b tag and 2 b tag SRs since a large fraction of  $t\bar{t} + \text{DM}$  events can contribute to the 1 b tag region if one of the b jets is not tagged. Only one NN is trained for all mediator mass hypotheses since the performance of the NN is found not to significantly improve when training on a single mass point compared to the ensemble of all mass points.

Various distributions are considered as inputs for the NN, and the following variables are found to improve the signal sensitivity (in addition to  $p_T^{\text{miss}}$  and  $m_{T2}^{\ell\ell}$ , mentioned previously):

- $|\Delta\phi(\ell, \bar{\ell})|$ , the opening angle between the two leptons. Since top quarks decay to a b quark and a  $W$  boson before hadronization, the spin information is propagated to the  $W$  boson and thus to the angular distribution of the leptons. For  $t/\bar{t} + \text{DM}$   $tW$ -channel processes, the (pseudo)scalar mediator is radiated from the top quark, affecting the final top polarization and therefore the correlation with the  $W$  boson. This observable contains information regarding the polarization correlation of the two particles and helps in separating the  $tW + \text{DM}$  from the SM single top quark  $tW$  production because of the effect of the mediator coupling to top quarks.

- $|\Delta\phi(\vec{p}_T^{\text{miss}}, \ell\bar{\ell}b)|$ , the opening angle between the two-lepton plus (leading) b-tagged jet and  $\vec{p}_T^{\text{miss}}$ . This variable allows differentiation of  $t/\bar{t}$ +DM events from the dominant  $t\bar{t}$  background since in  $t/\bar{t}$ +DM events the  $\ell\bar{\ell}b$  system is expected to be back-to-back with respect to  $\vec{p}_T^{\text{miss}}$ . This is not the case for the  $t\bar{t}$  production, which has another b jet present.

In the 2 b tag region, the four-momenta of the top quarks can be reconstructed from all the visible decay products of the  $t\bar{t}$  system. This can be achieved by constraining the neutrino four-momenta using the on-shell conditions for the top quarks, W bosons, and neutrinos along with the fact that the  $p_T^{\text{miss}}$  for  $t\bar{t}$  processes should correspond to the sum of the transverse momentum of the neutrinos. While this condition is valid for  $t\bar{t}$  processes, it does not hold for the  $t\bar{t}$ +DM signal where the DM particles produced also contribute to the total  $p_T^{\text{miss}}$  in the event. Therefore, a variant of traditional kinematic reconstruction algorithms [63] is used. This algorithm first applies the on-shell conditions to limit possible neutrino solutions to ellipses in momentum space and then attempts to assign the neutrino momenta as closely as possible to  $p_T^{\text{miss}}$ . If the entire  $p_T^{\text{miss}}$  can be assigned to neutrino momenta, the ellipses for the neutrino and anti-neutrino will intersect, and the point of intersection with the smallest  $m_{t\bar{t}}$  is taken as a solution for the neutrino momenta. Otherwise (as is usually the case because of the  $m_{T2}^{\ell\ell} > 80$  GeV selection), the ellipses do not intersect, and the point of closest approach is taken as a solution. Events that fail this kinematic reconstruction (which only happens if the on-shell conditions are not satisfied and represents about 5% of signal and background events) are not considered in this region. This reconstruction of the top quarks allows the use of additional discriminating variables in the 2 b tag region based on the top quark kinematic properties:

- $p_T^{\text{dark}}$ : this variable defines the amount of  $p_T^{\text{miss}}$  that cannot be assigned to the neutrinos in the top quark kinematic reconstruction, i.e., the distance of closest approach of the ellipses. This variable tends to have higher values for  $t\bar{t}$ +DM events with respect to  $t\bar{t}$  production because of the additional source of genuine  $p_T^{\text{miss}}$  from DM particles.
- $c_{\text{hel}}$ : the angle between the two leptons when boosted back into the rest frame of their parent top quarks. This quantity is sensitive to spin correlations.
- $|\Delta\phi(t, \bar{t})|$ : the top quark and antiquark are expected to be approximately back-to-back in the azimuthal plane for  $t\bar{t}$  production, while for  $t\bar{t}$ +DM events they are expected to recoil against the DM mediator.

The NNs were trained using Keras with a Tensorflow backend. Each contained three densely connected hidden layers using Rectified Linear Unit (ReLU) activation function, and an output layer with nodes for signal and background using the “softmax” activation function. Categorical cross-entropy was used for the activation function, and the network was trained for 50 epochs. The MC data sets were split into training, validation and test data sets; no evidence was observed for overtraining between the training and validation data sets, and no significant difference in the final distributions was observed between the test and training data sets. The NN score was used in the signal extraction, explained in more detail in Section 6, with a binning designed to maximize sensitivity while keeping a minimum of 5 expected background events in each bin.

#### 4.4 Control regions

After the event selection presented in Table 1 is applied, the leading SM background contributions in the different regions are evaluated. For the AH regions, the main backgrounds arise from single-lepton  $t\bar{t}$  and W+jets events in which the lepton is not identified, and Z boson pro-

Table 1: Final event selection requirements for the AH, SL, and DL SRs. For the SL channel, a categorization in terms of modified topness, with bins of  $t \leq 0$  and  $t > 0$ , is also applied after the event selection. The DL channel is split into SF  $e^+e^-/\mu^+\mu^-$  and DF  $e^\pm\mu^\mp$  regions.

|                                                            | All-hadronic SRs |                  |             | Single-lepton SRs |                  |             | Dilepton SRs |             |
|------------------------------------------------------------|------------------|------------------|-------------|-------------------|------------------|-------------|--------------|-------------|
|                                                            | $0\ell, 1b, 0FJ$ | $0\ell, 1b, 1FJ$ | $0\ell, 2b$ | $1\ell, 1b, 0FJ$  | $1\ell, 1b, 1FJ$ | $1\ell, 2b$ | $2\ell, 1b$  | $2\ell, 2b$ |
| $n_{\text{lep}}$                                           |                  | =0               |             |                   | =1               |             | =2           |             |
| $n_{\text{jet}}$                                           |                  | $\geq 3$         |             |                   | $\geq 2$         |             | $\geq 1$     |             |
| $n_b$                                                      | =1               | =1               | $\geq 2$    | =1                | =1               | $\geq 2$    | =1           | $\geq 2$    |
| Forward jets                                               | =0               | $\geq 1$         | —           | =0                | $\geq 1$         | —           | —            | —           |
| $p_T(j_1)/H_T$                                             | —                | —                | <0.5        | —                 | —                | —           | —            | —           |
| $p_T^{\text{miss}}$ [GeV]                                  |                  | >250             |             |                   | >250             |             | —            |             |
| $m_T$ [GeV]                                                |                  | —                |             |                   | >140             |             | —            |             |
| $m_{T2}^W$ [GeV]                                           |                  | —                |             |                   | >180             |             | —            |             |
| $\min \Delta\phi(j_{1,2}, \vec{p}_T^{\text{miss}})$ [rad.] |                  | >0.8             |             |                   | >0.8             |             | —            |             |
| $m_T^b$ [GeV]                                              |                  | >140             |             |                   | >140             |             | —            |             |
| $m_{\ell\ell}$ [GeV]                                       |                  | —                |             |                   | —                |             | >20          |             |
| $ m_{\ell\ell} - m_Z $ [GeV]                               |                  | —                |             |                   | —                |             | >15 (SF)     |             |
| $m_{T2}^{\ell\ell}$ [GeV]                                  |                  | —                |             |                   | —                |             | >80          |             |
| Pass $t\bar{t}$ reco                                       |                  | —                |             |                   | —                |             | —            | yes         |

duction in which the Z boson decays into two neutrinos, leading to genuine  $p_T^{\text{miss}}$ . In the SL SRs, the main backgrounds are dileptonic  $t\bar{t}$  events in which one of the leptons is not identified, and W+jets events. In the DL SRs, the backgrounds are dileptonic  $t\bar{t}$  and tW events, DY production, and  $t\bar{t}Z \rightarrow t\bar{t}\nu\bar{\nu}$  processes.

To improve the estimation of the backgrounds, control samples in data are used. In the AH and SL channels, CRs enhanced in the different background sources are used to derive correction factors as a function of  $p_T^{\text{miss}}$  from the comparison of the  $p_T^{\text{miss}}$  distribution between data and simulation. These corrections are extracted simultaneously across the CRs and SRs for each channel in a global fit, as explained in more detail in Section 6. The residual background processes are modeled with simulation.

In the DL channel, a validation region (VR) enriched in  $t\bar{t}$  and tW events is used to verify their modeling for all the input variables used in the NN. These distributions all show good agreement between data and simulations within uncertainties and therefore no correction factor is used. For the DY and  $t\bar{t}Z \rightarrow t\bar{t}\nu\bar{\nu}$  processes, dedicated CRs are used to derive corrections in the fit. For the DY process, these corrections are derived as a function of the NN score in a similar way to the AH and SL channels, while for  $t\bar{t}Z \rightarrow t\bar{t}\nu\bar{\nu}$ , only an overall normalization factor is considered.

#### 4.4.1 All-hadronic control regions

For the AH SRs, four independent sets of CRs are defined. The first set of CRs is designed to isolate single-lepton  $t\bar{t}$  processes by selecting events with exactly one lepton,  $n_{\text{jet}} \geq 3$ ,  $n_b \geq 1$ ,  $p_T^{\text{miss}} > 250$  GeV,  $\min \Delta\phi(j_{1,2}, \vec{p}_T^{\text{miss}}) \geq 0.8$ , and  $m_T < 140$  GeV to avoid overlap with the SL SRs.

The second set of CRs is enhanced in single-lepton W+jets processes by selecting events with exactly one lepton,  $n_{\text{jet}} \geq 3$ ,  $n_b = 0$ ,  $p_T^{\text{miss}} > 250$  GeV, and in order to avoid overlap with the SL W+jets CR,  $m_T < 140$  GeV.

Additional CRs are designed to model the background originating from Z+jets production, where the Z boson decays into a pair of neutrinos ( $Z \rightarrow \nu\bar{\nu}$ ). Here we use the Z boson decays to an OS SF dilepton pair ( $Z \rightarrow \ell\ell$ ) as proxy events to emulate the kinematic properties of the  $Z \rightarrow \nu\bar{\nu}$  process. Events are selected by requiring two leptons with SF (i.e.,  $e^+e^-$  or  $\mu^+\mu^-$ ) and OS that satisfy the requirement  $60 < m_{\ell\ell} < 120$  GeV on their invariant mass. Additionally,

events must contain at least three jets and events with b-tagged jets are vetoed ( $n_b = 0$ ). To reproduce the  $p_T$  spectrum of  $Z \rightarrow \nu\bar{\nu}$  events, the two leptons are added to the  $\vec{p}_T^{\text{miss}}$ , giving a quantity referred to as hadronic recoil, which is required to be greater than 250 GeV.

The fourth CR is introduced to estimate QCD multijet events from data to reduce their associated uncertainties. Though QCD multijet events are a minor background in the AH SRs, they are characterised by large systematic uncertainties. These large systematic uncertainties are due to the fact that to pass the  $p_T^{\text{miss}} > 250$  GeV selection, they need to have a significant jet mismeasurement, which is difficult to model accurately in simulation. The associated CR is defined by requiring zero leptons,  $n_{\text{jet}} \geq 3$ ,  $n_b \geq 1$ ,  $p_T^{\text{miss}} > 250$  GeV, and  $\min \Delta\phi(j_{1,2}, \vec{p}_T^{\text{miss}}) < 0.8$ . The last requirement selects events where the  $p_T^{\text{miss}}$  is aligned with one of the leading jets, pointing to a mismeasurement.

A summary of the different AH CRs can be found in the first four columns of Table 2.

#### 4.4.2 Single-lepton control regions

The first set of CRs is designed to estimate dileptonic  $t\bar{t}$  events by requiring exactly two leptons,  $n_{\text{jet}} \geq 2$ ,  $n_b \geq 1$ , and  $p_T^{\text{miss}} > 250$  GeV. Additionally, events are required to have  $m_{T2}^{\ell\ell} < 80$  GeV to maintain the orthogonality with the DL SRs. To statistically enhance these CRs, the  $m_T$ ,  $m_{T2}^W$ , and forward jet selections are removed.

The second set of CRs is defined to isolate W+jets events by requiring exactly one lepton,  $n_{\text{jet}} \geq 2$ ,  $n_b = 0$ ,  $p_T^{\text{miss}} > 250$  GeV, and  $m_T > 140$  GeV. The  $n_b = 0$  requirement makes this CR orthogonal to the SL SRs.

The selection criteria are summarized in the central columns of Table 2.

#### 4.4.3 Dilepton control regions

For the DL channel, a VR is used to check the agreement for the dominant  $t\bar{t}$  background between data and simulation for the distributions used as input variables to the NN. This region is defined with the same selections used in the DL SRs (exactly two leptons, at least one jet and b-tagged jet,  $m_{\ell\ell} > 20$  GeV, and outside the Z boson mass window for SF leptons), except for the inversion of the  $m_{T2}^{\ell\ell}$  selection requirement. No further correction to these processes in the fit is found to be necessary.

The DY process is found to be not well modeled in simulation. In particular, the  $p_T^{\text{miss}}$  distribution and the  $m_{T2}^{\ell\ell}$  distribution, which are the two most sensitive NN input variables, are not well described. A CR is defined with an identical selection to the DL SRs (exactly two leptons, at least one jet and b-tagged jet,  $m_{\ell\ell} > 20$  GeV, and  $m_{T2}^{\ell\ell} > 80$  GeV), except only SF ( $e^+e^-/\mu^+\mu^-$ ) leptons are considered inside the Z boson mass window  $76 < m_{\ell\ell} < 106$  GeV. Separate CRs corresponding to the 1 b tag and 2 b tag regions were considered, since this distribution is also known to be poorly modeled. However, the 2 b tag region was found to not be dominated by DY events even inside the Z boson mass window. Therefore, a single CR requiring exactly 1 b-tagged jet is used to predict the rate of DY in the 1 b tag SF SR as a function of the NN score.

To model the background originating from  $t\bar{t}Z \rightarrow t\bar{t}\nu\bar{\nu}$  production, a CR enriched in  $t\bar{t}Z$  processes, where the Z decays into electrons or muons, is used. To reduce the statistical uncertainty, a CR targeting SL  $t\bar{t}$  decays is used, which is defined by requiring exactly three leptons, with  $p_T > 25, 20, 20$  GeV respectively, along with at least three jets with  $p_T > 30$  GeV for the leading jet,  $p_T > 20$  GeV for all others, and at least one b-tagged jet. Furthermore, events are re-

quired to have OS SF lepton pair with  $|m_{\ell\ell} - m_Z| < 10 \text{ GeV}$ , and the remaining lepton to have  $p_T > 35 \text{ GeV}$  to suppress processes coming from DY with a misidentified lepton. To maximize the discrimination of  $t\bar{t}Z$  from the remaining backgrounds, the events are binned in terms of  $n_{\text{jet}}$  and  $n_b$ , since  $t\bar{t}Z$  has higher multiplicities of both compared to the other backgrounds.

A summary of the DL VR and CRs can be found in the last three columns of Table 2.

Table 2: CRs defined for the main backgrounds of the AH SRs (first 4 columns,  $t\bar{t}(1\ell)$ ,  $W$ +jets,  $Z \rightarrow \ell\bar{\ell}$ , QCD), the SL SRs (central two columns,  $t\bar{t}(2\ell)$  and  $W$ +jets), and the DL SRs (last 2 columns,  $t\bar{t}(2\ell)$  and  $t\bar{t}Z$ ). To increase the event counts, some selection criteria applied in the SRs are removed in the related CRs and hence are not listed. The  $p_T^{\text{miss}}$  selection for the  $Z \rightarrow \ell\bar{\ell}$  CR refers to the hadronic recoil.

|                                                            | AH                   |                 |               |            | SL                   |                 | DL                   |            |                |
|------------------------------------------------------------|----------------------|-----------------|---------------|------------|----------------------|-----------------|----------------------|------------|----------------|
|                                                            | $t\bar{t}(1\ell)$ CR | $W(\ell\nu)$ CR | $Z(2\ell)$ CR | QCD CR     | $t\bar{t}(2\ell)$ CR | $W(\ell\nu)$ CR | $t\bar{t}(2\ell)$ VR | DY CR      | $t\bar{t}Z$ CR |
| $n_b$                                                      | $\geq 1$             | $= 0$           | $= 0$         | $\geq 1$   | $\geq 1$             | $= 0$           | $\geq 1$             | $= 1$      | $\geq 1$       |
| $n_{\text{lep}}$                                           | $= 1$                | $= 1$           | $= 2$         | $= 0$      | $= 2$                | $= 1$           | $= 2$                | $= 2$ (SF) | $= 3$          |
| $n_{\text{jet}}$                                           | $\geq 3$             | $\geq 3$        | $\geq 3$      | $\geq 3$   | $\geq 2$             | $\geq 2$        | $\geq 1$             | $\geq 1$   | $\geq 3$       |
| $p_T^{\text{miss}}$ [GeV]                                  | $\geq 250$           | $\geq 250$      | $\geq 250$    | $\geq 250$ | $\geq 250$           | $\geq 250$      | —                    | —          | —              |
| $M_T$ [GeV]                                                | $\leq 140$           | $\leq 140$      | —             | —          | —                    | $\geq 140$      | —                    | —          | —              |
| $\min \Delta\phi(j_{1,2}, \vec{p}_T^{\text{miss}})$ [rad.] | $\geq 0.8$           | —               | —             | $< 0.8$    | —                    | —               | —                    | —          | —              |
| $m_{\ell\ell}$ [GeV]                                       | —                    | —               | [60, 120]     | —          | —                    | —               | $> 20$               | —          | —              |
| $ m_{\ell\ell} - m_Z $ [GeV]                               | —                    | —               | —             | —          | —                    | —               | $> 15$ (SF)          | $< 15$     | $< 10$ (OS SF) |
| $m_{T2}^{\ell\ell}$ [GeV]                                  | —                    | —               | —             | —          | $\leq 80$            | $\leq 80$       | $\leq 80$            | $\geq 80$  | —              |
| Included in fit?                                           | Yes                  | Yes             | Yes           | Yes        | Yes                  | Yes             | No                   | Yes        | Yes            |

## 5 Systematic uncertainties

Different sources of systematic uncertainties are considered, in most cases affecting both the signal and the background. The various sources are implemented in the limit calculation, described in detail later. We distinguish between two types of uncertainties, those that only affect the normalization of a process, and those that in addition affect its distribution.

The following uncertainties correspond to constrained normalization nuisance parameters in the fit discussed in Section 6 (unless otherwise noted, the uncertainty source is applied to all channels):

- *Luminosity*: the integrated luminosities for the 2016, 2017, and 2018 data-taking years have 1.2–2.5% individual uncertainties [64–66].
- *Single top quark background normalization (AH and SL channels)*: an uncertainty of 20% in the normalization of single top quark processes is considered, accounting for the uncertainty due to differences in generator predictions and the  $t\bar{t}/tW$  interference treatment in the predicted cross section.

The following sources of uncertainty affect the distributions as well as the normalization of the signal and background processes. They are applied to all search channels:

- *PDF uncertainties*: uncertainties in the choice of PDF are estimated by reweighting the samples to the NNPDF 3.0 [51] replicas for the 2016 data set and the NNPDF 3.1 [52] hessian variations for the 2017 and 2018 data sets, which were then combined as described in Ref. [67].
- *Factorization and renormalization scales*: the uncertainties in the choice of the factorization and renormalization scale parameters are taken into account by applying a set of weights that represent a change of these scales by a factor of 2 or 0.5. For processes where the rate is estimated directly from data in a CR (such as  $W$ +jets,  $Z$ +jets, and

QCD), these uncertainties are not considered;

- *Parton shower modeling*: variations in the parton shower initial and final state radiation (ISR and FSR) scales are also considered by independently varying these up and down by a factor of 2;
- *Pileup modeling*: the systematic uncertainty in the pileup modeling is taken into account by varying the total inelastic cross section entering the calculation of pileup distributions in simulation by  $\pm 4.6\%$  [68];
- *Trigger*: separate uncertainties are considered for the triggers employed in each channel. These are estimated using cross-checks of the methods used to compute them from control samples in data;
- *Lepton reconstruction and selection*: scale factors are applied to the MC processes to mimic the measured reconstruction and selection efficiencies of leptons in data. The uncertainties associated with these scale factors are binned in  $p_T$  and  $\eta$  and are of the order of  $\sim 2\%$  for electrons and muons;
- *Jet energy scale and resolution*: reconstructed jet four-momenta in the simulation are varied according to the uncertainties in the jet energy scale, which are split into different uncorrelated sources. Additionally, the  $p_T$  of the jets is stochastically varied within the resolution of the detector. These uncertainties are coherently propagated to all variables, including  $p_T^{\text{miss}}$  [69];
- *$p_T^{\text{miss}}$  mismodeling*: the effects of varying jet uncertainties are propagated to the calculation of the  $p_T^{\text{miss}}$ . Additionally, any unclustered energy, mostly coming from calorimeter deposits not assigned to any object, is varied within the associated measurement uncertainties and the impact of such variations is propagated to  $p_T^{\text{miss}}$ ;
- *b tagging efficiency scale factors*: the b tagging and light-flavor quark/gluon jet mistag efficiency scale factors and the respective uncertainties are measured in independent control samples [31] and propagated to the analysis;
- *W/Z+heavy-flavor fraction (AH and SL channels)*: the uncertainty in the fraction of W/Z+heavy-flavor (HF) jets in W+jets and Z+jets events is taken into account. The fraction of W/Z+jets events in which the jets include heavy flavour jets is varied within 20% separately and independently for the one b-tagged and two or more b-tagged jets cases [70–73].
- *Simulation sample size*: uncertainties originating from the limited size of the simulated signal and background samples are included by allowing each kinematic bin used in the signal extraction to fluctuate independently within its statistical uncertainty following Ref. [74];
- *Uncertainty related to ECAL mistiming*: partial mistiming of signals in the forward regions of the ECAL endcaps led to a minor reduction in trigger efficiency. Simulations are corrected to reproduce the behavior of the data and the uncertainty in these corrections is propagated to the distributions used in the signal extraction;
- *Electroweak and QCD K factors (AH and SL channels)*: uncertainties in the LO→NLO K factors calculated for W+jets and Z+jets processes are included to compensate for missing higher-order corrections. For QCD processes, the uncertainties arise from variations related to the factorization and renormalization scales. For electroweak processes, the magnitude of the missing higher-order corrections is estimated by calculating the difference between applying and omitting the LO→NLO electroweak K factors.

- *b-tagged jet multiplicity normalization (AH and SL channels)*: an uncertainty of 5% is considered for the normalization scale factors applied to both W+jets and Z+jets processes in 2017 and 2018 to correct the b-tagged jet multiplicity in the AH and SL channels;
- *Top quark  $p_T$  reweighting (DL channel)*: differential measurements of the top quark  $p_T$  spectrum in  $t\bar{t}$  events [75] show that the measured  $p_T$  spectrum is softer than in NLO simulation. Comparisons with NNLO QCD simulation suggest this difference is mainly due to missing higher-order corrections. A  $p_T$ -dependent scale factor is therefore applied to correct the difference between NNLO and NLO simulation in the DL channel. This correction is not needed in the other channels because of the  $p_T^{\text{miss}}$ -dependent CR corrections factors. A corresponding systematic uncertainty is estimated by evaluating the difference between applying and omitting the reweighting.

The above sources of systematic uncertainties are considered with various degrees of correlation across the different channels and the data-taking periods. A group of nuisance parameters, such as the theory-related uncertainties, which include the variation in the factorization and renormalization scales, the PDF choice, the parton showering scales, the uncertainty in the inelastic cross section, and others, are treated as fully correlated across all channels and periods of data-taking. Analogously, for the channel-dependent higher-order theory corrections to W+jets, Z+jets, and  $t\bar{t}$  processes, a commonly associated parameter is assigned across all periods. Experimental sources of uncertainties such as the lepton identification and isolation efficiencies, as well as the efficiency of the different triggers used, are also considered fully correlated. Similarly, a subgroup of size equal to 13 nuisance parameters out of the 27 sub-sources affecting the jet energy scale and the parameter associated with the ECAL mistiming uncertainty are assumed to be maximally correlated across all analysis categories. Another group of nuisance parameters is considered in the fit with a partial correlation scheme. This includes, for instance, uncertainty sources such as b tagging efficiency, for which a fraction of sub-sources are considered fully correlated and another fraction completely uncorrelated across data-taking periods. The remainder (14 parameters) of the group of jet energy scale sub-sources are also treated as fully correlated, making the jet energy scale a partially correlated uncertainty source overall. The uncertainty in the integrated luminosity is naturally correlated across channels but is kept uncorrelated among different data-taking periods. On the other side, among the group of fully uncorrelated uncertainty sources across periods, one finds the nuisance parameters associated with jet energy resolution and the unclustered component of the missing transverse momentum. These two, however, because of their common nature, are kept correlated across the three channels investigated here.

## 6 Signal extraction

The DM signal is extracted from a simultaneous fit to the  $p_T^{\text{miss}}$  distributions in the AH and SL channels, and to the NN output distributions for the DL channel. The following results have been determined using the CMS statistical analysis tool COMBINE [76], which is based on the ROOFIT [77] and ROOSTATS [78] frameworks. The main SM backgrounds, discussed in Section 4, are dileptonic  $t\bar{t}$  and W+jets events for the SL channels, single-lepton  $t\bar{t}$ , W+jets, and  $Z \rightarrow \nu\bar{\nu}$  events for the AH channels, and dileptonic  $t\bar{t}$ ,  $tW$ , and  $t\bar{t}Z \rightarrow t\bar{t}\nu\bar{\nu}$  for the DL case.

The effect of the systematic uncertainties is taken into account by introducing nuisance parameters. Uncertainties that affect the normalization only are modeled using nuisances with log-normal probability densities. These parameters are treated as correlated between bins of

the fit distribution and between the different CRs and SRs within each channel for the same year. The common sources are correlated across channels, with the exception of the  $b$  tagging efficiency scale factors, which are decorrelated between the AH, SL, and DL channels because of a significantly different phase space.

To improve the estimation of the primary backgrounds in the SL and AH channels, an unconstrained multiplicative parameter is separately assigned to each background for each individual bin of the  $p_T^{\text{miss}}$  distribution. A thorough explanation of this method, typically referred to as the “transfer factor” method in many other DM searches, can be found in Ref. [79]. Concretely, the unconstrained parameters tend to adjust the expected contribution of the corresponding background process as a function of  $p_T^{\text{miss}}$  simultaneously in the SRs and CRs for a particular channel, taking advantage of the similar distributions of the fitting variables across different regions. For example, in any given  $p_T^{\text{miss}}$  bin of the SL channel, a single multiplicative parameter for  $t\bar{t}$  links the  $t\bar{t}$  background in the  $t\bar{t}$  enhanced  $2\ell$  CR, the  $W$ +jets enhanced  $1\ell$  CR, and the SRs. This has the advantage that the larger expected sample of  $t\bar{t}$  events in the enhanced  $2\ell$  CR provides the primary, though not exclusive, control of the necessary corrections to that given process in the SR by comparing to the observation in the signal-depleted region. Similarly, a multiplicative parameter links the rates of the DY process in each NN bin between the DY CR and the DL 1  $b$  tag SF SR, while an overall normalization parameter is used to link the rate of  $t\bar{t}Z$  between the  $t\bar{t}Z$  CR and the DL SRs. The rates of processes that do not have a dedicated CR in the fit are taken from simulation. Potential contributions from DM signals are included in all CRs and SRs, and scaled by a signal strength modifier  $\mu = \sigma/\sigma_{\text{th}}$ , i.e. the ratio between the measured and the theoretical cross sections.

A simultaneous fit to the binned  $p_T^{\text{miss}}$  and NN distributions is performed combining all of the aforementioned regions for each of the signal points considered. The results obtained for the fit to the 100 GeV scalar mediator (with best-fit signal strength  $0.51\sigma_{\text{th}}$ ) are illustrated in Figs. 2, 3, and 4 for the AH, SL, and DL channels, respectively. The lower panels show the ratio of data and the SM prediction after performing the maximum likelihood fit.

## 7 Results

The results are interpreted using the signal model for scalar and pseudoscalar mediators with masses ranging from 50 to 500 GeV, with  $m_\chi = 1$  GeV and  $g_q = g_\chi = 1$ . The results are presented in terms of the signal strength parameter  $\mu$ , defined as the ratio between the measured and the theoretical cross sections for the  $t/\bar{t}$ +DM and  $t\bar{t}$ +DM production modes summed together. The theoretical cross sections for both signal models are obtained at LO.

The uncertainties from the AH and SL channels are found to be similar and dominate the combination, since these channels are most sensitive to the signal. In particular, the leading uncertainties are the  $b$ -tagging scale factor uncertainties (which can affect the number of  $W$ +jets and  $Z \rightarrow \nu\bar{\nu}$  events entering the SRs, and the uncertainties from the per-bin background estimation parameters, particularly for the  $t\bar{t}$  process. The next most significant uncertainties principally affect the DL channel: the normalization uncertainties of the  $t\bar{t}Z$  process and the last bin of the DY process, followed by the jet energy resolution uncertainties, which are observed to affect the shape of the  $t\bar{t}$   $m_{T2}^{\ell\ell}$  distribution in particular.

For masses of the mediator particle below 200 (300) GeV for the scalar (pseudoscalar) model, the leading contribution to the signal sensitivity stems from  $t\bar{t}$ +DM processes because of their larger cross sections with respect to  $t/\bar{t}$ +DM production. However, the  $t/\bar{t}$ +DM cross section ( $p_T^{\text{miss}}$  distribution) drops less rapidly as a function of the mediator particle mass ( $p_T^{\text{miss}}$ ) in




Figure 2: The main discriminant distribution  $p_T^{\text{miss}}$  in the three AH SRs: 1b 0FJ (top left), 1b 1FJ (top right), and 2b (bottom), after a signal plus background fit across all channels, assuming a 100 GeV scalar mediator. The solid histograms for the simulated SM backgrounds are summed cumulatively and rescaled to luminosity, while the signal is shown as a dashed pink line. The grey dashed area in the upper panel represents the total uncertainty in the cumulative sum of the simulated SM backgrounds and signal, while in the lower panel it represents only the total uncertainty in the backgrounds. The data are represented by solid points with the horizontal bar indicating the width of the bin and the vertical one the associated statistical uncertainty. The last bin contains overflow events. The lower panel shows the ratio of data to post-fit background.




Figure 3: The main discriminant distribution  $p_T^{\text{miss}}$  in the six SL SRs: 1b 0FJ ( $t \leq 0$ ) (top left), 1b 0FJ ( $t > 0$ ) (top right), 1b 1FJ ( $t \leq 0$ ) (center left), 1b 1FJ ( $t > 0$ ) (center right), 2b ( $t \leq 0$ ) (bottom left), and 2b ( $t > 0$ ) (bottom right), after a signal plus background fit across all channels, assuming a 100 GeV scalar mediator. The solid histograms for the simulated SM backgrounds are summed cumulatively and rescaled to luminosity, while the signal is shown as a dashed pink line. The grey dashed area in the upper panel represents the total uncertainty in the cumulative sum of the simulated SM backgrounds and signal, while in the lower panel it represents only the total uncertainty in the backgrounds. The data are represented by solid points with the horizontal bar indicating the width of the bin and the vertical one the associated statistical uncertainty. The last bin contains overflow events. The lower panel shows the ratio of data to post-fit background.




Figure 4: The main discriminant distribution NN in the the four DL SRs: 2b (DF) (top left), 2b (SF) (top right), 1b (DF) (bottom left), and 1b (SF) (bottom right), after a signal plus background fit across all channels, assuming a 100 GeV scalar mediator. The solid histograms for the simulated SM backgrounds are summed cumulatively and rescaled to luminosity, while the signal is shown as a dashed pink line. The grey dashed area in the upper panel represents the total uncertainty in the cumulative sum of the simulated SM backgrounds and signal, while in the lower panel it represents only the total uncertainty in the backgrounds. The data are represented by solid points with the horizontal bar indicating the width of the bin and the vertical one the associated statistical uncertainty. The last bin contains overflow events. The lower panel shows the ratio of data to post-fit background.

comparison to the  $t\bar{t} + \text{DM}$ .

Upper limits at 95% confidence level (CL) are computed using a modified frequentist approach with a test statistic based on the profile likelihood in the asymptotic approximation and the  $\text{CL}_s$  criterion [80–82]. The results are shown in Fig. 5 in terms of model-independent 95% CL limits on the production cross section for new physics processes for the DM scalar (left) and pseudoscalar (right) models. Cross section values ranging from 0.02 to 1 pb are probed depending on the assumption of the mediator of each new phenomenon. In terms of the DM model considered, we expect to exclude mediator masses below 410 (380) GeV for the benchmark cross sections of the scalar (pseudoscalar) hypothesis.

A signal-like excess is observed in data. This excess is most visible in the AH 2b SR, and three of the DL SRs. The size of the excess is found to be statistically consistent across all SRs. Because the signal kinematic properties have limited correlations with the mass of the mediator, this excess is consistent with all mediator mass hypotheses, with local significances between 1.4 and 1.9 standard deviations. The largest significance is obtained for the 150 GeV pseudoscalar mediator. The variations in local significances are understood to come from the differing relative sensitivities of the lepton channels to the mediator mass hypotheses. In particular, the signal hypotheses with the lowest mediator masses (50 and 100 GeV) have slightly softer kinematic properties, which are less consistent with this excess. Because of the observed excess, we only exclude mediator masses below 310 (320) GeV for the scalar (pseudoscalar) mediator




Figure 5: The model-independent 95% CL limits on production cross section for new physics processes for the scalar (left) and pseudoscalar (right) interactions. The expected limit is shown by the black dashed line with the 68 and 95% CL uncertainty bands in green and yellow, respectively, while the observed limit is shown by the solid black line. Theoretical LO cross section values for the DM model and their associated uncertainties are also presented (grey line).

As discussed in Section 1, these signatures can also be interpreted in terms of models with an ALP mediator  $A$ . Since the additional pseudoscalar Higgs boson considered in Eq. (2) and an ALP as introduced in Eq. (3) exhibit the same coupling structure, if the ALP has zero or negligible couplings to the gluon or electroweak gauge bosons [83], the existing limits on the production cross section for pseudoscalar interactions as presented in Fig. 5 can be directly translated into upper limits on the ALP coupling to the top quark  $c_t$ . Figure 6 shows the resulting limits on the ratio of the top quark coupling  $c_t$  to the ALP scale  $f_A$ , as a function of the




Figure 6: The 95% CL limits on the ratio of ALP-top coupling to the ALP decay constant ( $|c_t|/f_A$ ) as a function of mediator mass for the ALP mediator model. The expected limit is shown by the black dashed line with the 68 and 95% CL uncertainty bands in green and yellow, respectively, whereas the observed limit is shown by the solid black line.

ALP mass  $m_A$ , assuming  $c_t = c_\chi$ . For ALP masses above 350 GeV, it is possible for the mediator to decay back to top quarks and hence varying  $c_t$  will change the branching ratio of the ALP to DM; to resolve this ambiguity we take the coupling to DM  $c_\chi$  equal to the coupling to top quarks  $c_t$ .

## 8 Summary

A search for dark matter (DM) produced in association with a single top quark or a top quark pair produced in interactions with a neutral scalar or pseudoscalar mediator in proton-proton collisions at a center-of-mass energy of 13 TeV has been presented. The search was performed using data corresponding to an integrated luminosity of  $138 \text{ fb}^{-1}$  recorded by the CMS experiment between 2016 and 2018. This is the first search simultaneously optimized for the  $t/\bar{t}$ +DM and  $t\bar{t}$ +DM phase spaces across the zero lepton, single-lepton, and two lepton final states. In particular, lower jet multiplicities were considered with respect to the best LHC  $t\bar{t}$ +DM searches, resulting in an increase in the sensitivity to processes where DM is produced in association with a single top quark.

The results are interpreted within a simplified model in which a scalar or pseudoscalar mediator couples to the top quark and subsequently decays into two DM particles. Scalar and pseudoscalar mediator masses below 410 and 380 GeV, respectively, are expected to be excluded at 95% confidence level (CL) assuming a DM particle mass of 1 GeV and mediator couplings to fermions and DM particles equal to unity. This work represents a 40% improvement in sensitivity with respect to the previous combined CMS search for  $t/\bar{t}$ +DM and  $t\bar{t}$ +DM, which was the best LHC result with 2016 data. Of these improvements, up to 20% comes from analysis enhancements and inclusion of the dileptonic final state, and 20% from the larger data sample. A small signal-like excess is observed in data. Because the signal kinematic properties do not significantly depend on the mass of the mediator, this excess is consistent with all mediator mass hypotheses. The largest local significance for all mediator hypotheses is observed to be within

two standard deviations. Because of this excess, mediator masses are only excluded below 310 (320) GeV for the scalar (pseudoscalar) mediator. The results are also translated into model-independent 95% CL upper limits on the visible cross section of DM production in association with top quarks, ranging from 1 pb to 0.02 pb.

In addition, limits on the coupling of axion-like-particles (ALP) to top quarks are set for the first time. This is performed in the context of top quark(s) plus invisible signatures where the ALP couples to SM quarks as a mediator between the SM and fermionic DM particles.

## Acknowledgments

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, RVTT3 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIP and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MCIN/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Turkey); NASU (Ukraine); STFC (United Kingdom); DOE and NSF (USA).

Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, contract Nos. 675440, 724704, 752730, 758316, 765710, 824093, 101115353, 101002207, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, project no. 22rl-037 (Armenia); the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Beijing Municipal Science & Technology Commission, No. Z191100007219010 and Fundamental Research Funds for the Central Universities (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, grant FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), among others, under Germany's Excellence Strategy – EXC 2121 “Quantum Universe” – 390833306, and under project number 400140256 - GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project Number 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program - ÚNKP, the NKFIH research grants K 131991, K 133046, K 138136, K 143460, K 143477, K 146913, K 146914, K 147048, 2020-2.2.1-ED-2021-00181, TKP2021-NKTA-64, and 2021-4.1.2-NEMZ\_KI-2024-00036 (Hungary); the Council of Science and Industrial Research, India; ICSC – National Research Center for High Performance Computing, Big Data and Quantum Computing and FAIR – Future Artificial Intelligence Research, funded by the NextGenerationEU pro-

gram (Italy); the Latvian Council of Science; the Ministry of Education and Science, project no. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, grant CEECIND/01334/2018 (Portugal); the National Priorities Research Program by Qatar National Research Fund; MCIN/AEI/10.13039/501100011033, ERDF “a way of making Europe”, and the Programa Estatal de Fomento de la Investigación Científica y Técnica de Excelencia María de Maeztu, grant MDM-2017-0765 and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, grant B39G670016 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, contract C-1845; and the Weston Havens Foundation (USA).

## References

- [1] T. A. Porter, R. P. Johnson, and P. W. Graham, “Dark matter searches with astroparticle data”, *Ann. Rev. Astron. Astrophys.* **49** (2011) 155, doi:10.1146/annurev-astro-081710-102528, arXiv:1104.2836.
- [2] G. Bertone, D. Hooper, and J. Silk, “Particle dark matter: evidence, candidates and constraints”, *Phys. Rept.* **405** (2005) 279, doi:10.1016/j.physrep.2004.08.031, arXiv:hep-ph/0404175.
- [3] D. Abercrombie et al., “Dark matter benchmark models for early LHC run-2 searches: Report of the ATLAS/CMS dark matter forum”, *Phys. Dark Univ.* **27** (2020) 100371, doi:10.1016/j.dark.2019.100371, arXiv:1507.00966.
- [4] M. R. Buckley, D. Feld, and D. Goncalves, “Scalar simplified models for dark matter”, *Phys. Rev. D* **91** (2015) doi:10.1103/physrevd.91.015017, arXiv:1410.6497.
- [5] D. Pinna, A. Zucchetta, M. R. Buckley, and F. Canelli, “Single top quarks and dark matter”, *Phys. Rev. D* **96** (2017) 035031, doi:10.1103/PhysRevD.96.035031, arXiv:1701.05195.
- [6] LHC Dark Matter Working Group Collaboration, “LHC dark matter working group: Next-generation spin-0 dark matter models”, *Phys. Dark Univ.* **27** (2020) 100351, doi:10.1016/j.dark.2019.100351, arXiv:1810.09420.
- [7] M. Bauer, U. Haisch, and F. Kahlhoefer, “Simplified dark matter models with two Higgs doublets: I. pseudoscalar mediators”, *JHEP* **05** (2017) 138, doi:10.1007/JHEP05(2017)138, arXiv:1701.07427.
- [8] R. D. Peccei and H. R. Quinn, “CP conservation in the presence of instantons”, *Phys. Rev. Lett.* **38** (1977) 1440, doi:10.1103/PhysRevLett.38.1440.
- [9] R. D. Peccei and H. R. Quinn, “Constraints imposed by CP conservation in the presence of instantons”, *Phys. Rev. D* **16** (1977) 1791, doi:10.1103/PhysRevD.16.1791.
- [10] S. Weinberg, “A new light boson?”, *Phys. Rev. Lett.* **40** (1978) 223, doi:10.1103/PhysRevLett.40.223.
- [11] F. Wilczek, “Problem of strong  $P$  and  $T$  invariance in the presence of instantons”, *Phys. Rev. Lett.* **40** (1978) 279, doi:10.1103/PhysRevLett.40.279.

- 
- [12] M. J. Dolan, F. Kahlhoefer, C. McCabe, and K. Schmidt-Hoberg, “A taste of dark matter: Flavour constraints on pseudoscalar mediators”, *JHEP* **03** (2015) 171, doi:10.1007/JHEP03(2015)171, arXiv:1412.5174. [Erratum: doi:10.1007/JHEP07(2015)103].
- [13] A. Bharucha, F. Brümmer, N. Desai, and S. Mutzel, “Axion-like particles as mediators for dark matter: beyond freeze-out”, *JHEP* **02** (2023) 141, doi:10.1007/JHEP02(2023)141, arXiv:2209.03932.
- [14] S. Allen et al., “Electroweak axion portal to dark matter”, 2024. arXiv:2405.02403.
- [15] S. Blasi et al., “Top-philic ALP phenomenology at the LHC: the elusive mass-window”, *JHEP* **06** (2024) 077, doi:10.1007/JHEP06(2024)077, arXiv:2311.16048.
- [16] CMS Collaboration, “Combined searches for the production of supersymmetric top quark partners in proton–proton collisions at  $\sqrt{s} = 13$  TeV”, *Eur. Phys. J. C* **81** (2021) 970, doi:10.1140/epjc/s10052-021-09721-5, arXiv:2107.10892.
- [17] ATLAS Collaboration, “Constraints on spin-0 dark matter mediators and invisible Higgs decays using ATLAS 13 TeV pp collision data with two top quarks and missing transverse momentum in the final state”, *Eur. Phys. J. C* **83** (2023) 503, doi:10.1140/epjc/s10052-023-11477-z, arXiv:2211.05426.
- [18] ATLAS Collaboration, “Search for new phenomena with top-quark pairs and large missing transverse momentum using 140 fb<sup>-1</sup> of pp collision data at  $\sqrt{s} = 13$  TeV with the ATLAS detector”, *JHEP* **03** (2024) 139, doi:10.1007/JHEP03(2024)139, arXiv:2401.13430.
- [19] ATLAS Collaboration, “Search for dark matter produced in association with a single top quark and an energetic W boson in  $\sqrt{s} = 13$  TeV pp collisions with the ATLAS detector”, *Eur. Phys. J. C* **83** (2023), no. 7, 603, doi:10.1140/epjc/s10052-023-11582-z, arXiv:2211.13138.
- [20] ATLAS Collaboration, “Constraints on simplified dark matter models involving an s-channel mediator with the ATLAS detector in pp collisions at  $\sqrt{s} = 13$  TeV”, *Eur. Phys. J. C* **84** (2024), no. 10, 1102, doi:10.1140/epjc/s10052-024-13215-5, arXiv:2404.15930.
- [21] CMS Collaboration, “Search for dark matter produced in association with a single top quark or a top quark pair in proton-proton collisions at  $\sqrt{s} = 13$  TeV”, *JHEP* **03** (2019) 141, doi:10.1007/JHEP03(2019)141, arXiv:1901.01553.
- [22] HEPData record for this analysis, 2024. doi:10.17182/hepdata.154755.
- [23] CMS Collaboration, “The CMS experiment at the CERN LHC”, *JINST* **3** (2008) S08004, doi:10.1088/1748-0221/3/08/S08004.
- [24] CMS Collaboration, “The CMS trigger system”, *JINST* **12** (2017) P01020, doi:10.1088/1748-0221/12/01/P01020, arXiv:1609.02366.
- [25] CMS Collaboration, “Particle-flow reconstruction and global event description with the CMS detector”, *JINST* **12** (2017) P10003, doi:10.1088/1748-0221/12/10/P10003, arXiv:1706.04965.

- [26] CMS Collaboration, “Technical proposal for the Phase-II upgrade of the Compact Muon Solenoid”, CMS Technical Proposal CERN-LHCC-2015-010, CMS-TDR-15-02, 2015.
- [27] M. Cacciari, G. P. Salam, and G. Soyez, “The anti- $k_T$  jet clustering algorithm”, *JHEP* **04** (2008) 063, doi:10.1088/1126-6708/2008/04/063, arXiv:0802.1189.
- [28] M. Cacciari, G. P. Salam, and G. Soyez, “FastJet user manual”, *Eur. Phys. J. C* **72** (2012) 1896, doi:10.1140/epjc/s10052-012-1896-2, arXiv:1111.6097.
- [29] M. Cacciari, G. P. Salam, and G. Soyez, “The catchment area of jets”, *JHEP* **04** (2008) 005, doi:10.1088/1126-6708/2008/04/005, arXiv:0802.1188.
- [30] CMS Collaboration, “Jet algorithms performance in 13 TeV data”, CMS Physics Analysis Summary CMS-PAS-JME-16-003, 2017.
- [31] CMS Collaboration, “Identification of heavy-flavour jets with the CMS detector in pp collisions at 13 TeV”, *JINST* **13** (2018) P05011, doi:10.1088/1748-0221/13/05/P05011, arXiv:1712.07158.
- [32] P. Nason, “A new method for combining NLO QCD with shower Monte Carlo algorithms”, *JHEP* **11** (2004) 040, doi:10.1088/1126-6708/2004/11/040, arXiv:hep-ph/0409146.
- [33] S. Frixione, P. Nason, and C. Oleari, “Matching NLO QCD computations with parton shower simulations: the POWHEG method”, *JHEP* **11** (2007) 070, doi:10.1088/1126-6708/2007/11/070, arXiv:0709.2092.
- [34] S. Alioli, P. Nason, C. Oleari, and E. Re, “A general framework for implementing NLO calculations in shower Monte Carlo programs: the POWHEG box”, *JHEP* **06** (2010) 043, doi:10.1007/JHEP06(2010)043, arXiv:1002.2581.
- [35] S. Alioli, P. Nason, C. Oleari, and E. Re, “NLO single-top production matched with shower in POWHEG: s- and t-channel contributions”, *JHEP* **09** (2009) 111, doi:10.1088/1126-6708/2009/09/111, arXiv:0907.4076. [Erratum: doi:10.1007/JHEP02(2010)011].
- [36] E. Re, “Single-top Wt-channel production matched with parton showers using the POWHEG method”, *Eur. Phys. J. C* **71** (2011) 1547, doi:10.1140/epjc/s10052-011-1547-z, arXiv:1009.2450.
- [37] S. Frixione, P. Nason, and G. Ridolfi, “A positive-weight next-to-leading-order Monte Carlo for heavy flavour hadroproduction”, *JHEP* **09** (2007) 126, doi:10.1088/1126-6708/2007/09/126, arXiv:0707.3088.
- [38] M. Czakon et al., “Top-pair production at the LHC through NNLO QCD and NLO EW”, *JHEP* **10** (2017) 186, doi:10.1007/JHEP10(2017)186, arXiv:1705.04105.
- [39] R. Frederix and S. Frixione, “Merging meets matching in MC@NLO”, *JHEP* **12** (2012) 061, doi:10.1007/JHEP12(2012)061, arXiv:1209.6215.
- [40] J. Alwall et al., “The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations”, *JHEP* **07** (2014) 079, doi:10.1007/JHEP07(2014)079, arXiv:1405.0301.

- 
- [41] M. L. Mangano, M. Moretti, F. Piccinini, and M. Treccani, “Matching matrix elements and shower evolution for top-quark production in hadronic collisions”, *JHEP* **01** (2007) 013, doi:10.1088/1126-6708/2007/01/013, arXiv:hep-ph/0611129.
  - [42] A. Denner, S. Dittmaier, T. Kasprzik, and A. Muck, “Electroweak corrections to W+jet hadroproduction including leptonic W-boson decays”, *JHEP* **08** (2009) 075, doi:10.1088/1126-6708/2009/08/075, arXiv:0906.1656.
  - [43] A. Denner, S. Dittmaier, T. Kasprzik, and A. Muck, “Electroweak corrections to dilepton+jet production at hadron colliders”, *JHEP* **06** (2011) 069, doi:10.1007/JHEP06(2011)069, arXiv:1103.0914.
  - [44] A. Denner, S. Dittmaier, T. Kasprzik, and A. Maeck, “Electroweak corrections to monojet production at the LHC”, *Eur. Phys. J. C* **73** (2013) 2297, doi:10.1140/epjc/s10052-013-2297-x, arXiv:1211.5078.
  - [45] J. H. Kuhn, A. Kulesza, S. Pozzorini, and M. Schulze, “Electroweak corrections to hadronic photon production at large transverse momenta”, *JHEP* **03** (2006) 059, doi:10.1088/1126-6708/2006/03/059, arXiv:hep-ph/0508253.
  - [46] S. Kallweit et al., “NLO electroweak automation and precise predictions for W+multijet production at the LHC”, *JHEP* **04** (2015) 012, doi:10.1007/JHEP04(2015)012, arXiv:1412.5157.
  - [47] S. Kallweit et al., “NLO QCD+EW predictions for V+jets including off-shell vector-boson decays and multijet merging”, *JHEP* **04** (2016) 021, doi:10.1007/JHEP04(2016)021, arXiv:1511.08692.
  - [48] T. Gehrmann et al., “ $W^+W^-$  production at hadron colliders in next to next to leading order QCD”, *Phys. Rev. Lett.* **113** (2014) 212001, doi:10.1103/PhysRevLett.113.212001, arXiv:1408.5243.
  - [49] J. M. Campbell and R. K. Ellis, “An update on vector boson pair production at hadron colliders”, *Phys. Rev. D* **60** (1999) 113006, doi:10.1103/PhysRevD.60.113006, arXiv:hep-ph/9905386.
  - [50] A. Boveia et al., “Recommendations on presenting LHC searches for missing transverse energy signals using simplified s-channel models of dark matter”, *Phys. Dark Univ.* **27** (2020) 100365, doi:10.1016/j.dark.2019.100365, arXiv:1603.04156.
  - [51] NNPDF Collaboration, “Parton distributions for the LHC Run II”, *JHEP* **04** (2015) 040, doi:10.1007/JHEP04(2015)040, arXiv:1410.8849.
  - [52] NNPDF Collaboration, “Parton distributions from high-precision collider data”, *Eur. Phys. J. C* **77** (2017) 663, doi:10.1140/epjc/s10052-017-5199-5, arXiv:1706.00428.
  - [53] T. Sjöstrand et al., “An introduction to PYTHIA 8.2”, *Comput. Phys. Commun.* **191** (2015) 159, doi:10.1016/j.cpc.2015.01.024, arXiv:1410.3012.
  - [54] CMS Collaboration, “Event generator tunes obtained from underlying event and multiparton scattering measurements”, *Eur. Phys. J. C* **76** (2016) 155, doi:10.1140/epjc/s10052-016-3988-x, arXiv:1512.00815.

- [55] CMS Collaboration, “Extraction and validation of a new set of CMS PYTHIA8 tunes from underlying-event measurements”, *Eur. Phys. J. C* **80** (2020) 4, doi:10.1140/epjc/s10052-019-7499-4, arXiv:1903.12179.
- [56] GEANT4 Collaboration, “GEANT4—a simulation toolkit”, *Nucl. Instrum. Meth. A* **506** (2003) 250, doi:10.1016/S0168-9002(03)01368-8.
- [57] CMS Collaboration, “Performance of electron reconstruction and selection with the CMS detector in proton-proton collisions at  $\sqrt{s} = 8$  TeV”, *JINST* **10** (2015) P06005, doi:10.1088/1748-0221/10/06/P06005, arXiv:1502.02701.
- [58] CMS Collaboration, “Performance of CMS muon reconstruction in pp collision events at  $\sqrt{s} = 7$  TeV”, *JINST* **7** (2012) P10002, doi:10.1088/1748-0221/7/10/P10002, arXiv:1206.4071.
- [59] Y. Bai, H.-C. Cheng, J. Gallicchio, and J. Gu, “Stop the top background of the stop search”, *JHEP* **07** (2012) 110, doi:10.1007/JHEP07(2012)110, arXiv:1203.4813.
- [60] CMS Collaboration, “Search for direct top squark pair production in events with one lepton, jets, and missing transverse momentum at 13 TeV with the CMS experiment”, *JHEP* **05** (2020) 032, doi:10.1007/JHEP05(2020)032, arXiv:1912.08887.
- [61] C. G. Lester and D. J. Summers, “Measuring masses of semiinvisibly decaying particles pair produced at hadron colliders”, *Phys. Lett. B* **463** (1999) 99, doi:10.1016/S0370-2693(99)00945-4, arXiv:hep-ph/9906349.
- [62] CMS Collaboration, “Searches for supersymmetry using the  $m_{T2}$  variable in hadronic events produced in pp collisions at 8 TeV”, *JHEP* **05** (2015) 078, doi:10.1007/JHEP05(2015)078, arXiv:1502.04358.
- [63] B. A. Betchart, R. Demina, and A. Harel, “Analytic solutions for neutrino momenta in decay of top quarks”, *Nucl. Instrum. Meth. A* **736** (2014) 169, doi:10.1016/j.nima.2013.10.039, arXiv:1305.1878.
- [64] CMS Collaboration, “Precision luminosity measurement in proton-proton collisions at  $\sqrt{s} = 13$  TeV in 2015 and 2016 at CMS”, *Eur. Phys. J. C* **81** (2021) 800, doi:10.1140/epjc/s10052-021-09538-2, arXiv:2104.01927.
- [65] CMS Collaboration, “CMS luminosity measurement for the 2017 data-taking period at  $\sqrt{s} = 13$  TeV”, technical report, CERN, Geneva, 2018.
- [66] CMS Collaboration, “CMS luminosity measurement for the 2018 data-taking period at  $\sqrt{s} = 13$  TeV”, technical report, CERN, Geneva, 2019.
- [67] J. Butterworth et al., “PDF4LHC recommendations for LHC Run II”, *J. Phys. G* **43** (2016) 023001, doi:10.1088/0954-3899/43/2/023001, arXiv:1510.03865.
- [68] CMS Collaboration, “Measurement of the inelastic proton-proton cross section at  $\sqrt{s} = 13$  TeV”, *JHEP* **07** (2018) 161, doi:10.1007/JHEP07(2018)161, arXiv:1802.02613.
- [69] CMS Collaboration, “Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV”, *JINST* **12** (2017) P02014, doi:10.1088/1748-0221/12/02/P02014, arXiv:1607.03663.

- 
- [70] CMS Collaboration, “Measurements of differential cross sections for associated production of a W boson and jets in proton-proton collisions at  $\sqrt{s} = 8$  TeV”, *Phys. Rev. D* **95** (2017) 052002, doi:10.1103/PhysRevD.95.052002, arXiv:1610.04222.
  - [71] CMS Collaboration, “Measurement of the production cross section of a W boson in association with two b jets in pp collisions at  $\sqrt{s} = 8$  TeV”, *Eur. Phys. J. C* **77** (2017), no. 2, 92, doi:10.1140/epjc/s10052-016-4573-z, arXiv:1608.07561.
  - [72] CMS Collaboration, “Measurements of jet multiplicity and differential production cross sections of Z+jets events in proton-proton collisions at  $\sqrt{s} = 7$  TeV”, *Phys. Rev. D* **91** (2015), no. 5, 052008, doi:10.1103/PhysRevD.91.052008, arXiv:1408.3104.
  - [73] CMS Collaboration, “Measurements of the associated production of a Z boson and b jets in pp collisions at  $\sqrt{s} = 8$  TeV”, *Eur. Phys. J. C* **77** (2017), no. 11, 751, doi:10.1140/epjc/s10052-017-5140-y, arXiv:1611.06507.
  - [74] R. J. Barlow and C. Beeston, “Fitting using finite Monte Carlo samples”, *Comput. Phys. Commun.* **77** (1993) 219, doi:10.1016/0010-4655(93)90005-W.
  - [75] CMS Collaboration, “Measurement of differential cross sections for top quark pair production using the lepton+jets final state in proton-proton collisions at 13 TeV”, *Phys. Rev. D* **95** (2017) 092001, doi:10.1103/PhysRevD.95.092001, arXiv:1610.04191.
  - [76] CMS Collaboration, “The CMS statistical analysis and combination tool: COMBINE”, *Comput. Softw. Big Sci.* **8** (2024) 19, doi:10.1007/s41781-024-00121-4, arXiv:2404.06614.
  - [77] W. Verkerke and D. Kirkby, “The ROOFIT toolkit for data modeling”, in *Proc. 13th International Conference on Computing in High Energy and Nuclear Physics (CHEP 2003): La Jolla CA, United States, March 24–28, 2003*. 2003. arXiv:physics/0306116. [eConf C0303241 (2003) MOLT007].
  - [78] L. Moneta et al., “The RooStats project”, in *13th International Workshop on Advanced Computing and Analysis Techniques in Physics Research (ACAT2010)*, T. Speer et al., eds. SISSA, 2010. arXiv:1009.1003. PoS(ACAT2010)057. doi:10.22323/1.093.0057.
  - [79] “Dark sector searches with the CMS experiment”, *Physics Reports* **1115** (2025) 448, doi:10.1016/j.physrep.2024.09.013. Included in the special issue “ATLAS and CMS Run II Legacy results”.
  - [80] T. Junk, “Confidence level computation for combining searches with small statistics”, *Nucl. Inst. Meth. A* **434** (1999) 435, doi:10.1016/S0168-9002(99)00498-2, arXiv:hep-ex/9902226.
  - [81] A. L. Read, “Presentation of search results: the CL<sub>s</sub> technique”, *J. Phys. G* **28** (2002) 2693, doi:10.1088/0954-3899/28/10/313.
  - [82] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, “Asymptotic formulae for likelihood-based tests of new physics”, *Eur. Phys. J. C* **71** (2011) 1554, doi:10.1140/epjc/s10052-011-1554-0, arXiv:1007.1727. [Erratum: doi:10.1140/epjc/s10052-013-2501-z].
  - [83] A. Anuar et al., “ALP-ine quests at the LHC: hunting axion-like particles via peaks and dips in  $t\bar{t}$  production”, 2024. arXiv:2404.19014.

## A The CMS Collaboration

### Yerevan Physics Institute, Yerevan, Armenia

V. Chekhovsky, A. Hayrapetyan, V. Makarenko , A. Tumasyan<sup>1</sup> 

### Institut für Hochenergiephysik, Vienna, Austria

W. Adam , J.W. Andrejkovic, L. Benato , T. Bergauer , S. Chatterjee , K. Damanakis , M. Dragicevic , P.S. Hussain , M. Jeitler<sup>2</sup> , N. Krammer , A. Li , D. Liko , I. Mikulec , J. Schieck<sup>2</sup> , R. Schöfbeck<sup>2</sup> , D. Schwarz , M. Sonawane , W. Waltenberger , C.-E. Wulz<sup>2</sup> 

### Universiteit Antwerpen, Antwerpen, Belgium

T. Janssen , H. Kwon , T. Van Laer, P. Van Mechelen 

### Vrije Universiteit Brussel, Brussel, Belgium

N. Breugelmans, J. D'Hondt , S. Dansana , A. De Moor , M. Delcourt , F. Heyen, Y. Hong , S. Lowette , I. Makarenko , D. Müller , S. Tavernier , M. Tytgat<sup>3</sup> , G.P. Van Onsem , S. Van Putte , D. Vannerom 

### Université Libre de Bruxelles, Bruxelles, Belgium

B. Bilin , B. Clerbaux , A.K. Das, I. De Bruyn , G. De Lentdecker , H. Evard , L. Favart , P. Gianneios , A. Khalilzadeh, F.A. Khan , K. Lee , A. Malara , M.A. Shahzad, L. Thomas , M. Vanden Bemden , C. Vander Velde , P. Vanlaer 

### Ghent University, Ghent, Belgium

M. De Coen , D. Dobur , G. Gokbulut , J. Knolle , L. Lambrecht , D. Marckx , K. Skovpen , N. Van Den Bossche , J. van der Linden , L. Wezenbeek 

### Université Catholique de Louvain, Louvain-la-Neuve, Belgium

S. Bein , A. Benecke , A. Bethani , G. Bruno , C. Caputo , J. De Favereau De Jeneret , C. Delaere , I.S. Donertas , A. Giammanco , A.O. Guzel , Sa. Jain , V. Lemaitre, J. Lidrych , P. Mastrapasqua , T.T. Tran , S. Turkcapar 

### Centro Brasileiro de Pesquisas Fisicas, Rio de Janeiro, Brazil

G.A. Alves , E. Coelho , G. Correia Silva , C. Hensel , T. Menezes De Oliveira , C. Mora Herrera<sup>4</sup> , P. Rebello Teles , M. Soeiro, E.J. Tonelli Manganote<sup>5</sup> , A. Vilela Pereira<sup>4</sup> 

### Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

W.L. Aldá Júnior , M. Barroso Ferreira Filho , H. Brandao Malbouisson , W. Carvalho , J. Chinellato<sup>6</sup>, E.M. Da Costa , G.G. Da Silveira<sup>7</sup> , D. De Jesus Damiao , S. Fonseca De Souza , R. Gomes De Souza, T. Laux Kuhn<sup>7</sup> , M. Macedo , J. Martins , K. Mota Amarilo , L. Mundim , H. Nogima , J.P. Pinheiro , A. Santoro , A. Sznajder , M. Thiel 

### Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil

C.A. Bernardes<sup>7</sup> , L. Calligaris , T.R. Fernandez Perez Tomei , E.M. Gregores , I. Maietto Silverio , P.G. Mercadante , S.F. Novaes , B. Orzari , Sandra S. Padula 

### Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria

A. Aleksandrov , G. Antchev , R. Hadjiiska , P. Iaydjiev , M. Misheva , M. Shopova , G. Sultanov 

### University of Sofia, Sofia, Bulgaria

A. Dimitrov , L. Litov , B. Pavlov , P. Petkov , A. Petrov , E. Shumka 

**Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile**

S. Keshri , D. Laroze , S. Thakur 

**Beihang University, Beijing, China**

T. Cheng , T. Javaid , L. Yuan 

**Department of Physics, Tsinghua University, Beijing, China**

Z. Hu , Z. Liang, J. Liu

**Institute of High Energy Physics, Beijing, China**

G.M. Chen<sup>8</sup> , H.S. Chen<sup>8</sup> , M. Chen<sup>8</sup> , F. Iemmi , C.H. Jiang, A. Kapoor<sup>9</sup> , H. Liao , Z.-A. Liu<sup>10</sup> , R. Sharma<sup>11</sup> , J.N. Song<sup>10</sup>, J. Tao , C. Wang<sup>8</sup>, J. Wang , Z. Wang<sup>8</sup>, H. Zhang , J. Zhao 

**State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China**

A. Agapitos , Y. Ban , A. Carvalho Antunes De Oliveira , S. Deng , B. Guo, C. Jiang , A. Levin , C. Li , Q. Li , Y. Mao, S. Qian, S.J. Qian , X. Qin, X. Sun , D. Wang , H. Yang, Y. Zhao, C. Zhou 

**Guangdong Provincial Key Laboratory of Nuclear Science and Guangdong-Hong Kong Joint Laboratory of Quantum Matter, South China Normal University, Guangzhou, China**

S. Yang 

**Sun Yat-Sen University, Guangzhou, China**

Z. You 

**University of Science and Technology of China, Hefei, China**

K. Jaffel , N. Lu 

**Nanjing Normal University, Nanjing, China**

G. Bauer<sup>12</sup>, B. Li<sup>13</sup>, H. Wang , K. Yi<sup>14</sup> , J. Zhang 

**Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE) - Fudan University, Shanghai, China**

Y. Li

**Zhejiang University, Hangzhou, Zhejiang, China**

Z. Lin , C. Lu , M. Xiao 

**Universidad de Los Andes, Bogota, Colombia**

C. Avila , D.A. Barbosa Trujillo, A. Cabrera , C. Florez , J. Fraga , J.A. Reyes Vega

**Universidad de Antioquia, Medellin, Colombia**

J. Jaramillo , C. Rendón , M. Rodriguez , A.A. Ruales Barbosa , J.D. Ruiz Alvarez 

**University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia**

D. Giljanovic , N. Godinovic , D. Lelas , A. Sculac 

**University of Split, Faculty of Science, Split, Croatia**

M. Kovac , A. Petkovic , T. Sculac 

**Institute Rudjer Boskovic, Zagreb, Croatia**

P. Bargassa , V. Brigljevic , B.K. Chitroda , D. Ferencek , K. Jakovcic, A. Starodumov<sup>15</sup> , T. Susa 

**University of Cyprus, Nicosia, Cyprus**

A. Attikis , K. Christoforou , A. Hadjiagapiou, C. Leonidou , J. Mousa , C. Nicolaou, L. Paizanos, F. Ptochos , P.A. Razis , H. Rykaczewski, H. Saka , A. Stepennov 

**Charles University, Prague, Czech Republic**

M. Finger , M. Finger Jr. , A. Kveton 

**Escuela Politecnica Nacional, Quito, Ecuador**

E. Ayala 

**Universidad San Francisco de Quito, Quito, Ecuador**

E. Carrera Jarrin 

**Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt**

A.A. Abdelalim<sup>16,17</sup> , S. Elgammal<sup>18</sup>, A. Ellithi Kamel<sup>19</sup>

**Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt**

M. Abdullah Al-Mashad , M.A. Mahmoud 

**National Institute of Chemical Physics and Biophysics, Tallinn, Estonia**

K. Ehataht , M. Kadastik, T. Lange , C. Nielsen , J. Pata , M. Raidal , L. Tani , C. Veelken 

**Department of Physics, University of Helsinki, Helsinki, Finland**

K. Osterberg , M. Voutilainen 

**Helsinki Institute of Physics, Helsinki, Finland**

N. Bin Norjoharuddeen , E. Brücken , F. Garcia , P. Inkaew , K.T.S. Kallonen , T. Lampén , K. Lassila-Perini , S. Lehti , T. Lindén , M. Myllymäki , M.m. Rantanen , J. Tuominiemi 

**Lappeenranta-Lahti University of Technology, Lappeenranta, Finland**

H. Kirschenmann , P. Luukka , H. Petrow 

**IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France**

M. Besancon , F. Couderc , M. Dejardin , D. Denegri, J.L. Faure, F. Ferri , S. Ganjour , P. Gras , G. Hamel de Monchenault , M. Kumar , V. Lohezic , J. Malcles , F. Orlandi , L. Portales , A. Rosowsky , M.Ö. Sahin , A. Savoy-Navarro<sup>20</sup> , P. Simkina , M. Titov , M. Tornago 

**Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France**

F. Beaudette , G. Boldrini , P. Busson , A. Cappati , C. Charlot , M. Chiusi , T.D. Cuisset , F. Damas , O. Davignon , A. De Wit , I.T. Ehle , B.A. Fontana Santos Alves , S. Ghosh , A. Gilbert , R. Granier de Cassagnac , A. Hakimi , B. Harikrishnan , L. Kalipoliti , G. Liu , M. Nguyen , S. Obraztsov , C. Ochando , R. Salerno , J.B. Sauvan , Y. Sirois , G. Sokmen, L. Urda Gómez , E. Vernazza , A. Zabi , A. Zghiche 

**Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France**

J.-L. Agram<sup>21</sup> , J. Andrea , D. Apparú , D. Bloch , J.-M. Brom , E.C. Chabert , C. Collard , S. Falke , U. Goerlach , R. Haeberle , A.-C. Le Bihan , M. Meena , O. Poncet , G. Saha , M.A. Sessini , P. Van Hove , P. Vaucelle 

**Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules,**

**CNRS/IN2P3, Villeurbanne, France**A. Di Florio **Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France**

D. Amram , S. Beauceron , B. Blancon , G. Boudoul , N. Chanon , D. Contardo ,  
 P. Depasse , C. Dozen<sup>22</sup> , H. El Mamouni , J. Fay , S. Gascon , M. Gouzevitch ,  
 C. Greenberg , G. Grenier , B. Ille , E. Jourdhuy , I.B. Laktineh , M. Lethuillier ,  
 L. Mirabito , S. Perries , A. Purohit , M. Vander Donckt , P. Verdier , J. Xiao 

**Georgian Technical University, Tbilisi, Georgia**A. Khvedelidze<sup>23</sup> , I. Lomidze , Z. Tsamalaidze<sup>23</sup> **RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany**

V. Botta , S. Consuegra Rodríguez , L. Feld , K. Klein , M. Lipinski , D. Meuser ,  
 A. Pauls , D. Pérez Adán , N. Röwert , M. Teroerde 

**RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany**

S. Diekmann , A. Dodonova , N. Eich , D. Eliseev , F. Engelke , J. Erdmann , M. Erdmann , P. Fackeldey , B. Fischer , T. Hebbeker , K. Hoepfner , F. Ivone , A. Jung ,  
 M.y. Lee , F. Mausolf , M. Merschmeyer , A. Meyer , S. Mukherjee , D. Noll ,  
 F. Nowotny , A. Pozdnyakov , Y. Rath , W. Redjeb , F. Rehm , H. Reithler , V. Sarkisovi ,  
 A. Schmidt , C. Seth , A. Sharma , J.L. Spah , F. Torres Da Silva De Araujo<sup>24</sup> ,  
 S. Wiedenbeck , S. Zaleski 

**RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany**

C. Dziwok , G. Flügge , T. Kress , A. Nowack , O. Pooth , A. Stahl , T. Ziemons ,  
 A. Zotz 

**Deutsches Elektronen-Synchrotron, Hamburg, Germany**

H. Aarup Petersen , M. Aldaya Martin , J. Alimena , S. Amoroso , Y. An , J. Bach ,  
 S. Baxter , M. Bayatmakou , H. Becerril Gonzalez , O. Behnke , A. Belvedere ,  
 F. Blekman<sup>25</sup> , K. Borras<sup>26</sup> , A. Campbell , A. Cardini , F. Colombina , M. De Silva ,  
 G. Eckerlin , D. Eckstein , L.I. Estevez Banos , E. Gallo<sup>25</sup> , A. Geiser , V. Guglielmi ,  
 M. Guthoff , A. Hinzmann , L. Jeppe , B. Kaech , M. Kasemann , C. Kleinwort ,  
 R. Kogler , M. Komm , D. Krücker , W. Lange , D. Leyva Pernia , K. Lipka<sup>27</sup> ,  
 W. Lohmann<sup>28</sup> , F. Lorkowski , R. Mankel , I.-A. Melzer-Pellmann , M. Mendizabal Morentin ,  
 A.B. Meyer , G. Milella , K. Moral Figueroa , A. Mussgiller ,  
 L.P. Nair , J. Niedziela , A. Nürnberg , J. Park , E. Ranken , A. Raspereza ,  
 D. Rastorguev , J. Rübenach , L. Rygaard , M. Scham<sup>29,26</sup> , S. Schnake<sup>26</sup> , P. Schütze ,  
 C. Schwanenberger<sup>25</sup> , D. Selivanova , K. Sharko , M. Shchedrolosiev , D. Stafford ,  
 F. Vazzoler , A. Ventura Barroso , R. Walsh , D. Wang , Q. Wang , K. Wichmann ,  
 L. Wiens<sup>26</sup> , C. Wissing , Y. Yang , S. Zakharov , A. Zimmermann Castro Santos 

**University of Hamburg, Hamburg, Germany**

A. Albrecht , S. Albrecht , M. Antonello , S. Bollweg , M. Bonanomi , P. Connor ,  
 K. El Morabit , Y. Fischer , E. Garutti , A. Grohsjean , J. Haller , D. Hundhausen ,  
 H.R. Jabusch , G. Kasieczka , P. Keicher , R. Klanner , W. Korcari , T. Kramer ,  
 C.c. Kuo , V. Kutzner , F. Labe , J. Lange , A. Lobanov , C. Matthies , L. Moureaux ,  
 M. Mrowietz , A. Nigamova , Y. Nissan , A. Paasch , K.J. Pena Rodriguez , T. Quadfasel ,  
 B. Raciti , M. Rieger , D. Savoie , J. Schindler , P. Schleper , M. Schröder ,  
 J. Schwandt , M. Sommerhalder , H. Stadie , G. Steinbrück , A. Tews , B. Wiederspan ,  
 M. Wolf 

**Karlsruher Institut fuer Technologie, Karlsruhe, Germany**

S. Brommer , E. Butz , T. Chwalek , A. Dierlamm , G.G. Dincer , U. Elicabuk, N. Faltermann , M. Giffels , A. Gottmann , F. Hartmann<sup>30</sup> , R. Hofsaess , M. Horzela , U. Husemann , J. Kieseler , M. Klute , O. Lavoryk , J.M. Lawhorn , M. Link, A. Lintuluoto , S. Maier , S. Mitra , M. Mormile , Th. Müller , M. Neukum, M. Oh , E. Pfeffer , M. Presilla , G. Quast , K. Rabbertz , B. Regnery , N. Shadskiy , I. Shvetsov , H.J. Simonis , L. Sowa, L. Stockmeier, K. Tauqeer, M. Toms , B. Topko , N. Trevisani , R.F. Von Cube , M. Wassmer , S. Wieland , F. Wittig, R. Wolf , X. Zuo 

**Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece**

G. Anagnostou, G. Daskalakis , A. Kyriakis , A. Papadopoulos<sup>30</sup>, A. Stakia 

**National and Kapodistrian University of Athens, Athens, Greece**

G. Melachroinos, Z. Painesis , I. Paraskevas , N. Saoulidou , K. Theofilatos , E. Tziaferi , K. Vellidis , I. Zisopoulos 

**National Technical University of Athens, Athens, Greece**

G. Bakas , T. Chatzistavrou, G. Karapostoli , K. Kousouris , I. Papakrivopoulos , E. Siamarkou, G. Tsipolitis , A. Zacharopoulou

**University of Ioánnina, Ioánnina, Greece**

I. Bestintzanos, I. Evangelou , C. Foudas, C. Kamtsikis, P. Katsoulis, P. Kokkas , P.G. Kosmoglou Kioseoglou , N. Manthos , I. Papadopoulos , J. Strologas 

**HUN-REN Wigner Research Centre for Physics, Budapest, Hungary**

C. Hajdu , D. Horvath<sup>31,32</sup> , K. Márton, A.J. Rádl<sup>33</sup> , F. Sikler , V. Veszpremi 

**MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary**

M. Csanád , K. Farkas , A. Fehérkuti<sup>34</sup> , M.M.A. Gadallah<sup>35</sup> , Á. Kadlecik , P. Major , G. Pásztor , G.I. Veres 

**Faculty of Informatics, University of Debrecen, Debrecen, Hungary**

B. Ujvari , G. Zilizi 

**HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary**

G. Bencze, S. Czellar, J. Molnar, Z. Szillasi

**Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary**

T. Csorgo<sup>34</sup> , F. Nemes<sup>34</sup> , T. Novak 

**Panjab University, Chandigarh, India**

S. Bansal , S.B. Beri, V. Bhatnagar , G. Chaudhary , S. Chauhan , N. Dhingra<sup>36</sup> , A. Kaur , A. Kaur , H. Kaur , M. Kaur , S. Kumar , T. Sheokand, J.B. Singh , A. Singla 

**University of Delhi, Delhi, India**

A. Bhardwaj , A. Chhetri , B.C. Choudhary , A. Kumar , A. Kumar , M. Naimuddin , K. Ranjan , M.K. Saini, S. Saumya 

**Saha Institute of Nuclear Physics, HBNI, Kolkata, India**

S. Baradia , S. Barman<sup>37</sup> , S. Bhattacharya , S. Das Gupta, S. Dutta , S. Dutta, S. Sarkar

**Indian Institute of Technology Madras, Madras, India**

M.M. Ameen , P.K. Behera , S.C. Behera , S. Chatterjee , G. Dash , P. Jana 

P. Kalbhor , S. Kamble , J.R. Komaragiri<sup>38</sup> , D. Kumar<sup>38</sup> , T. Mishra , B. Parida<sup>39</sup> , P.R. Pujahari , N.R. Saha , A. Sharma , A.K. Sikdar , R.K. Singh , P. Verma , S. Verma , A. Vijay 

#### **Tata Institute of Fundamental Research-A, Mumbai, India**

S. Dugad, G.B. Mohanty , M. Shelake, P. Suryadevara

#### **Tata Institute of Fundamental Research-B, Mumbai, India**

A. Bala , S. Banerjee , S. Bhowmik , R.M. Chatterjee, M. Guchait , Sh. Jain , A. Jaiswal, B.M. Joshi , S. Kumar , G. Majumder , K. Mazumdar , S. Parolia , A. Thachayath 

#### **National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India**

S. Bahinipati<sup>40</sup> , C. Kar , D. Maity<sup>41</sup> , P. Mal , K. Naskar<sup>41</sup> , A. Nayak<sup>41</sup> , S. Nayak, K. Pal , P. Sadangi, S.K. Swain , S. Varghese<sup>41</sup> , D. Vats<sup>41</sup> 

#### **Indian Institute of Science Education and Research (IISER), Pune, India**

S. Acharya<sup>42</sup> , A. Alpana , S. Dube , B. Gomber<sup>42</sup> , P. Hazarika , B. Kansal , A. Laha , B. Sahu<sup>42</sup> , S. Sharma , K.Y. Vaish 

#### **Isfahan University of Technology, Isfahan, Iran**

H. Bakhshiansohi<sup>43</sup> , A. Jafari<sup>44</sup> , M. Zeinali<sup>45</sup> 

#### **Institute for Research in Fundamental Sciences (IPM), Tehran, Iran**

S. Bashiri, S. Chenarani<sup>46</sup> , S.M. Etesami , Y. Hosseini , M. Khakzad , E. Khazaie , M. Mohammadi Najafabadi , S. Tizchang<sup>47</sup> 

#### **University College Dublin, Dublin, Ireland**

M. Felcini , M. Grunewald 

#### **INFN Sezione di Bari<sup>a</sup>, Università di Bari<sup>b</sup>, Politecnico di Bari<sup>c</sup>, Bari, Italy**

M. Abbrescia<sup>a,b</sup> , A. Colaleo<sup>a,b</sup> , D. Creanza<sup>a,c</sup> , B. D'Anzi<sup>a,b</sup> , N. De Filippis<sup>a,c</sup> , M. De Palma<sup>a,b</sup> , W. Elmetenawee<sup>a,b,16</sup> , N. Ferrara<sup>a,b</sup> , L. Fiore<sup>a</sup> , G. Iaselli<sup>a,c</sup> , L. Longo<sup>a</sup> , M. Louka<sup>a,b</sup> , G. Maggi<sup>a,c</sup> , M. Maggi<sup>a</sup> , I. Margjeka<sup>a</sup> , V. Mastrapasqua<sup>a,b</sup> , S. My<sup>a,b</sup> , S. Nuzzo<sup>a,b</sup> , A. Pellecchia<sup>a,b</sup> , A. Pompili<sup>a,b</sup> , G. Pugliese<sup>a,c</sup> , R. Radogna<sup>a,b</sup> , D. Ramos<sup>a</sup> , A. Ranieri<sup>a</sup> , L. Silvestris<sup>a</sup> , F.M. Simone<sup>a,c</sup> , Ü. Sözbilir<sup>a</sup> , A. Stamerra<sup>a,b</sup> , D. Troiano<sup>a,b</sup> , R. Venditti<sup>a,b</sup> , P. Verwilligen<sup>a</sup> , A. Zaza<sup>a,b</sup> 

#### **INFN Sezione di Bologna<sup>a</sup>, Università di Bologna<sup>b</sup>, Bologna, Italy**

G. Abbiendi<sup>a</sup> , C. Battilana<sup>a,b</sup> , D. Bonacorsi<sup>a,b</sup> , P. Capiluppi<sup>a,b</sup> , A. Castro<sup>†a,b</sup> , F.R. Cavallo<sup>a</sup> , M. Cuffiani<sup>a,b</sup> , G.M. Dallavalle<sup>a</sup> , T. Diotallevi<sup>a,b</sup> , F. Fabbri<sup>a</sup> , A. Fanfani<sup>a,b</sup> , D. Fasanella<sup>a</sup> , P. Giacomelli<sup>a</sup> , L. Giommi<sup>a,b</sup> , C. Grandi<sup>a</sup> , L. Guiducci<sup>a,b</sup> , S. Lo Meo<sup>a,48</sup> , M. Lorusso<sup>a,b</sup> , L. Lunerti<sup>a</sup> , S. Marcellini<sup>a</sup> , G. Masetti<sup>a</sup> , F.L. Navarria<sup>a,b</sup> , G. Paggi<sup>a,b</sup> , A. Perrotta<sup>a</sup> , F. Primavera<sup>a,b</sup> , A.M. Rossi<sup>a,b</sup> , S. Rossi Tisbeni<sup>a,b</sup> , T. Rovelli<sup>a,b</sup> , G.P. Siroli<sup>a,b</sup> 

#### **INFN Sezione di Catania<sup>a</sup>, Università di Catania<sup>b</sup>, Catania, Italy**

S. Costa<sup>a,b,49</sup> , A. Di Mattia<sup>a</sup> , A. Lapertosa<sup>a</sup> , R. Potenza<sup>a,b</sup> , A. Tricomi<sup>a,b,49</sup> 

#### **INFN Sezione di Firenze<sup>a</sup>, Università di Firenze<sup>b</sup>, Firenze, Italy**

P. Assiouras<sup>a</sup> , G. Barbagli<sup>a</sup> , G. Bardelli<sup>a,b</sup> , B. Camaiani<sup>a,b</sup> , A. Cassese<sup>a</sup> , R. Ceccarelli<sup>a</sup> , V. Ciulli<sup>a,b</sup> , C. Civinini<sup>a</sup> , R. D'Alessandro<sup>a,b</sup> , E. Focardi<sup>a,b</sup> , T. Kello<sup>a</sup> , G. Latino<sup>a,b</sup> , P. Lenzi<sup>a,b</sup> , M. Lizzo<sup>a</sup> , M. Meschini<sup>a</sup> , S. Paoletti<sup>a</sup> , A. Papanastassiou<sup>a,b</sup> , G. Sguazzoni<sup>a</sup> , L. Viliani<sup>a</sup> 

**INFN Laboratori Nazionali di Frascati, Frascati, Italy**

L. Benussi , S. Bianco , S. Meola<sup>50</sup> , D. Piccolo 

**INFN Sezione di Genova<sup>a</sup>, Università di Genova<sup>b</sup>, Genova, Italy**

M. Alves Gallo Pereira<sup>a</sup> , F. Ferro<sup>a</sup> , E. Robutti<sup>a</sup> , S. Tosi<sup>a,b</sup> 

**INFN Sezione di Milano-Bicocca<sup>a</sup>, Università di Milano-Bicocca<sup>b</sup>, Milano, Italy**

A. Benaglia<sup>a</sup> , F. Brivio<sup>a</sup> , F. Cetorelli<sup>a,b</sup> , F. De Guio<sup>a,b</sup> , M.E. Dinardo<sup>a,b</sup> , P. Dini<sup>a</sup> , S. Gennai<sup>a</sup> , R. Gerosa<sup>a,b</sup> , A. Ghezzi<sup>a,b</sup> , P. Govoni<sup>a,b</sup> , L. Guzzi<sup>a</sup> , G. Lavizzari<sup>a,b</sup> , M.T. Lucchini<sup>a,b</sup> , M. Malberti<sup>a</sup> , S. Malvezzi<sup>a</sup> , A. Massironi<sup>a</sup> , D. Menasce<sup>a</sup> , L. Moroni<sup>a</sup> , M. Paganoni<sup>a,b</sup> , S. Palluotto<sup>a,b</sup> , D. Pedrini<sup>a</sup> , A. Perego<sup>a,b</sup> , B.S. Pinolini<sup>a</sup> , G. Pizzati<sup>a,b</sup> , S. Ragazzi<sup>a,b</sup> , T. Tabarelli de Fatis<sup>a,b</sup> 

**INFN Sezione di Napoli<sup>a</sup>, Università di Napoli 'Federico II'<sup>b</sup>, Napoli, Italy; Università della Basilicata<sup>c</sup>, Potenza, Italy; Scuola Superiore Meridionale (SSM)<sup>d</sup>, Napoli, Italy**

S. Buontempo<sup>a</sup> , A. Cagnotta<sup>a,b</sup> , F. Carnevali<sup>a,b</sup> , N. Cavallo<sup>a,c</sup> , F. Fabozzi<sup>a,c</sup> , A.O.M. Iorio<sup>a,b</sup> , L. Lista<sup>a,b,51</sup> , P. Paolucci<sup>a,30</sup> , B. Rossi<sup>a</sup> 

**INFN Sezione di Padova<sup>a</sup>, Università di Padova<sup>b</sup>, Padova, Italy; Università di Trento<sup>c</sup>, Trento, Italy**

R. Ardino<sup>a</sup> , P. Azzi<sup>a</sup> , N. Bacchetta<sup>a,52</sup> , D. Bisello<sup>a,b</sup> , P. Bortignon<sup>a</sup> , G. Bortolato<sup>a,b</sup> , A. Bragagnolo<sup>a,b</sup> , A.C.M. Bulla<sup>a</sup> , R. Carlin<sup>a,b</sup> , P. Checchia<sup>a</sup> , T. Dorigo<sup>a,53</sup> , F. Gasparini<sup>a,b</sup> , U. Gasparini<sup>a,b</sup> , S. Giorgetti<sup>a</sup> , E. Lusiani<sup>a</sup> , M. Margoni<sup>a,b</sup> , A.T. Meneguzzo<sup>a,b</sup> , M. Migliorini<sup>a,b</sup> , J. Pazzini<sup>a,b</sup> , P. Ronchese<sup>a,b</sup> , R. Rossin<sup>a,b</sup> , F. Simonetto<sup>a,b</sup> , M. Tosi<sup>a,b</sup> , A. Triossi<sup>a,b</sup> , S. Ventura<sup>a</sup> , M. Zanetti<sup>a,b</sup> , P. Zotto<sup>a,b</sup> , A. Zucchetta<sup>a,b</sup> , G. Zumerle<sup>a,b</sup> 

**INFN Sezione di Pavia<sup>a</sup>, Università di Pavia<sup>b</sup>, Pavia, Italy**

A. Braghieri<sup>a</sup> , S. Calzaferri<sup>a</sup> , D. Fiorina<sup>a</sup> , P. Montagna<sup>a,b</sup> , V. Re<sup>a</sup> , C. Riccardi<sup>a,b</sup> , P. Salvini<sup>a</sup> , I. Vai<sup>a,b</sup> , P. Vitulo<sup>a,b</sup> 

**INFN Sezione di Perugia<sup>a</sup>, Università di Perugia<sup>b</sup>, Perugia, Italy**

S. Ajmal<sup>a,b</sup> , M.E. Ascioti<sup>a,b</sup> , G.M. Bilei<sup>a</sup> , C. Carrivale<sup>a,b</sup> , D. Ciangottini<sup>a,b</sup> , L. Fanò<sup>a,b</sup> , V. Mariani<sup>a,b</sup> , M. Menichelli<sup>a</sup> , F. Moscatelli<sup>a,54</sup> , A. Rossi<sup>a,b</sup> , A. Santocchia<sup>a,b</sup> , D. Spiga<sup>a</sup> , T. Tedeschi<sup>a,b</sup> 

**INFN Sezione di Pisa<sup>a</sup>, Università di Pisa<sup>b</sup>, Scuola Normale Superiore di Pisa<sup>c</sup>, Pisa, Italy; Università di Siena<sup>d</sup>, Siena, Italy**

C. Aimè<sup>a</sup> , C.A. Alexe<sup>a,c</sup> , P. Asenov<sup>a,b</sup> , P. Azzurri<sup>a</sup> , G. Bagliesi<sup>a</sup> , R. Bhattacharya<sup>a</sup> , L. Bianchini<sup>a,b</sup> , T. Boccali<sup>a</sup> , E. Bossini<sup>a</sup> , D. Bruschini<sup>a,c</sup> , R. Castaldi<sup>a</sup> , M.A. Ciocci<sup>a,b</sup> , M. Cipriani<sup>a,b</sup> , V. D'Amante<sup>a,d</sup> , R. Dell'Orso<sup>a</sup> , S. Donato<sup>a</sup> , A. Giassi<sup>a</sup> , F. Ligabue<sup>a,c</sup> , A.C. Marini<sup>a</sup> , D. Matos Figueiredo<sup>a</sup> , A. Messineo<sup>a,b</sup> , S. Mishra<sup>a</sup> , V.K. Muraleedharan Nair Bindhu<sup>a,b,41</sup> , M. Musich<sup>a,b</sup> , S. Nandan<sup>a</sup> , F. Palla<sup>a</sup> , A. Rizzi<sup>a,b</sup> , G. Rolandi<sup>a,c</sup> , S. Roy Chowdhury<sup>a</sup> , T. Sarkar<sup>a</sup> , A. Scribano<sup>a</sup> , P. Spagnolo<sup>a</sup> , R. Tenchini<sup>a</sup> , G. Tonelli<sup>a,b</sup> , N. Turini<sup>a,d</sup> , F. Vaselli<sup>a,c</sup> , A. Venturi<sup>a</sup> , P.G. Verdini<sup>a</sup> 

**INFN Sezione di Roma<sup>a</sup>, Sapienza Università di Roma<sup>b</sup>, Roma, Italy**

P. Barria<sup>a</sup> , C. Basile<sup>a,b</sup> , F. Cavallari<sup>a</sup> , L. Cunqueiro Mendez<sup>a,b</sup> , D. Del Re<sup>a,b</sup> , E. Di Marco<sup>a,b</sup> , M. Diemoz<sup>a</sup> , F. Errico<sup>a,b</sup> , R. Gargiulo<sup>a,b</sup> , E. Longo<sup>a,b</sup> , L. Martikainen<sup>a,b</sup> , J. Mijuskovic<sup>a,b</sup> , G. Organtini<sup>a,b</sup> , F. Pandolfi<sup>a</sup> , R. Paramatti<sup>a,b</sup> , C. Quaranta<sup>a,b</sup> , S. Rahatlou<sup>a,b</sup> , C. Rovelli<sup>a</sup> , F. Santanastasio<sup>a,b</sup> , L. Soffi<sup>a</sup> , V. Vladimirov<sup>a,b</sup> 

**INFN Sezione di Torino<sup>a</sup>, Università di Torino<sup>b</sup>, Torino, Italy; Università del Piemonte Orientale<sup>c</sup>, Novara, Italy**

N. Amapane<sup>a,b</sup> , R. Arcidiacono<sup>a,c</sup> , S. Argiro<sup>a,b</sup> , M. Arneodo<sup>a,c</sup> , N. Bartosik<sup>a</sup> , R. Bellan<sup>a,b</sup> , C. Biino<sup>a</sup> , C. Borca<sup>a,b</sup> , N. Cartiglia<sup>a</sup> , M. Costa<sup>a,b</sup> , R. Covarelli<sup>a,b</sup> , N. Demaria<sup>a</sup> , L. Finco<sup>a</sup> , M. Grippo<sup>a,b</sup> , B. Kiani<sup>a,b</sup> , F. Legger<sup>a</sup> , F. Luongo<sup>a,b</sup> , C. Mariotti<sup>a</sup> , L. Markovic<sup>a,b</sup> , S. Maselli<sup>a</sup> , A. Mecca<sup>a,b</sup> , L. Menzio<sup>a,b</sup> , P. Meridiani<sup>a</sup> , E. Migliore<sup>a,b</sup> , M. Monteno<sup>a</sup> , R. Mulargia<sup>a</sup> , M.M. Obertino<sup>a,b</sup> , G. Ortona<sup>a</sup> , L. Pacher<sup>a,b</sup> , N. Pastrone<sup>a</sup> , M. Pelliccioni<sup>a</sup> , M. Ruspà<sup>a,c</sup> , F. Siviero<sup>a,b</sup> , V. Sola<sup>a,b</sup> , A. Solano<sup>a,b</sup> , A. Staiano<sup>a</sup> , C. Tarricone<sup>a,b</sup> , D. Trocino<sup>a</sup> , G. Umoret<sup>a,b</sup> , R. White<sup>a,b</sup> 

**INFN Sezione di Trieste<sup>a</sup>, Università di Trieste<sup>b</sup>, Trieste, Italy**

J. Babbar<sup>a,b</sup> , S. Belforte<sup>a</sup> , V. Candelise<sup>a,b</sup> , M. Casarsa<sup>a</sup> , F. Cossutti<sup>a</sup> , K. De Leo<sup>a</sup> , G. Della Ricca<sup>a,b</sup> 

**Kyungpook National University, Daegu, Korea**

S. Dogra , J. Hong , J. Kim, D. Lee, H. Lee, S.W. Lee , C.S. Moon , Y.D. Oh , M.S. Ryu , S. Sekmen , B. Tae, Y.C. Yang 

**Department of Mathematics and Physics - GWNu, Gangneung, Korea**

M.S. Kim 

**Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea**

G. Bak , P. Gwak , H. Kim , D.H. Moon 

**Hanyang University, Seoul, Korea**

E. Asilar , J. Choi<sup>55</sup> , D. Kim , T.J. Kim , J.A. Merlin, Y. Ryou

**Korea University, Seoul, Korea**

S. Choi , S. Han, B. Hong , K. Lee, K.S. Lee , S. Lee , J. Yoo 

**Kyung Hee University, Department of Physics, Seoul, Korea**

J. Goh , S. Yang 

**Sejong University, Seoul, Korea**

Y. Kang , H. S. Kim , Y. Kim, S. Lee

**Seoul National University, Seoul, Korea**

J. Almond, J.H. Bhyun, J. Choi , J. Choi, W. Jun , J. Kim , Y.W. Kim , S. Ko , H. Lee , J. Lee , J. Lee , B.H. Oh , S.B. Oh , H. Seo , U.K. Yang, I. Yoon 

**University of Seoul, Seoul, Korea**

W. Jang , D.Y. Kang, S. Kim , B. Ko, J.S.H. Lee , Y. Lee , I.C. Park , Y. Roh, I.J. Watson 

**Yonsei University, Department of Physics, Seoul, Korea**

S. Ha , K. Hwang , B. Kim , H.D. Yoo 

**Sungkyunkwan University, Suwon, Korea**

M. Choi , M.R. Kim , H. Lee, Y. Lee , I. Yu 

**College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait**

T. Beyrouthy , Y. Gharbia 

**Kuwait University - College of Science - Department of Physics, Safat, Kuwait**

F. Alazemi 

**Riga Technical University, Riga, Latvia**

K. Dreimanis , A. Gaile , C. Munoz Diaz , D. Osite , G. Pikurs, A. Potrebko ,  
M. Seidel , D. Sidiropoulos Kontos 

**University of Latvia (LU), Riga, Latvia**

N.R. Strautnieks 

**Vilnius University, Vilnius, Lithuania**

M. Ambrozias , A. Juodagalvis , A. Rinkevicius , G. Tamulaitis 

**National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia**

I. Yusuff<sup>56</sup> , Z. Zolkapli

**Universidad de Sonora (UNISON), Hermosillo, Mexico**

J.F. Benitez , A. Castaneda Hernandez , H.A. Encinas Acosta, L.G. Gallegos Maríñez,  
M. León Coello , J.A. Murillo Quijada , A. Sehrawat , L. Valencia Palomo 

**Centro de Investigacion y de Estudios Avanzados del IPN, Mexico City, Mexico**

G. Ayala , H. Castilla-Valdez , H. Crotte Ledesma, E. De La Cruz-Burelo , I. Heredia-  
De La Cruz<sup>57</sup> , R. Lopez-Fernandez , J. Mejia Guisao , C.A. Mondragon Herrera,  
A. Sánchez Hernández 

**Universidad Iberoamericana, Mexico City, Mexico**

C. Oropeza Barrera , D.L. Ramirez Guadarrama, M. Ramírez García 

**Benemerita Universidad Autonoma de Puebla, Puebla, Mexico**

I. Bautista , F.E. Neri Huerta , I. Pedraza , H.A. Salazar Ibarguen , C. Uribe Estrada 

**University of Montenegro, Podgorica, Montenegro**

I. Bubanja , N. Raicevic 

**University of Canterbury, Christchurch, New Zealand**

P.H. Butler 

**National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan**

A. Ahmad , M.I. Asghar, A. Awais , M.I.M. Awan, H.R. Hoorani , W.A. Khan 

**AGH University of Krakow, Krakow, Poland**

V. Avati, A. Bellora , L. Forthomme , L. Grzanka , M. Malawski , K. Piotrkowski

**National Centre for Nuclear Research, Swierk, Poland**

H. Bialkowska , M. Bluj , M. Górski , M. Kazana , M. Szleper , P. Zalewski 

**Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland**

K. Bunkowski , K. Doroba , A. Kalinowski , M. Konecki , J. Krolikowski ,  
A. Muhammad 

**Warsaw University of Technology, Warsaw, Poland**

P. Fokow , K. Pozniak , W. Zabolotny 

**Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal**

M. Araujo , D. Bastos , C. Beirão Da Cruz E Silva , A. Boletti , M. Bozzo ,  
T. Camporesi , G. Da Molin , P. Faccioli , M. Gallinaro , J. Hollar , N. Leonardo ,  
G.B. Marozzo , A. Petrilli , M. Pisano , J. Seixas , J. Varela , J.W. Wulff 

**Faculty of Physics, University of Belgrade, Belgrade, Serbia**

P. Adzic , P. Milenovic 

**VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia**

D. Devetak, M. Dordevic , J. Milosevic , L. Nadderd , V. Rekovic, M. Stojanovic 

**Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain**

J. Alcaraz Maestre , Cristina F. Bedoya , J.A. Brochero Cifuentes , Oliver M. Carretero , M. Cepeda , M. Cerrada , N. Colino , B. De La Cruz , A. Delgado Peris , A. Escalante Del Valle , D. Fernández Del Val , J.P. Fernández Ramos , J. Flix , M.C. Fouz , O. Gonzalez Lopez , S. Goy Lopez , J.M. Hernandez , M.I. Josa , J. Llorente Merino , C. Martin Perez , E. Martin Viscasillas , D. Moran , C. M. Morcillo Perez , Á. Navarro Tobar , C. Perez Dengra , A. Pérez-Calero Yzquierdo , J. Puerta Pelayo , I. Redondo , J. Sastre , J. Vazquez Escobar 

**Universidad Autónoma de Madrid, Madrid, Spain**

J.F. de Trocóniz 

**Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain**

B. Alvarez Gonzalez , J. Cuevas , J. Fernandez Menendez , S. Folgueras , I. Gonzalez Caballero , P. Leguina , E. Palencia Cortezon , J. Prado Pico , V. Rodríguez Bouza , A. Soto Rodríguez , A. Trapote , C. Vico Villalba , P. Vischia 

**Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain**

S. Blanco Fernández , I.J. Cabrillo , A. Calderon , J. Duarte Campderros , M. Fernandez , G. Gomez , C. Lasasa García , R. Lopez Ruiz , C. Martinez Rivero , P. Martinez Ruiz del Arbol , F. Matorras , P. Matorras Cuevas , E. Navarrete Ramos , J. Piedra Gomez , L. Scodellaro , I. Vila , J.M. Vizan Garcia 

**University of Colombo, Colombo, Sri Lanka**

B. Kailasapathy<sup>58</sup> , D.D.C. Wickramarathna 

**University of Ruhuna, Department of Physics, Matara, Sri Lanka**

W.G.D. Dharmaratna<sup>59</sup> , K. Liyanage , N. Perera 

**CERN, European Organization for Nuclear Research, Geneva, Switzerland**

D. Abbaneo , C. Amendola , E. Auffray , G. Auzinger , J. Baechler, D. Barney , A. Bermúdez Martínez , M. Bianco , A.A. Bin Anuar , A. Bocci , L. Borgonovi , C. Botta , E. Brondolin , C.E. Brown , C. Caillol , G. Cerminara , N. Chernyavskaya , D. d'Enterria , A. Dabrowski , A. David , A. De Roeck , M.M. Defranchis , M. Deile , M. Dobson , G. Franzoni , W. Funk , S. Giani, D. Gigi, K. Gill , F. Glege , M. Glowacki, J. Hegeman , J.K. Heikkilä , B. Huber , V. Innocente , T. James , P. Janot , O. Kaluzinska , O. Karacheban<sup>28</sup> , G. Karathanasis , S. Laurila , P. Lecoq , E. Leutgeb , C. Lourenço , M. Magherini , L. Malgeri , M. Mannelli , M. Matthewman, A. Mehta , F. Meijers , S. Mersi , E. Meschi , V. Milosevic , F. Monti , F. Moortgat , M. Mulders , I. Neutelings , S. Orfanelli, F. Pantaleo , G. Petrucciani , A. Pfeiffer , M. Pierini , H. Qu , D. Rabady , B. Ribeiro Lopes , F. Riti , M. Rovere , H. Sakulin , R. Salvatico , S. Sanchez Cruz , S. Scarfi , C. Schwick, M. Selvaggi , A. Sharma , K. Shchelina , P. Silva , P. Sphicas<sup>60</sup> , A.G. Stahl Leitner , A. Steen , S. Summers , D. Treille , P. Tropea , D. Walter , J. Wanczyk<sup>61</sup> , J. Wang, S. Wuchterl , P. Zehetner , P. Zejd , W.D. Zeuner

**PSI Center for Neutron and Muon Sciences, Villigen, Switzerland**

T. Bevilacqua<sup>62</sup> , L. Caminada<sup>62</sup> , A. Ebrahimi , W. Erdmann , R. Horisberger 

Q. Ingram , H.C. Kaestli , D. Kotlinski , C. Lange , M. Missiroli<sup>62</sup> , L. Noehte<sup>62</sup> , T. Rohe , A. Samalan

**ETH Zurich - Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland**

T.K. Aarrestad , M. Backhaus , G. Bonomelli , A. Calandri , C. Cazzaniga , K. Datta , P. De Bryas Dexmiers D'archiac<sup>61</sup> , A. De Cosa , G. Dissertori , M. Dittmar, M. Donegà , F. Eble , M. Galli , K. Gedia , F. Glessgen , C. Grab , N. Härringer , T.G. Harte, D. Hits , W. Lustermann , A.-M. Lyon , R.A. Manzoni , M. Marchegiani , L. Marchese , A. Mascellani<sup>61</sup> , F. Nessi-Tedaldi , F. Pauss , V. Perovic , S. Pigazzini , B. Ristic , R. Seidita , J. Steggemann<sup>61</sup> , A. Tarabini , D. Valsecchi , R. Wallny 

**Universität Zürich, Zurich, Switzerland**

C. Amsler<sup>63</sup> , P. Bäertschi , M.F. Canelli , K. Cormier , M. Huwiler , W. Jin , A. Jofrehei , B. Kilminster , S. Leontsinis , S.P. Liehti , A. Macchiolo , P. Meiring , F. Meng , J. Motta , A. Reimers , P. Robmann, M. Senger , E. Shokr, F. Stäger , R. Tramontano 

**National Central University, Chung-Li, Taiwan**

C. Adloff<sup>64</sup>, D. Bhowmik, C.M. Kuo, W. Lin, P.K. Rout , P.C. Tiwari<sup>38</sup> 

**National Taiwan University (NTU), Taipei, Taiwan**

L. Ceard, K.F. Chen , Z.g. Chen, A. De Iorio , W.-S. Hou , T.h. Hsu, Y.w. Kao, S. Karmakar , G. Kole , Y.y. Li , R.-S. Lu , E. Paganis , X.f. Su , J. Thomas-Wilsker , L.s. Tsai, D. Tzionou, H.y. Wu, E. Yazgan 

**High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand**

C. Asawatangkuldee , N. Srimanobhas , V. Wachirapusanand 

**Tunis El Manar University, Tunis, Tunisia**

Y. Maghrbi 

**Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey**

D. Agyel , F. Boran , F. Dolek , I. Dumanoglu<sup>65</sup> , E. Eskut , Y. Guler<sup>66</sup> , E. Gurpinar Guler<sup>66</sup> , C. Isik , O. Kara, A. Kayis Topaksu , Y. Komurcu , G. Onengut , K. Ozdemir<sup>67</sup> , A. Polatoz , B. Tali<sup>68</sup> , U.G. Tok , E. Uslan , I.S. Zorbakir 

**Middle East Technical University, Physics Department, Ankara, Turkey**

M. Yalvac<sup>69</sup> 

**Bogazici University, Istanbul, Turkey**

B. Akgun , I.O. Atakisi , E. Gülmez , M. Kaya<sup>70</sup> , O. Kaya<sup>71</sup> , S. Tekten<sup>72</sup> 

**Istanbul Technical University, Istanbul, Turkey**

A. Cakir , K. Cankocak<sup>65,73</sup> , S. Sen<sup>74</sup> 

**Istanbul University, Istanbul, Turkey**

O. Aydilek<sup>75</sup> , B. Haciasahinoglu , I. Hos<sup>76</sup> , B. Kaynak , S. Ozkorucuklu , O. Potok , H. Sert , C. Simsek , C. Zorbilmez 

**Yildiz Technical University, Istanbul, Turkey**

S. Cerci , B. Isildak<sup>77</sup> , D. Sunar Cerci , T. Yetkin 

**Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine**

A. Boyaryntsev , B. Grynyov 

**National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine**L. Levchuk **University of Bristol, Bristol, United Kingdom**D. Anthony , J.J. Brooke , A. Bundock , F. Bury , E. Clement , D. Cussans , H. Flacher , J. Goldstein , H.F. Heath , M.-L. Holmberg , L. Kreczko , S. Paramesvaran , L. Robertshaw, V.J. Smith , K. Walkingshaw Pass**Rutherford Appleton Laboratory, Didcot, United Kingdom**A.H. Ball, K.W. Bell , A. Belyaev<sup>78</sup> , C. Brew , R.M. Brown , D.J.A. Cockerill , C. Cooke , A. Elliot , K.V. Ellis, K. Harder , S. Harper , J. Linacre , K. Manolopoulos, D.M. Newbold , E. Olaiya, D. Petyt , T. Reis , A.R. Sahasransu , G. Salvi , T. Schuh, C.H. Shepherd-Themistocleous , I.R. Tomalin , K.C. Whalen , T. Williams **Imperial College, London, United Kingdom**I. Andreou , R. Bainbridge , P. Bloch , O. Buchmuller, C.A. Carrillo Montoya , G.S. Chahal<sup>79</sup> , D. Colling , J.S. Dancu, I. Das , P. Dauncey , G. Davies , M. Della Negra , S. Fayer, G. Fedi , G. Hall , A. Howard, G. Iles , C.R. Knight , P. Krueper, J. Langford , K.H. Law , J. León Holgado , L. Lyons , A.-M. Magnan , B. Maier , S. Mallios, M. Mieskolainen , J. Nash<sup>80</sup> , M. Pesaresi , P.B. Pradeep, B.C. Radburn-Smith , A. Richards, A. Rose , K. Savva , C. Seez , R. Shukla , A. Tapper , K. Uchida , G.P. Uttley , T. Virdee<sup>30</sup> , M. Vojinovic , N. Wardle , D. Winterbottom **Brunel University, Uxbridge, United Kingdom**J.E. Cole , A. Khan, P. Kyberd , I.D. Reid **Baylor University, Waco, Texas, USA**S. Abdullin , A. Brinkerhoff , E. Collins , M.R. Darwish , J. Dittmann , K. Hatakeyama , V. Hegde , J. Hiltbrand , B. McMaster , J. Samudio , S. Sawant , C. Sutantawibul , J. Wilson **Catholic University of America, Washington, DC, USA**R. Bartek , A. Dominguez , A.E. Simsek , S.S. Yu **The University of Alabama, Tuscaloosa, Alabama, USA**B. Bam , A. Buchot Perraguin , R. Chudasama , S.I. Cooper , C. Crovella , S.V. Gleyzer , E. Pearson, C.U. Perez , P. Rumerio<sup>81</sup> , E. Usai , R. Yi **Boston University, Boston, Massachusetts, USA**A. Akpinar , C. Cosby , G. De Castro, Z. Demiragli , C. Erice , C. Fangmeier , C. Fernandez Madrazo , E. Fontanesi , D. Gastler , F. Golf , S. Jeon , J. O'cain, I. Reed , J. Rohlf , K. Salyer , D. Sperka , D. Spitzbart , I. Suarez , A. Tsatsos , A.G. Zecchinelli **Brown University, Providence, Rhode Island, USA**G. Barone , G. Benelli , D. Cutts , L. Gouskos , M. Hadley , U. Heintz , K.W. Ho , J.M. Hogan<sup>82</sup> , T. Kwon , G. Landsberg , K.T. Lau , J. Luo , S. Mondal , T. Russell, S. Sagir<sup>83</sup> , X. Shen , M. Stamenkovic , N. Venkatasubramanian**University of California, Davis, Davis, California, USA**S. Abbott , B. Barton , C. Brainerd , R. Breedon , H. Cai , M. Calderon De La Barca Sanchez , M. Chertok , M. Citron , J. Conway , P.T. Cox , R. Erbacher , F. Jensen , O. Kukral , G. Mocellin , M. Mulhearn , S. Ostrom 

W. Wei , S. Yoo , F. Zhang 

#### University of California, Los Angeles, California, USA

K. Adamidis , M. Bachtis , D. Campos , R. Cousins , A. Datta , G. Flores Avila , J. Hauser , M. Ignatenko , M.A. Iqbal , T. Lam , Y.f. Lo , E. Manca , A. Nunez Del Prado , D. Saltzberg , V. Valuev 

#### University of California, Riverside, Riverside, California, USA

R. Clare , J.W. Gary , G. Hanson 

#### University of California, San Diego, La Jolla, California, USA

A. Aportela , A. Arora , J.G. Branson , S. Cittolin , S. Cooperstein , D. Diaz , J. Duarte , L. Giannini , Y. Gu , J. Guiang , R. Kansal , V. Krutelyov , R. Lee , J. Letts , M. Masciovecchio , F. Mokhtar , S. Mukherjee , M. Pieri , D. Primosch , M. Quinnan , V. Sharma , M. Tadel , E. Vourliotis , F. Würthwein , Y. Xiang , A. Yagil 

#### University of California, Santa Barbara - Department of Physics, Santa Barbara, California, USA

A. Barzdukas , L. Brennan , C. Campagnari , K. Downham , C. Grieco , M.M. Hussain , J. Incandela , J. Kim , A.J. Li , P. Masterson , H. Mei , J. Richman , S.N. Santpur , U. Sarica , R. Schmitz , F. Setti , J. Sheplock , D. Stuart , T.Á. Vámi , X. Yan , D. Zhang 

#### California Institute of Technology, Pasadena, California, USA

S. Bhattacharya , A. Bornheim , O. Cerri , J. Mao , H.B. Newman , G. Reales Gutiérrez , M. Spiropulu , J.R. Vlimant , C. Wang , S. Xie , R.Y. Zhu 

#### Carnegie Mellon University, Pittsburgh, Pennsylvania, USA

J. Alison , S. An , P. Bryant , M. Cremonesi , V. Dutta , T. Ferguson , T.A. Gómez Espinosa , A. Harilal , A. Kallil Tharayil , C. Liu , T. Mudholkar , S. Murthy , P. Palit , K. Park , M. Paulini , A. Roberts , A. Sanchez , W. Terrill 

#### University of Colorado Boulder, Boulder, Colorado, USA

J.P. Cumalat , W.T. Ford , A. Hart , A. Hassani , N. Manganelli , J. Pearkes , C. Savard , N. Schonbeck , K. Stenson , K.A. Ulmer , S.R. Wagner , N. Zipper , D. Zuolo 

#### Cornell University, Ithaca, New York, USA

J. Alexander , X. Chen , D.J. Cranshaw , J. Dickinson , J. Fan , X. Fan , S. Hogan , P. Kotamnives , J. Monroy , M. Oshiro , J.R. Patterson , M. Reid , A. Ryd , J. Thom , P. Wittich , R. Zou 

#### Fermi National Accelerator Laboratory, Batavia, Illinois, USA

M. Albrow , M. Alyari , O. Amram , G. Apollinari , A. Apresyan , L.A.T. Bauerdick , D. Berry , J. Berryhill , P.C. Bhat , K. Burkett , J.N. Butler , A. Canepa , G.B. Cerati , H.W.K. Cheung , F. Chlebana , G. Cummings , I. Dutta , V.D. Elvira , J. Freeman , A. Gandrakota , Z. Gecse , L. Gray , D. Green , A. Grummer , S. Grünendahl , D. Guerrero , O. Gutsche , R.M. Harris , T.C. Herwig , J. Hirschauer , B. Jayatilaka , S. Jindariani , M. Johnson , U. Joshi , T. Klijsma , B. Klima , K.H.M. Kwok , S. Lammel , C. Lee , D. Lincoln , R. Lipton , T. Liu , K. Maeshima , D. Mason , P. McBride , P. Merkel , S. Mrenna , S. Nahn , J. Ngadiuba , D. Noonan , S. Norberg , V. Papadimitriou , N. Pastika , K. Pedro , C. Pena<sup>84</sup> , F. Ravera , A. Reinsvold Hall<sup>85</sup> , L. Ristori , M. Safdari , E. Sexton-Kennedy , N. Smith 

A. Soha , L. Spiegel , S. Stoynev , J. Strait , L. Taylor , S. Tkaczyk , N.V. Tran , L. Uplegger , E.W. Vaandering , I. Zoi 

#### **University of Florida, Gainesville, Florida, USA**

C. Aruta , P. Avery , D. Bourilkov , P. Chang , V. Cherepanov , R.D. Field, C. Huh , E. Koenig , M. Kolosova , J. Konigsberg , A. Korytov , K. Matchev , N. Menendez , G. Mitselmakher , K. Mohrman , A. Muthirakalayil Madhu , N. Rawal , S. Rosenzweig , Y. Takahashi , J. Wang 

#### **Florida State University, Tallahassee, Florida, USA**

T. Adams , A. Al Kadhimi , A. Askew , S. Bower , R. Hashmi , R.S. Kim , S. Kim , T. Kolberg , G. Martinez, H. Prosper , P.R. Prova, M. Wulansatiti , R. Yohay , J. Zhang

#### **Florida Institute of Technology, Melbourne, Florida, USA**

B. Alsufyani , S. Butalla , S. Das , T. Elkafrawy<sup>86</sup> , M. Hohlmann , E. Yanes

#### **University of Illinois Chicago, Chicago, Illinois, USA**

M.R. Adams , A. Baty , C. Bennett, R. Cavanaugh , R. Escobar Franco , O. Evdokimov , C.E. Gerber , M. Hawksworth, A. Hingrajiya, D.J. Hofman , J.h. Lee , D. S. Lemos , C. Mills , S. Nanda , G. Oh , B. Ozek , D. Pilipovic , R. Pradhan , E. Prifti, T. Roy , S. Rudrabhatla , N. Singh, M.B. Tonjes , N. Varelas , M.A. Wadud , Z. Ye , J. Yoo 

#### **The University of Iowa, Iowa City, Iowa, USA**

M. Alhusseini , D. Blend, K. Dilsiz<sup>87</sup> , L. Emediato , G. Karaman , O.K. Köseyan , J.-P. Merlo, A. Mestvirishvili<sup>88</sup> , O. Neogi, H. Ogul<sup>89</sup> , Y. Onel , A. Penzo , C. Snyder, E. Tiras<sup>90</sup> 

#### **Johns Hopkins University, Baltimore, Maryland, USA**

B. Blumenfeld , L. Corcodilos , J. Davis , A.V. Gritsan , L. Kang , S. Kyriacou , P. Maksimovic , M. Roguljic , J. Roskes , S. Sekhar , M. Swartz 

#### **The University of Kansas, Lawrence, Kansas, USA**

A. Abreu , L.F. Alcerro Alcerro , J. Anguiano , S. Arteaga Escatel , P. Baringer , A. Bean , Z. Flowers , D. Grove , J. King , G. Krintiras , M. Lazarovits , C. Le Mahieu , J. Marquez , M. Murray , M. Nickel , M. Pitt , S. Popescu<sup>91</sup> , C. Rogan , C. Royon , S. Sanders , C. Smith , G. Wilson 

#### **Kansas State University, Manhattan, Kansas, USA**

B. Allmond , R. Gujju Gurunadha , A. Ivanov , K. Kaadze , Y. Maravin , J. Natoli , D. Roy , G. Sorrentino 

#### **University of Maryland, College Park, Maryland, USA**

A. Baden , A. Belloni , J. Bistany-riebman, Y.M. Chen , S.C. Eno , N.J. Hadley , S. Jabeen , R.G. Kellogg , T. Koeth , B. Kronheim, Y. Lai , S. Lascio , A.C. Mignerey , S. Nabili , C. Palmer , C. Papageorgakis , M.M. Paranjpe, E. Popova<sup>92</sup> , A. Shevelev , L. Wang , L. Zhang 

#### **Massachusetts Institute of Technology, Cambridge, Massachusetts, USA**

C. Baldenegro Barrera , J. Bendavid , S. Bright-Thonney , I.A. Cali , P.c. Chou , M. D'Alfonso , J. Eysermans , C. Freer , G. Gomez-Ceballos , M. Goncharov, G. Grosso, P. Harris, D. Hoang, D. Kovalskyi , J. Krupa , L. Lavezzo , Y.-J. Lee , K. Long , C. McGinn , A. Novak , M.I. Park , C. Paus , C. Reissel , C. Roland , G. Roland , S. Rothman , G.S.F. Stephans , Z. Wang , B. Wyslouch , T. J. Yang 

**University of Minnesota, Minneapolis, Minnesota, USA**

B. Crossman , C. Kapsiak , M. Krohn , D. Mahon , J. Mans , B. Marzocchi ,  
M. Revering , R. Rusack , R. Saradhy , N. Strobbe 

**University of Nebraska-Lincoln, Lincoln, Nebraska, USA**

K. Bloom , D.R. Claes , G. Haza , J. Hossain , C. Joo , I. Kravchenko , A. Rohilla ,  
J.E. Siado , W. Tabb , A. Vagnerini , A. Wightman , F. Yan , D. Yu 

**State University of New York at Buffalo, Buffalo, New York, USA**

H. Bandyopadhyay , L. Hay , H.w. Hsia , I. Iashvili , A. Kalogeropoulos ,  
A. Kharchilava , M. Morris , D. Nguyen , S. Rappoccio , H. Rejeb Sfar, A. Williams ,  
P. Young 

**Northeastern University, Boston, Massachusetts, USA**

G. Alverson , E. Barberis , J. Bonilla , B. Bylsma, M. Campana , J. Dervan ,  
Y. Haddad , Y. Han , I. Israr , A. Krishna , P. Levchenko , J. Li , M. Lu ,  
R. Mccarthy , D.M. Morse , V. Nguyen , T. Orimoto , A. Parker , L. Skinnari ,  
E. Tsai , D. Wood 

**Northwestern University, Evanston, Illinois, USA**

S. Dittmer , K.A. Hahn , D. Li , Y. Liu , M. McGinnis , Y. Miao , D.G. Monk ,  
M.H. Schmitt , A. Taliercio , M. Velasco

**University of Notre Dame, Notre Dame, Indiana, USA**

G. Agarwal , R. Band , R. Bucci, S. Castells , A. Das , R. Goldouzian , M. Hildreth ,  
K. Hurtado Anampa , T. Ivanov , C. Jessop , K. Lannon , J. Lawrence , N. Loukas ,  
L. Lutton , J. Mariano, N. Marinelli, I. Mcalister, T. McCauley , C. Mcgrady , C. Moore ,  
Y. Musienko<sup>23</sup> , H. Nelson , M. Osherson , A. Piccinelli , R. Ruchti , A. Townsend ,  
Y. Wan, M. Wayne , H. Yockey, M. Zarucki , L. Zygalá 

**The Ohio State University, Columbus, Ohio, USA**

A. Basnet , M. Carrigan , L.S. Durkin , C. Hill , M. Joyce , M. Nunez Ornelas , K. Wei,  
D.A. Wenzl, B.L. Winer , B. R. Yates 

**Princeton University, Princeton, New Jersey, USA**

H. Bouchamaoui , K. Coldham, P. Das , G. Dezoort , P. Elmer , A. Frankenthal ,  
B. Greenberg , N. Haubrich , K. Kennedy, G. Kopp , S. Kwan , D. Lange ,  
A. Loeliger , D. Marlow , I. Ojalvo , J. Olsen , F. Simpson , D. Stickland , C. Tully ,  
L.H. Vage

**University of Puerto Rico, Mayaguez, Puerto Rico, USA**

S. Malik , R. Sharma

**Purdue University, West Lafayette, Indiana, USA**

A.S. Bakshi , S. Chandra , R. Chawla , A. Gu , L. Gutay, M. Jones , A.W. Jung ,  
A.M. Koshy, M. Liu , G. Negro , N. Neumeister , G. Paspalaki , S. Piperov ,  
V. Scheurer, J.F. Schulte , A. K. Viridi , F. Wang , A. Wildridge , W. Xie , Y. Yao 

**Purdue University Northwest, Hammond, Indiana, USA**

J. Dolen , N. Parashar , A. Pathak 

**Rice University, Houston, Texas, USA**

D. Acosta , A. Agrawal , T. Carnahan , K.M. Ecklund , P.J. Fernández Manteca ,  
S. Freed, P. Gardner, F.J.M. Geurts , I. Krommydas , W. Li , J. Lin , O. Miguel Colin ,  
B.P. Padley , R. Redjimi, J. Rotter , E. Yigitbasi , Y. Zhang 

**University of Rochester, Rochester, New York, USA**

A. Bodek , P. de Barbaro , R. Demina , J.L. Dulemba , A. Garcia-Bellido , O. Hindrichs , A. Khukhunaishvili , N. Parmar , P. Parygin<sup>92</sup> , R. Taus 

**Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA**

B. Chiarito, J.P. Chou , S.V. Clark , D. Gadkari , Y. Gershtein , E. Halkiadakis , M. Heindl , C. Houghton , D. Jaroslawski , S. Konstantinou , I. Laflotte , A. Lath , R. Montalvo, K. Nash, J. Reichert , P. Saha , S. Salur , S. Schnetzer, S. Somalwar , R. Stone , S.A. Thayil , S. Thomas, J. Vora 

**University of Tennessee, Knoxville, Tennessee, USA**

D. Ally , A.G. Delannoy , S. Fiorendi , S. Higginbotham , T. Holmes , A.R. Kanunganti , N. Karunarathna , L. Lee , E. Nibigira , S. Spanier 

**Texas A&M University, College Station, Texas, USA**

D. Aebi , M. Ahmad , T. Akhter , K. Androsov<sup>61</sup> , O. Bouhali<sup>93</sup> , R. Eusebi , J. Gilmore , T. Huang , T. Kamon<sup>94</sup> , H. Kim , S. Luo , R. Mueller , D. Overton , A. Safonov 

**Texas Tech University, Lubbock, Texas, USA**

N. Akchurin , J. Damgov , Y. Feng , N. Gogate , Y. Kazhykarim, K. Lamichhane , S.W. Lee , C. Madrid , A. Mankel , T. Peltola , I. Volobouev 

**Vanderbilt University, Nashville, Tennessee, USA**

E. Appelt , Y. Chen , S. Greene, A. Gurrola , W. Johns , R. Kunnawalkam Elayavalli , A. Melo , D. Rathjens , F. Romeo , P. Sheldon , S. Tuo , J. Velkovska , J. Viinikainen 

**University of Virginia, Charlottesville, Virginia, USA**

B. Cardwell , H. Chung, B. Cox , J. Hakala , R. Hirosky , A. Ledovskoy , C. Mantilla , C. Neu , C. Ramón Álvarez 

**Wayne State University, Detroit, Michigan, USA**

S. Bhattacharya , P.E. Karchin 

**University of Wisconsin - Madison, Madison, Wisconsin, USA**

A. Aravind , S. Banerjee , K. Black , T. Bose , E. Chavez , S. Dasu , P. Everaerts , C. Galloni, H. He , M. Herndon , A. Herve , C.K. Koraka , A. Lanaro, R. Loveless , J. Madhusudanan Sreekala , A. Mallampalli , A. Mohammadi , S. Mondal, G. Parida , L. Pétré , D. Pinna, A. Savin, V. Shang , V. Sharma , W.H. Smith , D. Teague, H.F. Tsoi , W. Vetens , A. Warden 

**Authors affiliated with an international laboratory covered by a cooperation agreement with CERN**

S. Afanasiev , V. Alexakhin , D. Budkouski , I. Golutvin<sup>†</sup> , I. Gorbunov , V. Karjavine , O. Kodolova<sup>95,92</sup> , V. Korenkov , A. Lanev , A. Malakhov , V. Matveev<sup>96</sup> , A. Nikitenko<sup>97,95</sup> , V. Palichik , V. Perelygin , M. Savina , V. Shalaev , S. Shmatov , S. Shulha , V. Smirnov , O. Teryaev , N. Voytishin , B.S. Yuldashev<sup>†98</sup>, A. Zarubin , I. Zhizhin , Yu. Andreev , A. Dermenev , S. Gninenko , N. Golubev , A. Karneyeu , D. Kirpichnikov , M. Kirsanov , N. Krasnikov , I. Tlisova , A. Toropin 

**Authors affiliated with an institute formerly covered by a cooperation agreement with CERN**

G. Gavrilo , V. Golovtcov , Y. Ivanov , V. Kim<sup>99</sup> , V. Murzin , V. Oreshkin , D. Sosnov , V. Sulimov , L. Uvarov , A. Vorobyev<sup>†</sup>, T. Aushev , K. Ivanov , V. Gavrilov , N. Lychkovskaya , V. Popov , A. Zhokin , R. Chistov<sup>99</sup> , M. Danilov<sup>99</sup> 

S. Polikarpov<sup>99</sup> , V. Andreev , M. Azarkin , M. Kirakosyan, A. Terkulov , E. Boos , V. Bunichev , M. Dubinin<sup>84</sup> , L. Dudko , A. Ershov , V. Klyukhin , M. Perfilov , V. Savrin , P. Volkov , G. Vorotnikov , V. Blinov<sup>99</sup>, T. Dimova<sup>99</sup> , A. Kozyrev<sup>99</sup> , O. Radchenko<sup>99</sup> , Y. Skovpen<sup>99</sup> , V. Kachanov , S. Slabospitskii , A. Uzunian , A. Babaev , V. Borshch , D. Druzhkin 

†: Deceased

<sup>1</sup>Also at Yerevan State University, Yerevan, Armenia

<sup>2</sup>Also at TU Wien, Vienna, Austria

<sup>3</sup>Also at Ghent University, Ghent, Belgium

<sup>4</sup>Also at Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil

<sup>5</sup>Also at FACAMP - Faculdades de Campinas, Sao Paulo, Brazil

<sup>6</sup>Also at Universidade Estadual de Campinas, Campinas, Brazil

<sup>7</sup>Also at Federal University of Rio Grande do Sul, Porto Alegre, Brazil

<sup>8</sup>Also at University of Chinese Academy of Sciences, Beijing, China

<sup>9</sup>Also at China Center of Advanced Science and Technology, Beijing, China

<sup>10</sup>Also at University of Chinese Academy of Sciences, Beijing, China

<sup>11</sup>Also at China Spallation Neutron Source, Guangdong, China

<sup>12</sup>Now at Henan Normal University, Xinxiang, China

<sup>13</sup>Also at University of Shanghai for Science and Technology, Shanghai, China

<sup>14</sup>Now at The University of Iowa, Iowa City, Iowa, USA

<sup>15</sup>Also at an institute formerly covered by a cooperation agreement with CERN

<sup>16</sup>Also at Helwan University, Cairo, Egypt

<sup>17</sup>Now at Zewail City of Science and Technology, Zewail, Egypt

<sup>18</sup>Now at British University in Egypt, Cairo, Egypt

<sup>19</sup>Now at Cairo University, Cairo, Egypt

<sup>20</sup>Also at Purdue University, West Lafayette, Indiana, USA

<sup>21</sup>Also at Université de Haute Alsace, Mulhouse, France

<sup>22</sup>Also at Istinye University, Istanbul, Turkey

<sup>23</sup>Also at an international laboratory covered by a cooperation agreement with CERN

<sup>24</sup>Also at The University of the State of Amazonas, Manaus, Brazil

<sup>25</sup>Also at University of Hamburg, Hamburg, Germany

<sup>26</sup>Also at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany

<sup>27</sup>Also at Bergische University Wuppertal (BUW), Wuppertal, Germany

<sup>28</sup>Also at Brandenburg University of Technology, Cottbus, Germany

<sup>29</sup>Also at Forschungszentrum Jülich, Jülich, Germany

<sup>30</sup>Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland

<sup>31</sup>Also at HUN-REN ATOMKI - Institute of Nuclear Research, Debrecen, Hungary

<sup>32</sup>Now at Universitatea Babes-Bolyai - Facultatea de Fizica, Cluj-Napoca, Romania

<sup>33</sup>Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary

<sup>34</sup>Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary

<sup>35</sup>Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt

<sup>36</sup>Also at Punjab Agricultural University, Ludhiana, India

<sup>37</sup>Also at University of Visva-Bharati, Santiniketan, India

<sup>38</sup>Also at Indian Institute of Science (IISc), Bangalore, India

<sup>39</sup>Also at Amity University Uttar Pradesh, Noida, India

<sup>40</sup>Also at IIT Bhubaneswar, Bhubaneswar, India

<sup>41</sup>Also at Institute of Physics, Bhubaneswar, India

<sup>42</sup>Also at University of Hyderabad, Hyderabad, India

- <sup>43</sup>Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany
- <sup>44</sup>Also at Isfahan University of Technology, Isfahan, Iran
- <sup>45</sup>Also at Sharif University of Technology, Tehran, Iran
- <sup>46</sup>Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran
- <sup>47</sup>Also at Department of Physics, Faculty of Science, Arak University, ARAK, Iran
- <sup>48</sup>Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy
- <sup>49</sup>Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy
- <sup>50</sup>Also at Università degli Studi Guglielmo Marconi, Roma, Italy
- <sup>51</sup>Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy
- <sup>52</sup>Also at Fermi National Accelerator Laboratory, Batavia, Illinois, USA
- <sup>53</sup>Also at Lulea University of Technology, Lulea, Sweden
- <sup>54</sup>Also at Consiglio Nazionale delle Ricerche - Istituto Officina dei Materiali, Perugia, Italy
- <sup>55</sup>Also at Institut de Physique des 2 Infinis de Lyon (IP2I ), Villeurbanne, France
- <sup>56</sup>Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia
- <sup>57</sup>Also at Consejo Nacional de Ciencia y Tecnología, Mexico City, Mexico
- <sup>58</sup>Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka
- <sup>59</sup>Also at Saegis Campus, Nugegoda, Sri Lanka
- <sup>60</sup>Also at National and Kapodistrian University of Athens, Athens, Greece
- <sup>61</sup>Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland
- <sup>62</sup>Also at Universität Zürich, Zurich, Switzerland
- <sup>63</sup>Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria
- <sup>64</sup>Also at Laboratoire d'Annecy-le-Vieux de Physique des Particules, IN2P3-CNRS, Annecy-le-Vieux, France
- <sup>65</sup>Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey
- <sup>66</sup>Also at Konya Technical University, Konya, Turkey
- <sup>67</sup>Also at Izmir Bakircay University, Izmir, Turkey
- <sup>68</sup>Also at Adiyaman University, Adiyaman, Turkey
- <sup>69</sup>Also at Bozok Universitetesi Rektörlüğü, Yozgat, Turkey
- <sup>70</sup>Also at Marmara University, Istanbul, Turkey
- <sup>71</sup>Also at Milli Savunma University, Istanbul, Turkey
- <sup>72</sup>Also at Kafkas University, Kars, Turkey
- <sup>73</sup>Now at Istanbul Okan University, Istanbul, Turkey
- <sup>74</sup>Also at Hacettepe University, Ankara, Turkey
- <sup>75</sup>Also at Erzincan Binali Yildirim University, Erzincan, Turkey
- <sup>76</sup>Also at Istanbul University - Cerrahpasa, Faculty of Engineering, Istanbul, Turkey
- <sup>77</sup>Also at Yildiz Technical University, Istanbul, Turkey
- <sup>78</sup>Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom
- <sup>79</sup>Also at IPPP Durham University, Durham, United Kingdom
- <sup>80</sup>Also at Monash University, Faculty of Science, Clayton, Australia
- <sup>81</sup>Also at Università di Torino, Torino, Italy
- <sup>82</sup>Also at Bethel University, St. Paul, Minnesota, USA
- <sup>83</sup>Also at Karamanoğlu Mehmetbey University, Karaman, Turkey
- <sup>84</sup>Also at California Institute of Technology, Pasadena, California, USA
- <sup>85</sup>Also at United States Naval Academy, Annapolis, Maryland, USA

---

<sup>86</sup>Also at Ain Shams University, Cairo, Egypt

<sup>87</sup>Also at Bingol University, Bingol, Turkey

<sup>88</sup>Also at Georgian Technical University, Tbilisi, Georgia

<sup>89</sup>Also at Sinop University, Sinop, Turkey

<sup>90</sup>Also at Erciyes University, Kayseri, Turkey

<sup>91</sup>Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania

<sup>92</sup>Now at another institute formerly covered by a cooperation agreement with CERN

<sup>93</sup>Also at Texas A&M University at Qatar, Doha, Qatar

<sup>94</sup>Also at Kyungpook National University, Daegu, Korea

<sup>95</sup>Also at Yerevan Physics Institute, Yerevan, Armenia

<sup>96</sup>Also at another international laboratory covered by a cooperation agreement with CERN

<sup>97</sup>Also at Imperial College, London, United Kingdom

<sup>98</sup>Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan

<sup>99</sup>Also at another institute formerly covered by a cooperation agreement with CERN