

Search for charged lepton flavor violating Z and Z' boson decays in proton-proton collisions at $\sqrt{s} = 13$ TeV

A. Hayrapetyan *et al.*^{*}
(CMS Collaboration)



(Received 9 August 2025; accepted 16 October 2025; published 15 December 2025)

A search for flavor violating decays of the Z boson to charged leptons is performed using data from proton-proton collisions at $\sqrt{s} = 13$ TeV collected with the CMS detector at the LHC, corresponding to an integrated luminosity of 138 fb^{-1} . Each of the decays $Z \rightarrow e\mu$, $Z \rightarrow e\tau$, and $Z \rightarrow \mu\tau$ is considered. The data are consistent with the backgrounds expected from standard model processes. For the $Z \rightarrow e\mu$ channel the observed (expected) 95% confidence level upper limit on the branching fraction is $1.9(2.0) \times 10^{-7}$, which is the most stringent direct limit to date on this process; the corresponding limits for the $Z \rightarrow e\tau$ and $Z \rightarrow \mu\tau$ channels are $13.8(11.4) \times 10^{-6}$ and $12.0(5.3) \times 10^{-6}$, respectively. Additionally, the $e\mu$ final state is used to search for lepton flavor violating decays of Z' resonances in the mass range from 110 to 500 GeV. No significant excess is observed above the predicted background levels.

DOI: [10.1103/8lw4-6lyk](https://doi.org/10.1103/8lw4-6lyk)

I. INTRODUCTION

The standard model (SM) has been highly successful in describing nature on the small distance scale. An important exception is the neutrino sector, where in the SM neutrinos are massless, while the observation of neutrino flavor oscillations [1–4] indicates that neutrinos have small but nonzero masses, and that their individual flavor is not a conserved quantity. Generation number violation has also been observed in the quark sector, but not to date among charged leptons. In an extended version of the SM that includes neutrino masses, the anticipated rates of charged lepton flavor violation (CLFV) induced by neutrino mixing are vanishingly small; e.g., the branching fraction $\mathcal{B}(Z \rightarrow e\mu) \approx \mathcal{O}(10^{-60})$ [5]. Therefore any observation of a CLFV process would be direct evidence of physics beyond the SM (BSM).

Potentially measurable rates of CLFV in various processes, including Z boson decay, are predicted in BSM scenarios such as the seesaw mechanism, and supersymmetry and leptoquark models [6,7]. In addition, new particles predicted by these and other BSM models may exhibit CLFV through their decays. The new particles include additional gauge bosons such as a Z' boson, which would generally be expected to be heavier than the Z boson [8].

Axionlike particles have also been suggested as potential BSM particles that may decay into flavor-violating states [9,10].

Searches for CLFV have been conducted in low-energy processes including $\mu \rightarrow e\gamma$ [11], $\mu \rightarrow eee$ [12], $\mu^- N(A, Z) \rightarrow e^- N(A, Z)$ [13], $e^+ e^- \rightarrow \mu^+ (e^+) \tau^-$ [14,15], $\tau \rightarrow \ell \gamma$ ($\ell = e, \mu$) [16], and $\tau \rightarrow \ell' \ell^+ \ell^-$ [17]. Limits from these reactions lead to indirect constraints on the Z boson branching fractions $\mathcal{B}(Z \rightarrow e\mu) < 10^{-12}$, $\mathcal{B}(Z \rightarrow e\tau) < 10^{-7}$, and $\mathcal{B}(Z \rightarrow \mu\tau) < 2 \times 10^{-7}$ [6]. However, direct measurements of CLFV Z decays are needed to rule out anomalous couplings or cancellations that could evade these limits [18]. Searches for such decays have been performed by experiments at the CERN LEP [19,20] and LHC [21–23] colliders. Current limits on the branching fractions for $Z \rightarrow e\mu$, $Z \rightarrow e\tau$, and $Z \rightarrow \mu\tau$ are 2.6×10^{-7} [23], 5.0×10^{-6} [22], and 6.5×10^{-6} [22], respectively.

This paper presents a search for decays of the Z boson in the CLFV channels $Z \rightarrow e\mu$, $Z \rightarrow e\tau$, and $Z \rightarrow \mu\tau$, together with a scan of the $e\mu$ invariant mass over the range 110–500 GeV to search for potential heavy neutral resonances. The proton-proton (pp) collision data at $\sqrt{s} = 13$ TeV were collected with the CMS detector in 2016–2018, and correspond to an integrated luminosity of 138 fb^{-1} . New multivariate techniques extend the reach beyond previous results.

The signal extraction makes use of two main variables: the invariant mass of the (visible) daughters of the $Z^{(\prime)}$ boson candidate, and a multivariate discriminator, implemented as a boosted decision tree (BDT), that combines kinematic observables chosen to discriminate signal from background. For the $e\mu$ channel, where the $Z^{(\prime)}$ boson is

^{*}Full author list given at the end of the article.

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

fully reconstructed, we extract the signal yield with a fit to the $e\mu$ mass distributions, relying mainly on the sidebands to determine the background under the signal peak. These distributions are taken from separate subsamples that vary in purity, as measured by the BDT discriminator. To distinguish signal from background in the $e\tau$ and $\mu\tau$ channels, where the peak in the invariant mass distribution of the visible Z boson daughters is less sharply defined, we instead rely primarily on distributions in the BDT discriminant. These distributions are taken from subsamples defined by ranges in the visible invariant mass chosen according to the dominance of the signal or of various backgrounds.

In Sec. II we describe the CMS detector and particle reconstruction, and in Sec. III the simulation of signal and background processes. The event selection is given in Sec. IV, followed by the signal extraction for the $Z \rightarrow e\mu$ and $Z \rightarrow \ell\tau$ channels in Secs. V and VI, respectively. Systematic uncertainties are discussed in Sec. VII, and the results in Sec. VIII. We conclude with a summary in Sec. IX. Tabulated results are provided in the HEPData record for this analysis [24].

II. THE CMS DETECTOR AND EVENT RECONSTRUCTION

The central feature of the CMS apparatus is a superconducting solenoid of 6 m internal diameter, providing a magnetic field of 3.8 T. Within the solenoid volume are a silicon pixel and strip tracker, a lead tungstate crystal electromagnetic calorimeter (ECAL), and a brass and scintillator hadron calorimeter (HCAL), each composed of a barrel and two end cap sections. Forward calorimeters extend the coverage in pseudorapidity (η) provided by the barrel and end cap detectors. The ECAL consists of 75 848 lead tungstate crystals, which provide pseudorapidity coverage of $|\eta| < 1.48$ in its barrel region and $1.48 < |\eta| < 3.00$ in the two end cap regions. Preshower detectors consisting of two planes of silicon sensors interleaved with a total of three radiation lengths of lead are located in front of each end cap detector. Muons are measured in gas-ionization detectors embedded in the steel flux-return yoke outside the solenoid. A more detailed description of the CMS detector, together with a definition of the coordinate system used and the relevant kinematic variables, is reported in Refs. [25,26].

The silicon tracker used in 2016 measured charged particles within the range $|\eta| < 2.5$. For nonisolated particles of transverse momentum p_T between 1 and 10 GeV and $|\eta| < 1.4$, the track resolutions were typically 1.5% in p_T and 25–90 (45–150) μm in the transverse (longitudinal) impact parameter [27]. At the start of 2017, a new pixel detector was installed [28], which provided expanded acceptance, to $|\eta| < 3.0$, and improved transverse impact parameter resolution of 20–75 μm [29]. The primary vertex is taken to be the vertex corresponding to the hardest

scattering in the event, evaluated using tracking information alone, as described in Sec. 9.4.1 of Ref. [30].

The global event reconstruction (also called particle-flow event reconstruction [31]) aims to reconstruct and identify each individual particle in an event, with an optimized combination of all subdetector information. In this process, the identification (ID) of the particle type (photon, electron, muon, charged hadron, neutral hadron) plays an important role in the determination of the particle direction and energy. Photons are identified as ECAL energy clusters not linked to the extrapolation of any charged-particle trajectory to the ECAL. Electrons are identified as a primary charged-particle track and potentially many ECAL energy clusters corresponding to this track extrapolation to the ECAL and to possible bremsstrahlung photons emitted along the way through the tracker material. Muons are identified as tracks in the central tracker consistent with either a track or several hits in the muon system, and associated with calorimeter deposits compatible with the muon hypothesis. Charged hadrons are identified as charged-particle tracks neither identified as electrons, nor as muons. Finally, neutral hadrons are identified as HCAL energy clusters not linked to any charged hadron trajectory, or as a combined ECAL and HCAL energy excess with respect to the expected charged-hadron energy deposit.

For each event, hadronic jets are clustered from these reconstructed particles using the infrared- and collinear-safe anti- k_T algorithm [32,33] with a distance parameter of 0.4. Jet momentum is determined as the vectorial sum of all particle momenta in the jet, and is found from simulation to be, on average, within 5–10% of the true momentum over the entire spectrum and detector acceptance.

Additional pp interactions within the same or nearby bunch crossings (pileup) can contribute additional tracks and calorimetric energy depositions to the jet momentum. To mitigate this effect, charged particles identified to be originating from pileup vertices are discarded, and an offset correction is applied to correct for remaining contributions [34].

Jet energy corrections are derived from simulation to bring the measured response of jets equal, on average, to that of particle-level jets. *In situ* measurements of the momentum balance in dijet, γ + jets, Z + jets, and strong production of jets (QCD multijet) events are used to account for any residual differences in the jet energy scale between data obtained with the detector (“data”) and simulation [35]. The jet energy resolution amounts typically to 15–20% at 30 GeV, 10% at 100 GeV, and 5% at 1 TeV [35]. Additional selection criteria are applied to each jet to remove jets potentially dominated by anomalous contributions from various subdetector component or reconstruction failures.

Jets associated with b hadron decays are identified within the tracker acceptance of $|\eta| < 2.5$ using the DeepJet deep neural network ID algorithms [36–38]. We use both tight and

loose working points of this algorithm. Their efficiencies for b jets (light-flavor quark and gluon jets) have been measured in $t\bar{t}$ events to be 84 (10)% for the loose working point and 52 (0.1)% for the tight working point.

Hadronic tau lepton decays (τ_h) are reconstructed from jets using the hadrons-plus-strips algorithm [39], which combines 1 or 3 tracks with energy deposits in the calorimeters to identify the various hadronic τ decay modes. Neutral pions are reconstructed as clusters with dynamic size in (η, ϕ) , from reconstructed electrons and photons; the size of the dimension in the azimuthal angle ϕ (“strip”) varies as a function of the p_T of the electron or photon candidate, because of the dispersion in the magnetic field. To distinguish between genuine τ_h decays and jets originating from the hadronization of quarks or gluons, and from electrons or muons, we use the DeepTau algorithm [40]. Information from all individual reconstructed particles near the τ_h axis is combined with properties of the τ_h candidate and the event. The rate of a jet to be misidentified as a τ_h candidate depends on the p_T and quark flavor of the jet. In a data sample enriched in $W + \text{jets}$ events this misidentification (misID) rate is measured to be about 0.25%, with an ID efficiency for genuine τ_h of 60%, at the working point and kinematic regime of our signal regions (SRs) [40]. The rate for electrons (muons) to be misidentified as τ_h is 2.60 (0.03)% for a genuine τ_h ID efficiency of 80 (> 99)%.

The electron momentum is estimated by combining the energy measurement in the ECAL with the corresponding momentum measurement in the tracker [41]. The momentum resolution for electrons with $p_T \approx 45$ GeV from $Z \rightarrow ee$ decays ranges from 1.6 to 5%. The resolution is generally better in the barrel region than in the end caps, and it also depends on the bremsstrahlung energy emitted by the electron as it traverses the material upstream of the ECAL [41,42].

Muons are measured in the range $|\eta| < 2.4$, with detection planes made using three technologies: drift tubes, cathode strip chambers, and resistive-plate chambers. The efficiency for the reconstruction and ID of muons is greater than 96%. Once these muon candidates are matched to tracks measured in the silicon tracker, the relative p_T resolution for muons with p_T up to 100 GeV is 1% in the barrel and 3% in the end caps. The p_T resolution in the barrel is below 7% for muons with p_T up to 1 TeV [43].

The missing transverse momentum vector $\vec{p}_T^{\text{miss}}$ is computed as the negative vector sum of the transverse momenta of all the particle-flow candidates in an event; its magnitude is denoted as p_T^{miss} . The $\vec{p}_T^{\text{miss}}$ is modified to account for corrections to the energy scale of the reconstructed jets in the event. To reduce the dependence of $\vec{p}_T^{\text{miss}}$ on pileup, we apply the pileup-per-particle-identification algorithm [44], which weights the particle-flow candidates entering the $\vec{p}_T^{\text{miss}}$ sum by their probability of originating from the primary vertex [45].

Events of interest are selected with a two-tiered trigger system. The first level, composed of custom hardware processors, uses information from the calorimeters and muon detectors to select events at a rate of around 100 kHz within a fixed latency of about 4 μs [46]. The second level, known as the high-level trigger, comprises a farm of processors running a version of the full event reconstruction software optimized for fast processing, and reduces the event rate to around 1 kHz before data storage [47,48].

III. SIMULATED EVENT SAMPLES

Monte Carlo simulated samples are used to optimize the analysis and to model various background sources. Events arising from $W + \text{jets}$ are simulated, for the 2016 (2017–2018) data taking periods, with the MadGraph5_aMC@NLO 2.2.2 (2.4.2) [49,50] event generator at leading order (LO) accuracy. Drell-Yan (DY), $W\gamma$, and WW events are simulated with the same generators at next-to-LO (NLO) accuracy. The POWHEG v2.0 [51–60] program at NLO is used to generate events from $t\bar{t}$, tW , single top quark, and H boson production (where H denotes the 125 GeV Higgs boson), as well as diboson events originating from WW , WH , or ZH production. The WZ and ZZ events are generated at LO using the PYTHIA 8.205 [61] package. The simulated background samples are normalized using cross section calculations performed at NLO or next-to-NLO (NNLO) precision [62–66].

Signal processes are generated at LO using PYTHIA. For each CLFV decay channel ($Z \rightarrow e\mu$, $Z \rightarrow e\tau$, or $Z \rightarrow \mu\tau$), a tree-level coupling is added, with no coupling to the other decay channels. Events for the off- Z mass scan are generated with PYTHIA, including only the contribution of a vector boson Z' , and not considering any mixing with the Z boson or photon. The decay amplitude is calculated using tree-level couplings to the $e\mu$ final state with no other decay channels. The resonances are generated in the narrow-width approximation, i.e., with a width of 0.1 GeV, far below the experimental resolution. Events are simulated with masses of 100, 125, 150, 175, 200, 300, 400, and 500 GeV.

In all samples the hadronization, as well as the underlying event, are modeled with PYTHIA using the CUETP8M1 [67] (CP5 [68]) tune for the 2016 (2017–18) data-taking years. The LO (NLO) simulations use the MLM (FxFx) scheme for matching the products of the matrix-element calculations to the parton shower simulation [69]. The parton distribution functions used with the matrix-element generators are taken from the NNPDF3.0 [70] set at LO or NLO, except for the WW samples for the 2017–2018 data-taking years where NNPDF3.1 [71] is used; for PYTHIA the NNPDF2.3LO [72] set is used.

The CMS detector response is simulated using Geant4 [73]. Simulated samples are reconstructed with the same software packages as used for collision data. To account for effects of pileup, minimum-bias events simulated with PYTHIA are added to the hard-scattering process, with a

distribution in the number of vertices matching that in the data.

IV. EVENT SELECTION

An event is required to pass either a single-electron or single-muon trigger. To ensure that the trigger is fully efficient ($> 95\%$) for the events selected for analysis, an electron (muon) matched to the single-electron (muon) trigger is required to have p_T exceeding a threshold of 29 (25) GeV for data collected in 2016, 35 (28) GeV for data collected in 2017, and 34 (25) GeV for data collected in 2018.

For all three search channels we make a selection of events with two lepton candidates (e , μ , or τ_h) of different flavor and opposite charge. These events are sorted into three separate data channels; $e\mu$ [from which the $Z^{(\prime)} \rightarrow e\mu$, $Z \rightarrow e\tau_\mu$, and $Z \rightarrow \mu\tau_e$ candidates are selected], $e\tau_h$, and $\mu\tau_h$. To suppress SM backgrounds that produce pairs of leptons, events are rejected if they contain additional electrons or muons, and for the $e\tau_h$ and $\mu\tau_h$ channels, also if they contain additional τ_h candidates. In the search for the $Z \rightarrow e\tau$ or $Z \rightarrow \mu\tau$ signal we include both a hadronic τ decay category ($e\tau_h$ or $\mu\tau_h$, respectively) and a different-flavor leptonic τ decay category ($e\tau_\mu$ or $\mu\tau_e$). As each CLFV Z boson decay signal is the target of a separate search, there is no requirement that the $Z \rightarrow e\tau_\mu$, $Z \rightarrow \mu\tau_e$, and $Z \rightarrow e\mu$ samples be nonoverlapping.

To select a light lepton ℓ (e or μ) that is produced in the decay of a Z boson or tau lepton and not from a hadron within a quark jet, we construct the relative isolation variable I_{rel}^ℓ , defined as a function of the lepton's transverse momentum p_T^ℓ as

$$I_{\text{rel}}^\ell = \frac{\sum p_T^{\text{charged}} + \max[0, \sum p_T^{\text{neutral}} + \sum p_T^\gamma - p_T^{\text{PU}}]}{p_T^\ell}, \quad (1)$$

where the sums include the particles, excluding the lepton itself, within a cone of size $\Delta R \equiv \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2} = 0.4(0.3)$ about the electron (muon). Here p_T^{charged} refers to charged hadrons, p_T^{neutral} to neutral hadrons, p_T^γ to photons, and p_T^{PU} to the estimated neutral contribution from pileup. Electrons and muons are required to pass a tight isolation requirement, $I_{\text{rel}}^\ell < 0.15$.

To select prompt electrons and muons we require their transverse and longitudinal impact parameters, d_{xy} and d_z , to be less than 0.2 and 0.5 cm, respectively. For further suppression of the background from nonprompt electrons (primarily from QCD multijet and W + jets events), we require electrons to have $p_T > 15$ GeV. Electron candidates with ECAL clusters in the gap region between the barrel and end cap ($1.444 < |\eta| < 1.566$) are rejected.

The τ_h candidates are required to have $p_T > 20$ GeV and $|\eta| < 2.3$, with $|d_z| < 0.2$ cm. They are further required to

pass the following working points of the DeepTau tagging algorithm [40] for the suppression of: muons (“tight”); jets [“tight” for the SR, “very loose” for the control regions (CRs) used to estimate the misID background]; and electrons (“loose,” except “very tight” for the $e\tau_h$ channel SR to suppress the background from $Z \rightarrow ee$ where one of the electrons is misidentified as a τ_h candidate).

Jets are required to have $p_T > 20$ GeV and $|\eta| < 3$. Where jets are used in the event selection, jets within $\Delta R < 0.3$ of identified leptons are excluded from consideration.

Additional requirements specific to the individual channels are described in the corresponding sections below.

V. SEARCH FOR $Z \rightarrow e\mu$

The $Z \rightarrow e\mu$ search uses the $e\mu$ data channel, requiring exactly one electron and one muon candidate. Both leptons are required to have $|\eta| < 2.4$, and $p_T > 20$ GeV in addition to the trigger threshold requirement noted in Sec. IV above. The invariant mass $m_{e\mu}$ of the dilepton system is required to satisfy $70 < m_{e\mu} < 110$ GeV; this range includes sidebands around the Z boson mass (91.1880 ± 0.0020 GeV [74]) for estimation of the background. The sources of backgrounds near the Z boson mass are WW , $t\bar{t}$, DY production of $\tau\tau$, nonprompt $e\mu$, and $Z \rightarrow \mu\mu^* \rightarrow \mu\mu\gamma$ production.




FIG. 1. Invariant mass of the $e\mu$ system for data (points, with bars denoting statistical uncertainty) and simulated background (stacked filled histograms) for events passing the baseline selection. In the legend, “W, multi-V” refers to W + jets events having a jet that is misidentified as a lepton, together with multiple vector boson production. The hatched histogram shows a hypothetical $Z \rightarrow e\mu$ signal normalized to a branching fraction of 10^{-5} . The lower panel shows the ratio of the data to simulated background yields, with the statistical (combined systematic and statistical) uncertainty in the simulated yield indicated by the filled (hatched) gray band.

Events from $t\bar{t}$ (as well as subdominant single t quark) production are suppressed by rejecting events with jets that satisfy the loose b tagging [38]. To suppress the $Z \rightarrow \tau\tau$ contribution we require, in addition to the requirements on the track impact parameters given in Sec. IV, that their significances satisfy $|d_{xy}|/\sigma_{d_{xy}} < 3$ and $|d_z|/\sigma_{d_z} < 4.7$, with $\sigma_{d_{xy}}$ and σ_{d_z} the respective uncertainties. The efficiency reduction from these requirements on the track impact parameters is $\approx 5\%$ on the signal, and $\approx 20\%$ on those backgrounds that have tau leptons in the final state.

To extract the signal yield we exploit the narrow resonance peak of the signal against a mainly smoothly falling background in the $m_{e\mu}$ spectrum. The background shape arises from the sharply falling dominant contribution of $Z \rightarrow \tau_e \tau_\mu$, combined with a nearly flat spectrum from WW , $t\bar{t}$, and nonprompt $e\mu$ events. The distribution in $m_{e\mu}$ of the events surviving the baseline selection described above is shown in Fig. 1, for the data and for the simulated background components. The shape of the simulated signal, normalized to a branching fraction of 10^{-5} , is superimposed.

The final selection in the $Z \rightarrow e\mu$ search makes use of a BDT trained to distinguish signal from background events. The signal is then extracted from a fit of signal and background probability density functions (PDFs) to the $m_{e\mu}$ distribution, for three SRs defined as ranges of the BDT discriminant value.

A. The BDT implementation for $Z \rightarrow e\mu$

While DY production of τ pairs is one of the main backgrounds, these events are concentrated in the region below the signal resonance. In our SRs, both for the Z and Z' cases, events from $t\bar{t}$ and WW production are dominant at the signal resonance mass. Therefore, a single BDT is trained using simulated events for the $Z \rightarrow e\mu$ signal and for the background. The background is represented by the WW simulation, which is similar to $t\bar{t}$ after our baseline selection. We select disjoint training and testing samples, restricting $m_{e\mu}$ for the training sample to the Z -boson peak region, 86–96 GeV, to limit the influence of any correlations of the BDT features with $m_{e\mu}$. The BDT algorithm is implemented with the XGBoost [75] package. The hyperparameters of the BDT are optimized with a random grid search. The input features aim to exploit the difference between signal and background in p_T^{miss} and other kinematic variables. For the signal process, the Z boson is produced predominantly near threshold and decays to visible daughters, which are roughly collinear and have similar p_T , with little p_T^{miss} . Backgrounds with leptonically decaying W bosons give rise to a characteristic distribution in the transverse mass m_T formed from the W daughter lepton and p_T^{miss} . The transverse mass $m_T(1,2)$ is defined for a system of two particles (or of two systems of particles) with transverse momenta $p_T^{(1,2)}$ and azimuthal separation $\Delta\phi$ as

$$m_T(1,2) = \sqrt{2p_T^{(1)}p_T^{(2)}(1 - \cos \Delta\phi)}. \quad (2)$$

Since the signal extraction in the $Z \rightarrow e\mu$ search treats the BDT and $m_{e\mu}$ as independent variables, we include only features having small correlations with $m_{e\mu}$ in the BDT. From all the features tested, those that have the most discriminating power are selected:

- (1) $p_T^{\text{trailing}}/p_T^{\text{leading}}$;
- (2) $p_T^{e\mu}$;
- (3) $\eta_{e\mu}$;
- (4) $m_T(p_T^{\text{miss}}, \text{leading})$;
- (5) $m_T(p_T^{\text{miss}}, \text{trailing})$;
- (6) $|\Delta\phi(p_T^{\text{miss}}, e\mu)|$; and
- (7) p_T^{miss} .




FIG. 2. The upper plot shows unity-normalized distributions of the $Z \rightarrow e\mu$ BDT score for simulated events from the BDT test samples satisfying $70 < m_{e\mu} < 110$ GeV. The blue and red histograms represent the signal and WW background, respectively. The lower plot displays the unity-normalized distribution of $m_{e\mu}$ for events in the $t\bar{t}$ data CR used to check for BDT mass-spectrum bias, for several BDT thresholds; the vertical bars show the statistical uncertainties.

The terms “leading” and “trailing” denote the higher- and lower- p_T lepton, respectively.

The resulting distributions in the discriminant value (BDT score) for the BDT test events in the range 70–110 GeV in $m_{e\mu}$ are shown for signal and background in the upper plot of Fig. 2. We see from this plot that substantial discrimination between signal and background is achieved.

The $m_{e\mu}$ distribution for the WW background test sample confirms that no structure is introduced by the training. To check this conclusion in data, we construct a CR enriched in $t\bar{t}$ (and WW) events by applying the SR selection but replacing the veto on any loose b -tagged jet with a requirement of one tight b -tagged jet. The lower plot in Fig. 2 shows the $m_{e\mu}$ shape for events in this CR for each of the BDT thresholds used to define the BDT binning. The




FIG. 3. Fits of the data sidebands with the background functions for the BDT score ranges 0.3–0.7 (upper left), 0.7–0.9 (upper right), and 0.9–1.0 (lower). In the upper panel of each plot, the black (blue) points with bars show the data with statistical uncertainties in the sideband (signal) region, the solid red line shows the average background prediction, and the green curve shows the $Z \rightarrow \mu\mu^* \rightarrow \mu\mu\gamma$ background component. The lower panel in each plot shows the ratio of the difference between the data and the average background prediction to the uncertainty in the data, and the gray band shows the spread of the background estimates from the separate families of parametric functions.

distribution is seen to be free of structure within uncertainties. The BDT-score ranges chosen to define the SRs are 0.3–0.7, 0.7–0.9, and 0.9–1.0. This binning optimized the sensitivity of the simultaneous fit to the $m_{e\mu}$ distributions in all of the SRs.

B. Fit model for $Z \rightarrow e\mu$

For the events within one of the BDT score categories, parametric functions of $m_{e\mu}$ are defined to describe the signal and backgrounds in that category. The parameters of these functions are then determined from a fit to the binned data distribution, performed simultaneously in all categories.

The $Z \rightarrow e\mu$ signal distribution is described by a double-sided Crystal Ball function [76,77]. This function has a Gaussian core with independent power law tails defined such that the function is continuous and differentiable at all points. Its parameters are determined from a fit to simulated signal events, and fixed in the fit to the data.

The nonresonant background spectrum is modeled by analytic functions obtained directly from fits to the data, which removes from the model any dependence on simulation. The forms of these functions are established with fits to the data in the $m_{e\mu}$ sidebands, defined as $70 < m_{e\mu} < 86$ GeV and $96 < m_{e\mu} < 110$ GeV. Rather than choosing a single function that best fits the data sidebands for each region, we employ the “discrete profiling” method [78] in which an envelope of all well-fitting functions considered are included in the background model. The fit is then free to choose the background function that minimizes the likelihood at each point of the fit, profiling over the discrete nuisance parameter that represents the function choice.

Each of the individual functions included in the background model envelope is constructed from a broad, low-mass Gaussian component to represent the tail of the $Z \rightarrow \tau\tau$ spectrum, and an additional function, chosen through statistical tests based on CRs in data, for the nonresonant processes. This function has the form

$$f(x) = f_g g(x|\mu, \sigma) + (1 - f_g) h_n(x|\vec{\theta}), \quad (3)$$

where g is the Gaussian PDF with mean μ and standard deviation σ , f_g its fraction in the model, and h_n is the n th-order PDF of the function family h with parameters $\vec{\theta}$. The free parameters in the fit to the data are f_g , μ , σ , and $\vec{\theta}$. The h_n function is chosen from among the Chebychev polynomial, exponential sum, or power law sum families. The background functions that pass a χ^2 -probability test for fit consistency, $p(\chi^2) > 1\%$, are added to the model, and an F-test [79], $p(\chi_n^2 - \chi_{n+1}^2, \Delta n_d) < 5\%$, is performed to determine the maximum function order for each family. Here Δn_d is the number of degrees of freedom added by a step to the next higher-order function. Tests performed with large numbers of simulated background events confirm that the (zero) signal strength extracted from the fit is free of bias.

The model in Eq. (3) does not adequately describe the background from $Z \rightarrow \mu\mu^* \rightarrow \mu\mu\gamma$ events, which occur in cases where a muon loses a significant fraction of its energy to final-state radiation, thereby failing to reach the muon detectors, and the resulting photon is misidentified as an electron. This gives rise to a broad peak below the Z mass region that is not differentiated by the BDT since the background comes from true Z boson decays. Therefore, this background needs to be accounted for separately in the final fit. It is modeled with a double-sided Crystal Ball function, determined from fits to simulated $Z \rightarrow \mu\mu$ events that are reconstructed as $Z \rightarrow e\mu$. We find satisfactory agreement between the simulation and data in a sample of pairs reconstructed as $e\mu$, but having the same sign (SS) of charge. These events have a large fraction of electron candidates from misID background and exhibit a shape similar to that of the opposite-sign background.

The background model envelope from the fit to the $m_{e\mu}$ sidebands is shown in Fig. 3 for each BDT category. The data points in blue in the figure are not revealed or used for the determination of the background shape.

C. Application to the Z' mass scan

For the $e\mu$ final state, a scan for new neutral resonances Z' above the Z boson mass is also performed, using the same strategy and selection as in the $Z \rightarrow e\mu$ search. The extended-range $m_{e\mu}$ distribution in data and simulation is




FIG. 4. Invariant mass of the $e\mu$ system for data (points) and simulated background (stacked filled histograms), for events that pass the baseline selection except for that selection’s upper limit on the invariant mass. The lower panel shows the ratio of the data to simulated yields, with the statistical (combined systematic and statistical) uncertainty in the simulated yield indicated by the filled (hatched) gray band.

shown in Fig. 4. The most important difference in procedure for this scan is that a sliding window in $m_{e\mu}$ is used for successive scan points. With this strategy, potential masses of Z' are explored from 110 to 500 GeV, the upper limit being chosen because there are not enough events for a reliable fit above that value. The size of each fit range is computed as twenty standard deviations of the mass resolution for the signal point considered. The low end of the lowest fit range is 95 GeV, which evades the $Z \rightarrow \mu\mu$ and $Z \rightarrow \tau\tau$ background components, and thus allows the model to be simplified to just the function family h of Eq. (3). For each mass point we define two BDT bins, 0.3–0.7 and 0.7–1.0, combining the upper two bins used for the $Z \rightarrow e\mu$ search because of their smaller populations.

VI. SEARCH FOR $Z \rightarrow e\tau$ AND $Z \rightarrow \mu\tau$

The $Z \rightarrow e\tau$ and $Z \rightarrow \mu\tau$ searches each consider hadronic and leptonic τ -daughter categories. In addition to the common selection described in Sec. IV, in all four categories we require $40 < m_{\ell\tau} < 170$ GeV, where $m_{\ell\tau}$ is the invariant mass of the system comprising a prompt light-lepton candidate and the visible daughter(s) of a tau lepton. To reduce backgrounds from $\tau\tau$, $W + \text{jets}$, WW , and $t\bar{t}$ production, we also require $m_T(\tau, p_T^{\text{miss}})/m_{\ell\tau} < 0.8$ and $m_T(\ell\tau, p_T^{\text{miss}}) < 70(90)$ GeV, in the hadronic (leptonic) final-state categories.

To further reduce the $\tau\tau$ background, the prompt light lepton candidate must satisfy $p_T > 28(20)$ GeV in the hadronic (leptonic) final-state categories. Events are rejected if they contain b -tagged jets, based on the tight (loose) working point of the b jet ID for the hadronic (leptonic) τ -decay categories, to suppress backgrounds from t quark processes. The QCD multijet background becomes significant at low electron p_T in the $Z \rightarrow e\tau_\mu$ category, and so we require $p_T^e > p_T^\mu$ in that category. Additionally, the light leptons are required to have $|\eta^\ell| < 2.2(2.4)$ in the hadronic (leptonic) τ decay categories, for compatibility with the $\tau\tau$ background estimation method described in Sec. VIA 1 below.

The search strategy for final states involving a tau lepton needs to take into account that the Z boson can be only partially reconstructed because of the neutrino(s) in the τ decay. The signal and background spectra in $m_{\ell\tau}$ are less distinct than those of $m_{e\mu}$. Here the background components are modeled separately, using data in CRs to the extent possible. New variables derived from the visible daughters are developed to approximate the mass of the Z boson. These and other discriminating variables are again combined into a BDT to optimize the signal extraction.

A. Background estimates for the $Z \rightarrow e\tau$ and $Z \rightarrow \mu\tau$ channels

The $Z \rightarrow e\tau$ and $Z \rightarrow \mu\tau$ searches rely on signal and background templates (binned PDFs). The dominant background

sources are $\tau\tau$ production and those arising from misidentified leptons. Other sources are leptonic W boson decays, such as those from WW and $t\bar{t}$ processes.

1. Background from $\tau\tau$

The production of tau lepton pairs, most importantly from DY $Z \rightarrow \tau\tau$, is a background in all of the $Z \rightarrow \ell\tau$ channels. Neutrinos associated with the decays of both tau leptons contribute to p_T^{miss} , and the visible invariant mass spectrum of the dilepton system peaks below the Z boson mass, with a high-side tail. The τ candidate in a $Z \rightarrow e\tau$ or $Z \rightarrow \mu\tau$ signal event would be kinematically identical to one from the $Z \rightarrow \tau\tau$ decay, while the prompt lepton would have generally higher p_T than the daughter from a leptonically decaying τ lepton in the $Z \rightarrow \tau\tau$ event, and there would be no contribution to p_T^{miss} from that leg of the Z boson decay.

The SM processes that produce tau lepton pairs are identical to those that produce muon pairs, and the latter are very cleanly reconstructed in the detector. Therefore we use an “embedding” technique [80] to model $\tau\tau$ backgrounds, except for $H \rightarrow \tau\tau$ production, which is treated separately. In this method, we remove the recorded detector deposits associated with the muons from a data sample of muon pair production, and substitute those derived from simulated $\tau\tau$ events, matching the kinematic properties of the simulated tau lepton with those of the muon it replaces. The result is a hybrid data sample where the τ decay and reconstruction are simulated, but the τ production kinematics and the rest of the event (pileup, p_T^{miss} , jets, etc.) are taken directly from data. This approach significantly reduces simulation uncertainties, where the only remaining simulation-based uncertainties are those related to the decay and reconstruction of the tau leptons.

2. The τ_h misID background

The background from jets misidentified as τ_h ($j \rightarrow \tau_h$) is estimated in the $\mu\tau_h$ and $e\tau_h$ channels using a “misID-factor” method, similar to those used in Refs. [81,82]. The method uses an application region (AR) enriched in $j \rightarrow \tau_h$ events, which has the same kinematic selection as the SR but with an inverted jet-suppression ID (“antijet-ID”) requirement (very loose and not tight, using working points of the discriminator [40]). Shape-dependent transfer factors (TFs) are used to relate the rate of the $j \rightarrow \tau_h$ events in the AR to an estimate of the corresponding background rate in the SR.

Since the TFs depend on properties that might be experimentally inaccessible (e.g., the underlying parton flavor), and depend on the background process (e.g., $W + \text{jets}$, QCD multijets), they are measured separately for each background source. An overall TF is then evaluated by combining these factors weighted by the process composition in the CR:

$$f(\vec{\beta}) = \sum_{p \in \text{processes}} c_p(\vec{\beta}) f_p(\vec{\beta}).$$

The parameters $\vec{\beta}$ are derived from the kinematic properties of the τ_h and prompt ℓ candidates along with p_T^{miss} , and the functions f , f_p , and c_p are, respectively, the overall TF to be evaluated for a given event in the AR, the TF for process p , and the expected composition fraction of process p in the AR.

The process-dependent TFs f_p are measured for $W + \text{jets}$, QCD multijet, and $t\bar{t}$ background. The $W + \text{jets}$ TFs are measured in a high- $m_T(\ell\tau, p_T^{\text{miss}})$ region, and the QCD multijet TFs are measured in a region with SS $\ell\tau_h$ pairs and $0.05 < I_{\text{rel}}^\ell < 0.15$. The $t\bar{t}$ TFs are determined using simulation, as there is not a suitable measurement region in data. These measurement regions are referred to as the determination regions. The weights c_p are estimated as functions of $m_T(\ell\tau, p_T^{\text{miss}})$, using simulated $j \rightarrow \tau_h$ events for $W + \text{jets}$ and $t\bar{t}$; for QCD multijet background we use an SS data CR, correcting for non-QCD contributions in this CR by subtracting their simulated yields.

The f_p TFs are measured as functions of the τ_h candidate p_T independently for each of four τ_h identified decay modes, one- or three-prong with zero or one π^0 . There are also subleading kinematic dependencies in the TFs. Corrections for these are derived by comparing the observation with the estimate in the tight antijet-ID subset of the determination region. This is done for the light-lepton p_T , the $\tau_h |\eta|$, and $|\Delta\phi(\ell, p_T^{\text{miss}})|$. Additional corrections to account for the CR selection bias are measured in alternate CRs and applied as functions of I_{rel}^ℓ , $m_{\ell\tau}$, and the BDT score defined in Sec. VIC.

3. Light-lepton misID background

Electron and muon candidates in the $e\tau_\mu$ and $\mu\tau_e$ selections arising from misID of other particles are estimated using a method similar to the $j \rightarrow \tau_h$ estimation. This background arises mainly from $W + \text{jets}$ events, where one lepton comes from the W boson and one is a misidentified jet or a nonprompt lepton originating from a jet, and from QCD multijet events, where each lepton is either a misidentified jet or a nonprompt lepton originating from a jet.

The AR for the light-lepton misID background is the same as the SR, except with an SS selection. The determination regions for the TFs are the same as for the SRs and ARs, except with a loose isolation requirement ($0.15 < I_{\text{rel}}^\mu < 0.50$) on the muon candidate. The TFs are measured as functions of $\Delta R(e, \mu)$ separately in categories of 0, 1, and > 1 jets. Subleading corrections to the (p_T^e, p_T^μ) distribution are measured in this determination region, and CR bias corrections for the (p_T^e, p_T^μ) and BDT score (Sec. VIC) distributions are measured in a loose electron isolation region.

4. Backgrounds estimated from simulation

The background from leptonic W decays in $t\bar{t}$ and WW production is relevant in all search channels, though it is most impactful at high mass in the $e\mu$ data channel, and is estimated from simulation. The embedded $\tau\tau$ samples already include the small contribution from $t\bar{t}$ and WW decays to $\tau\tau$, and so these are omitted from the simulated background model. The remaining $t\bar{t}$ and WW background processes are kept small by the event selection, which exploits several properties that distinguish them from signal. Both $t\bar{t}$ and WW have large p_T^{miss} associated with each lepton leg, unlike the signal processes, and the $t\bar{t}$ background is significantly suppressed by the veto of events with b -tagged jets.

There are also small background contributions from other diboson and triboson processes, but these have small cross sections and are suppressed by the rejection of events with additional light leptons. The $H \rightarrow \tau\tau$ and $H \rightarrow WW$ processes are also modeled with simulation, because these processes yield too few $\mu\mu p_T^{\text{miss}}$ events in data for application of the τ embedding method.

Photons misidentified as electrons and light leptons misidentified as τ_h candidates are also modeled using simulated samples. The background from $DY \ Z \rightarrow ee$ and $Z \rightarrow \mu\mu$ events where one light lepton is misidentified as a τ_h candidate must be carefully separated from the signal as the prompt-lepton candidate leg is identical to that of the CLFV Z signal.

B. Discriminating variables for $Z \rightarrow e\tau$ and $Z \rightarrow \mu\tau$

Because of the neutrinos in the τ decay, the invariant mass of the visible daughters in the $Z \rightarrow \ell\tau$ channels is a less effective discriminating variable than in the construction of $Z \rightarrow e\mu$ against the backgrounds, of which $Z \rightarrow \tau\tau$ is the largest. However, we can exploit the facts that since the decaying boson is much more massive than the τ , the τ is typically produced with a significant Lorentz boost, leading to a highly collimated decay, and that one of the Z boson daughters for the signal decay is a light lepton unaccompanied by neutrinos. We utilize two different approaches to take advantage of these features.

1. Collinear mass estimate

The first approach is the collinear mass m^{col} , derived with the collimated τ decay approximation, in which $\vec{p}_T^{\text{miss}}$ projected onto the visible tau lepton $\vec{p}_T$ equals the $\vec{p}_T$ of the neutrino(s). With the further assumption that the neutrino three-momentum lies along the three-momentum of the visible τ daughter(s), we then obtain the neutrino four-vector,

$$p^\nu = \frac{\vec{p}_T^{\text{miss}} \cdot \hat{p}_T^{\text{vis}}}{p_T^{\text{vis}}} p^{\text{vis}}, \quad (4)$$




FIG. 5. The $\mu\tau_h$ (upper) and $\mu\tau_e$ (lower) m^{col} distributions, for the data (black markers with bars showing the statistical uncertainties) and the simulated backgrounds (filled stacked histograms). The hatched blue histogram shows the shape of the signal, normalized to a branching fraction of 10^{-3} , for comparison. The lower panel shows the ratio of the data to simulated yields, with the statistical (combined systematic and statistical) uncertainty in the simulated yield indicated by the filled (hatched) gray band.

where p^{vis} is the four-momentum of the visible τ daughter(s), and p_T^{vis} and $\hat{p}_T^{\text{vis}}$ are the magnitude and direction unit vector of its transverse momentum components, respectively. The sign of $\vec{p}_T^{\text{miss}} \cdot \hat{p}_T^{\text{vis}}$ indicates whether the neutrino vector is parallel or antiparallel to the visible τ daughter's momentum. The collinear mass is then the mass of the $p^\ell + p^{\text{vis}} + p^\nu = p^\ell + p^\tau$ system. The m^{col} distribution is shown

in Fig. 5 for the $Z \rightarrow \mu\tau$ search. The true $\Delta\phi$ between the visible τ decay products and the associated neutrinos is typically small, but because of uncertainties in the p_T^{miss} reconstruction this angle can be larger than $\pi/2$, leading to antiparallel neutrino configurations.

2. The tau lepton momentum scale variables $\alpha_{1,2}$

Another approach is to use collinear approximations for the τ decays along with assumptions [83] that the Z boson is on shell and produced at threshold, i.e., that the Z boson daughters' p_T vanishes. The latter constraint implies equality of the Z daughters' p_T , $p_T^{(1)} = p_T^{(2)}$. This relation can be restated in terms of the four-momentum $p^{(i)}$ of the tau lepton and that of its daughters, defined as $p^{\text{vis},i} = p^{(i)}/\alpha_i$:

$$\alpha_1 p_T^{\text{vis},1} = \alpha_2 p_T^{\text{vis},2}. \quad (5)$$

The on-shell Z boson assumption combined with four-momentum conservation in the decay provides a second relation between the α_i :

$$\begin{aligned} m_Z^2 &= m_1^2 + m_2^2 + 2p^{(1)} \cdot p^{(2)} \approx 2p^{(1)} \cdot p^{(2)} \\ &= 2\alpha_1\alpha_2 p^{\text{vis},1} \cdot p^{\text{vis},2}. \end{aligned} \quad (6)$$

From these equations we find

$$\alpha_{1,2} = \sqrt{\left(\frac{p_T^{\text{vis},(2,1)}}{p_T^{\text{vis},(1,2)}}\right) \frac{m_Z^2}{2p^{\text{vis},1} \cdot p^{\text{vis},2}}}. \quad (7)$$

For the signal decay $Z \rightarrow \ell\tau$ we have $\alpha_1 = \alpha_\ell = 1$, whereas for the $Z \rightarrow \tau\tau$ background we have in general $\alpha_i > 1$ for both Z boson daughters. Figure 6 shows these estimates for $Z \rightarrow \mu\tau_h$, where for the signal events α_μ is centered at one, as expected, and provides considerable discrimination against both $Z \rightarrow \tau\tau$ and τ_h misID backgrounds.

C. The BDTs for the $Z \rightarrow e\tau$ and $Z \rightarrow \mu\tau$ decay channels

A BDT is trained for each final state of each signal, for a total of four BDTs ($Z \rightarrow e\tau_h$, $Z \rightarrow e\tau_\mu$, $Z \rightarrow \mu\tau_h$, and $Z \rightarrow \mu\tau_e$). The ROOT TMVA toolkit [84] is used for this training. The BDTs are constructed with input features that describe the kinematic properties of the Z boson candidate and of p_T^{miss} :

- (1) $m_{\ell\tau}$;
- (2) m^{col} ;
- (3) α_ℓ and α_τ ;
- (4) p_T^ℓ/p_T^τ ;
- (5) $p_T^{\ell\tau}$;
- (6) $\Delta\phi_{\ell\tau}$;
- (7) $m_T(\ell, p_T^{\text{miss}})$;
- (8) $\Delta\phi(\tau, p_T^{\text{miss}})$; and
- (9) p_T^{jet} .




FIG. 6. Distributions of $\mu\tau_h$ signal and estimated backgrounds in α_μ (upper), α_τ (lower), for the data (black markers with bars showing the statistical uncertainties) and the simulated backgrounds (filled stacked histograms). The hatched blue histogram shows the shape of the signal, normalized to a branching fraction of 10^{-3} , for comparison. The lower panel shows the ratio of the data to simulated yields, with the statistical (combined systematic and statistical) uncertainty in the simulated yield indicated by the filled (hatched) gray band.

The leading jet p_T is included here, as b quark initiated jets in $t\bar{t}$ decays that either fail the b tagging or are beyond the b tagging region of $|\eta| < 2.4$ typically have higher p_T than jets associated with the signal process.

The BDT training uses $\tau\tau$ simulations rather than the embedding samples. Data events from the loose antijet ID (SS) region are used in the training to model the $j \rightarrow \tau_h$ (nonprompt $e\mu$) background in the hadronic (leptonic) τ

TABLE I. Regions in $m_{\ell\tau}$ for the $Z \rightarrow \mu\tau$ and $Z \rightarrow e\tau$ fits.

Region	$m_{\ell\tau}$ bounds [GeV]	
	$Z \rightarrow \ell\tau_h$	$Z \rightarrow \ell\tau_\ell$
$\tau\tau$	[40, 60]	[40, 50]
Signal-like	[60, 85]	[50, 100]
$Z \rightarrow \ell\ell$	[85, 100]	...
misID ($j \rightarrow \tau_h$ and Nonprompt $e\mu$)	[100, 170]	[100, 170]

categories. Events used for training, whether simulated events or CR data, are excluded from use in the background template estimates, where the remaining events or data are scaled to compensate for the removed events. To improve the discrimination against rare $Z \rightarrow \ell\ell$ background events in which one of the leptons is misidentified as one of a different flavor, these events are given a larger weight in the training. The BDT score distribution $O(x)$ is transformed using the signal BDT score cumulative distribution function $p(x) = \int_{-\infty}^x O(x')dx' / \int_{-\infty}^{+\infty} O(x')dx'$ to produce a useful shape that is less impacted by changes in the underlying machine-learning technique or hyperparameters.

D. The $Z \rightarrow e\tau$ and $Z \rightarrow \mu\tau$ mass categories

Several of the backgrounds in the $Z \rightarrow e\tau$ and $Z \rightarrow \mu\tau$ search have large systematic uncertainties that impact the search sensitivity. These are:

- (1) $\tau\tau$ embedding uncertainties;
- (2) $j \rightarrow \tau_h$ and nonprompt $e\mu$ estimates from data; and
- (3) $Z \rightarrow \ell\ell \rightarrow \ell\tau_h$ misID rate uncertainties.

To improve background uncertainty constraints and isolate background- or signal-enriched regions, the $Z \rightarrow e\tau$ and $Z \rightarrow \mu\tau$ fits are split into the $m_{\ell\tau}$ mass regions described in Table I. The leptonic τ signal decays typically have lower mass than the τ_h decays because of the extra neutrino produced, which leads to differences in the $\tau\tau$ region selection. Additionally, $Z \rightarrow \ell\ell$ is a significant background only in the hadronic channels, so there is no CR for these processes in the leptonic channels. Events with low transformed BDT score are removed from the final analysis, and the transformed BDT binning is chosen to ensure that each bin contains an adequate number of events for the final fit.

VII. SYSTEMATIC UNCERTAINTIES

The main sources of systematic uncertainty arise from the simulation (e.g., simulated object reconstruction efficiency or energy scale), from the background model uncertainties (e.g., validity of the model in the SR), and statistical uncertainties because of limited event counts in the simulated samples and CRs. These uncertainties are discussed in the following paragraphs and summarized in Table II.

In the $Z \rightarrow e\mu$ search, the most significant sources are the purely statistical uncertainties from the parameters of the background functions in the fit. These are followed by

uncertainties arising from the discrete collection of background functions included in the envelope. The signal model uncertainties are negligible compared with those of the background model.

The largest sources of systematic uncertainty in the $Z \rightarrow e\tau$ and $Z \rightarrow \mu\tau$ searches are the statistical uncertainties in the background model templates and the light-lepton energy scale uncertainties. The model-template statistical uncertainties arise largely from the $Z \rightarrow ee$ and $Z \rightarrow \mu\mu$ simulation samples used to estimate the $\ell \rightarrow \tau_h$ backgrounds, along with those from the nonprompt $e\mu$ background estimate based on CRs in the data, where the subtraction of prompt leptons from the data has a large statistical uncertainty. The statistical uncertainties in the $j \rightarrow \tau_h$ and embedded $\tau\tau$ estimates are much smaller

TABLE II. Sources of uncertainty and their impacts on the measured branching fraction, for each of the Z decay channels, and on the product of the production cross section and branching fraction for the Z' resonance scan. The uncertainty ranges for the Z' resonance scan are ordered from the lowest to the highest Z' mass point. Entries to which the specified uncertainty does not apply are denoted with “...”.

Uncertainty source	$\mathcal{B}(Z \rightarrow e\mu)$ [10^{-8}]	$\mathcal{B}(Z \rightarrow e\tau)$ [10^{-6}]	$\mathcal{B}(Z \rightarrow \mu\tau)$ [10^{-6}]	$\sigma(pp \rightarrow Z' + X) \times \mathcal{B}(Z' \rightarrow e\mu)$ [fb]
Electron and muon ID and trigger	0.1	0.3	0.1	0.1–0.01
Electron energy scale	0.2	3.5	0.2	0.6–0.01
Muon energy scale	0.1	0.1	0.9	0.3–0.01
Tau ID	...	0.3	0.2	...
Tau energy scale	...	0.4	0.4	...
$(e, \mu) \rightarrow \tau_h$	...	0.4	0.1	...
Jet energy, p_T^{miss}	0.3	0.8	0.2	0.1–0.01
b tagging	< 0.1	0.2	0.2	< (0.1–0.01)
Pileup	0.1	0.2	0.1	< (0.1–0.01)
Integrated luminosity	0.3	0.1	0.1	0.1–0.01
Theory	0.1	0.5	0.2	0.1–(< 0.01)
Parametric background	7.9	...	...	0.3–0.06
Envelope	2.5	...	...	0.1–0.04
$Z \rightarrow \mu\mu$ yield	0.5	...	...	...
Embedding energy resolution	...	1.1	0.2	...
Embedding normalization	...	1.5	0.2	...
$j \rightarrow \tau_h$	...	2.8	0.9	...
Nonprompt $e\mu$	...	2.7	0.4	...
Template event counts	< 0.1	3.1	1.5	0.1–(< 0.01)
Total systematic	8.3	5.7	2.2	0.8–0.08
Statistical	5.6	1.9	1.6	1.9–0.18




FIG. 7. For the $Z \rightarrow e\mu$ search, the invariant mass fit results for the BDT score ranges 0.3–0.7 (upper), 0.7–0.9 (middle), and 0.9–1.0 (lower). In each plot, the upper panel shows the data (points with bars showing the statistical uncertainties) together with the fit distribution curve (red) and its separate signal (blue dotted) $Z \rightarrow \mu\mu$ (yellow dash-dotted) and continuum background (purple dashed) components, and the lower panel shows the deviations of the data from the fit function divided by the data uncertainty.




FIG. 8. Upper limits at 95% CL on the branching fraction $B(Z \rightarrow e\mu)$, for each BDT score range and for the final combined fit. The observed limits are denoted by the markers, while the expected limits with their 68% and 95% uncertainties are denoted by the horizontal dashed lines and green and yellow bands, respectively.

because of their larger CR populations. The impact of the electron and muon energy scale uncertainties is large because the p_T of the prompt ℓ is highly discriminating against leptonic $Z \rightarrow \tau\tau$ decays, as well as being incorporated into several of the variables in the BDTs [dilepton mass and p_T , collinear mass, α_i , $m_T(\ell, p_T^{\text{miss}})$, and the p_T ratio]. The uncertainties in the background estimate from CRs in data are also significant, mainly because of the corrections for the kinematic variation in the TF

measurement regions and the statistical uncertainties in these measurement regions.

The normalization of the simulated signal and minor background yields depends on the integrated luminosity, pileup corrections, and the theoretical modeling. The integrated luminosities for the 2016, 2017, and 2018 data-taking years have individual uncertainties of 1.2–2.5% [85–87], while the overall uncertainty for the 2016–2018 period is 1.6%. To evaluate the uncertainty associated with the pileup reweighting, we vary the value of the total inelastic cross section by 5% [88]. Uncertainties arising from the theoretical model include the cross sections, the parton distribution functions, and the renormalization and factorization scales. These uncertainties are propagated to the observed signal yields and are summarized in the row labeled “Theory” in Table II.

All uncertainties are included in the maximum likelihood fit to extract the branching fractions, which is implemented with the Combine statistical analysis package [89]. For an uncertainty that does not contain shape information, the uncertainty is implemented as a rate uncertainty with a log-normal PDF. Uncertainties that impact the shape of the signal or background model are handled differently in the $Z \rightarrow e\mu$ and $Z \rightarrow \ell\tau$ searches. In the $Z \rightarrow e\mu$ search, the effect is split into a rate uncertainty and an impact on the pure shape effect. A nuisance parameter is added as a shift in the corresponding model parameter drawn from a Gaussian PDF centered on zero and with a width defined by the systematic source’s effect on that model parameter. In the $Z \rightarrow e\tau$ and $Z \rightarrow \mu\tau$ searches, shape uncertainties are introduced via “continuous morphing” [90] parametrized by upward and downward uncertainty templates. In the case of the $Z \rightarrow e\mu$ background envelope, the discrete background function index parameter is treated as a free parameter. The fit selects

TABLE III. The measured branching fraction with its significance (signif.) and observed and expected 95% CL upper limits, for each of the $Z \rightarrow e\mu$, $Z \rightarrow e\tau$, and $Z \rightarrow \mu\tau$ decay channels. The prior best published limits are also given for comparison. Included are results for the separate BDT bins for $Z \rightarrow e\mu$ and the separate τ decay subchannels for $Z \rightarrow e\tau$ and $Z \rightarrow \mu\tau$.

Channel	Branching fraction	Signif. [σ]	Observed (expected) limit	Prior (expected) limit
$Z \rightarrow e\mu$	$-0.1_{-1.0}^{+1.0} \times 10^{-7}$	−0.1	$1.9(2.0_{-0.6}^{+0.8}) \times 10^{-7}$	$2.6(2.4) \times 10^{-7}$ [23]
$0.3 < \text{BDT} < 0.7$	$-3.4_{-2.6}^{+2.8} \times 10^{-7}$	−1.2	$5.7(8.2_{-2.0}^{+2.5}) \times 10^{-7}$	...
$0.7 < \text{BDT} < 0.9$	$0.4_{-1.4}^{+1.5} \times 10^{-7}$	+0.3	$3.2(2.9_{-0.8}^{+1.2}) \times 10^{-7}$	...
$0.9 < \text{BDT} < 1.0$	$0.0_{-2.0}^{+1.5} \times 10^{-7}$	+0.0	$3.0(3.0_{-0.8}^{+1.2}) \times 10^{-7}$	...
$Z \rightarrow e\tau$	$3.2_{-6.0}^{+6.1} \times 10^{-6}$	+0.5	$13.8(11.4_{-3.2}^{+4.7}) \times 10^{-6}$	$5.0(6.0) \times 10^{-6}$ [22]
$Z \rightarrow e\tau_h$	$6.3_{-8.2}^{+8.4} \times 10^{-6}$	+0.8	$21.3(16.1_{-4.6}^{+6.7}) \times 10^{-6}$	$8.1(8.1) \times 10^{-6}$ [22]
$Z \rightarrow e\tau_\mu$	$1.2_{-8.1}^{+7.9} \times 10^{-6}$	+0.2	$16.2(15.3_{-4.2}^{+6.1}) \times 10^{-6}$	$7.0(8.9) \times 10^{-6}$ [22]
$Z \rightarrow \mu\tau$	$7.5_{-2.7}^{+2.7} \times 10^{-6}$	+2.7	$12.0(5.3_{-1.5}^{+2.1}) \times 10^{-6}$	$6.5(5.3) \times 10^{-6}$ [22]
$Z \rightarrow \mu\tau_h$	$7.2_{-2.8}^{+2.8} \times 10^{-6}$	+2.5	$11.9(5.6_{-1.6}^{+2.2}) \times 10^{-6}$	$9.5(6.1) \times 10^{-6}$ [22]
$Z \rightarrow \mu\tau_e$	$7.5_{-7.9}^{+7.1} \times 10^{-6}$	+1.0	$19.5(14.4_{-3.8}^{+5.1}) \times 10^{-6}$	$7.2(10.0) \times 10^{-6}$ [22]

the background function that maximizes the likelihood at any given point, profiling this discrete nuisance parameter [78].

VIII. RESULTS

The branching fraction is extracted from a maximum likelihood fit of the background plus signal model to the data distribution in $m_{e\mu}$, for the $Z \rightarrow e\mu$ search, or in the BDT discriminant, for the $Z \rightarrow \ell\tau$ searches.

For the $Z \rightarrow e\mu$ search, the parameters of the background model developed from the $m_{e\mu}$ sidebands (Sec. V B) are free

in the fit to the full distributions. The distributions for separate fits in each BDT region are shown along with the data in Fig. 7. No excess of events beyond the SM expectation is observed. We set upper limits on the branching fraction following the modified frequentist construction CL_s [91–93] implemented with the CMS statistical analysis tool Combine [89]. The observed and expected upper limits are shown in Fig. 8 and Table III from the fits in each BDT score category and for the joint fit combining all categories, along with the 68% and 95% uncertainties. The observed limit of $\mathcal{B}(Z \rightarrow e\mu) < 1.9 \times 10^{-7}$ represents the most stringent direct limit to date on this process.




FIG. 9. Transformed BDT score distributions for the $Z \rightarrow e\tau_h$ channels in the $m_{e\tau}$ ranges: (upper left) 40–60 GeV, “ $\tau\tau$ ”; (upper right) 60–85 GeV, “signal-like”; (lower left) 85–100 GeV, “ $Z \rightarrow \ell\ell$ ”; (lower right) 100–170 GeV, “misID.” In each plot, the upper panel shows the data (points), the total yield from the signal + background fit (red open histogram), the signal component (blue open histogram), and the background components (stacked filled histograms). The middle panel shows the ratio to the background of the data (points with bars showing the statistical uncertainty in the data) and the combined signal + background (blue dotted histogram). The shaded band shows the systematic uncertainty in the background estimate. The lower panel shows the pull defined in the text (light-blue filled histogram).




FIG. 10. Transformed BDT score distributions for the $Z \rightarrow e\tau_\mu$ channels in the $m_{e\tau}$ ranges: (upper left) 40–50 GeV, “ $\tau\tau$ ”; (upper right) 50–100 GeV, “signal-like”; (lower) 100–170 GeV, “misID.” In each plot, the upper panel shows the data (points), the total yield from the signal + background fit (red open histogram), the signal component (blue open histogram), and the background components (stacked filled histograms). The middle panel shows the ratio to the background of the data (points with bars showing the statistical uncertainty in the data) and the combined signal + background (blue dotted histogram). The shaded band shows the systematic uncertainty in the background estimate. The lower panel shows the pull defined in the text (light-blue filled histogram).

The transformed BDT score distributions in the invariant mass and tau lepton final-state categories are shown in Figs. 9 and 10 for the $Z \rightarrow e\tau$ search and in Figs. 11 and 12 for the $Z \rightarrow \mu\tau$ search. The pull appearing in the distributions in the lower panels for each BDT bin is the difference between the observed and fit yields divided by the uncertainty $\sqrt{\sigma_{\text{obs}}^2 - \sigma_{\text{fit}}^2}$. The minus sign in the definition of this uncertainty accounts for the correlation between the observed and fit yields, which arises because the observed yield is included in the fit [94]. The large pulls that appear in some bins in the lower row of Fig. 9 reflect non-Gaussian

uncertainties in one or more of the nuisance parameters. The observed yield in the $e\tau$ ($\mu\tau$) channel is consistent with the SM background prediction at the 0.5(2.7) σ level. The observed and expected 95% CL upper limits on the branching fractions for $Z \rightarrow e\tau$ and $Z \rightarrow \mu\tau$ are shown in Fig. 13 and Table III, along with the 68% and 95% uncertainties in the expected limit.

Figure 14 shows the $e\mu$ mass scan fit results for two example mass points. No statistically significant excess is observed over the SM background. We report 95% CL upper limits on the production cross section $\sigma(pp \rightarrow Z' + X)$




FIG. 11. Transformed BDT score distributions for the $Z \rightarrow \mu\tau_h$ channels in the $m_{\mu\tau}$ ranges: (upper left) 40–60 GeV, “ $\tau\tau$ ”; (upper right) 60–85 GeV, “signal-like”; (lower left) 85–100 GeV, “ $Z \rightarrow \ell\ell$ ”; (lower right) 100–170 GeV, “misID.” In each plot, the upper panel shows the data (points), the total yield from the signal + background fit (red open histogram), the signal component (blue open histogram), and the background components (stacked filled histograms). The middle panel shows the ratio to the background of the data (points with bars showing the statistical uncertainty in the data) and the combined signal + background (blue dotted histogram). The shaded band shows the systematic uncertainty in the background estimate. The lower panel shows the pull defined in the text (light-blue filled histogram).

times the branching fraction $\mathcal{B}(Z' \rightarrow e\mu)$ to the $e\mu$ final state. The observed and expected limits for the entire mass scan in the $e\mu$ final state are shown in Fig. 15. All observed deviations from the background-only hypothesis have local p -values greater than 1%. The global p -value is greater than 39% after accounting for the look-elsewhere effect [95].

The search presented here for CLFV Z boson decays represents an approximately 20% improvement in sensitivity to the direct $Z \rightarrow e\mu$ decay process, with respect to the best previous search [23]. The $Z \rightarrow \mu\tau$ search has

similar sensitivity as the prior limit, coming mainly from the $Z \rightarrow \mu\tau_h$ channel. The observed $Z \rightarrow \mu\tau$ limit is weaker than the expected one, mainly because of the 2.5σ excess we find in the $Z \rightarrow \mu\tau_h$ channel. The $Z \rightarrow e\tau$ search presented here is limited by high electron trigger thresholds, lower electron identification efficiencies, and large rates of $j \rightarrow \tau_h$ and $e \rightarrow \tau_h$ backgrounds, as compared with the $Z \rightarrow \mu\tau$ search, and the $Z \rightarrow e\tau$ search in Ref. [22]. Additionally, the electron energy scale has much larger uncertainties in the $\tau\tau$ embedding simulation that significantly impacts the $Z \rightarrow e\tau$ search sensitivity.




FIG. 12. Transformed BDT score distributions for the $Z \rightarrow \mu\tau_e$ channels in the $m_{\mu\tau}$ ranges: (upper left) 40–50 GeV, “ $\tau\tau$ ”; (upper right) 50–100 GeV, “signal-like”; (lower) 100–170 GeV, “misID”. In each plot, the upper panel shows the data (points), the total yield from the signal + background fit (red open histogram), the signal component (blue open histogram), and the background components (stacked filled histograms). The middle panel shows the ratio to the background of the data (points with bars showing the statistical uncertainty in the data) and the combined signal + background (blue dotted histogram). The shaded band shows the systematic uncertainty in the background estimate. The lower panel shows the pull defined in the text (light-blue filled histogram).

As seen in Fig. 15, the $Z' \rightarrow e\mu$ search yields an expected 95% CL upper limit on $\sigma(pp \rightarrow Z' + X)\mathcal{B}(Z' \rightarrow e\mu)$ of 4 fb at 110 GeV and 0.4 fb at 500 GeV, with an approximately exponential relation between the mass and the expected sensitivity. The observed limits are consistent with these. Searches for $Z' \rightarrow e\mu$ over various mass ranges

above 200 GeV have been reported by ATLAS [96,97] and CMS [98,99]. Where the ranges overlap, the present limits represent an improvement of at least a factor of five over these, and are either the first or the most stringent expected limits to date on the $Z' \rightarrow e\mu$ process within the covered mass range.

IX. SUMMARY

A search is presented for flavor violating decays of the Z boson to charged leptons, and for the presence of a heavier vector boson Z' exhibiting such decays. The data from proton-proton collisions at $\sqrt{s} = 13$ TeV were collected with the CMS detector at the LHC, and correspond to an integrated luminosity of 138 fb^{-1} . The specific decay modes considered are $Z^{(\prime)} \rightarrow e\mu$, $Z \rightarrow e\tau$, and $Z \rightarrow \mu\tau$. No significant excess of events over backgrounds from standard model processes is observed. Observed (expected)

upper limits of 1.9×10^{-7} , 1.38×10^{-5} , and 1.20×10^{-5} (2.0×10^{-7} , 1.14×10^{-5} , and 0.53×10^{-5}) at 95% confidence level are set on the branching fractions for $Z \rightarrow e\mu$, $Z \rightarrow e\tau$, and $Z \rightarrow \mu\tau$, respectively. The limit for $Z \rightarrow e\mu$ is the most restrictive to date, while for $Z \rightarrow \mu\tau$ the sensitivity in terms of the expected limit is the same as that of the previous best limit. All of these limits are consistent with expectations from the standard model, and with constraints inferred from low-energy experimental limits. For Z' boson masses in the range 110–500 GeV, upper limits are set on the cross section times the branching fraction to $e\mu$ that




FIG. 13. Observed and expected 95% CL upper limit by category, as well as for the final combined fit, for the $Z \rightarrow e\tau$ (upper) and $Z \rightarrow \mu\tau$ (lower) searches. The observed limits are denoted by the markers, while the expected limits with their 68% and 95% uncertainties are denoted by the horizontal dashed lines and green and yellow bands, respectively.




FIG. 14. Distributions in $m_{e\mu}$ for the scan points 111 GeV (upper row) and 496 GeV (lower row) from the Z' search. In each row the BDT score range for the left (right) plot is 0.3–0.7 (0.7–1.0). In each plot, the upper panel shows the data (points with bars showing the statistical uncertainty) together with the fit distribution curve (red solid) and its separate signal (blue dotted) and background (red dotted) components, the middle panel shows the background subtracted data with the fit signal distribution, and the lower panel shows the deviations of the data from the fit function divided by the fit uncertainty.




FIG. 15. Expected and observed 95% CL upper limits on $\sigma(pp \rightarrow Z' + X)\mathcal{B}(Z' \rightarrow e\mu)$ for Z' masses between 110 and 500 GeV. The solid black line connects filled circles representing the observed upper limits at the scan points, and the dashed line with filled error bands shows the expected limit.

range from 0.3 to 7 fb, and are the most restrictive to date for this mass range. Future studies can benefit from additional data, since even the systematic uncertainties arise mainly from statistical ones in control samples.

ACKNOWLEDGMENTS

We congratulate our colleagues in the CERN accelerator departments for the excellent performance of the LHC and thank the technical and administrative staffs at CERN and at other CMS institutes for their contributions to the success of the CMS effort. In addition, we gratefully acknowledge the computing centers and personnel of the Worldwide LHC Computing Grid and other centers for delivering so effectively the computing infrastructure essential to our analyses. Finally, we acknowledge the enduring support for the construction and operation of the LHC, the CMS detector, and the supporting computing infrastructure provided by the following funding agencies: SC (Armenia), BMBWF and FWF (Austria); FNRS and FWO (Belgium); CNPq, CAPES, FAPERJ, FAPERGS, and FAPESP (Brazil); MES and BNSF (Bulgaria); CERN; CAS, MoST, and NSFC (China); MINCIENCIAS (Colombia); MSES and CSF (Croatia); RIF (Cyprus); SENESCYT (Ecuador); ERC PRG, TARISTU24-TK10 and MoER TK202 (Estonia); Academy of Finland, MEC, and HIP (Finland); CEA and CNRS/IN2P3 (France); SRNSF (Georgia); BMBF, DFG, and HGF (Germany); GSRI (Greece); NKFIH (Hungary); DAE and DST (India); IPM (Iran); SFI (Ireland); INFN (Italy); MSIT and NRF (Republic of Korea); MES (Latvia); LMTLT (Lithuania); MOE and UM (Malaysia); BUAP, CINVESTAV, CONACYT, LNS, SEP, and UASLP-FAI (Mexico); MOS (Montenegro); MBIE (New Zealand); PAEC (Pakistan); MES and NSC (Poland); FCT (Portugal); MESTD (Serbia); MICIU/AEI and PCTI (Spain); MOSTR (Sri Lanka); Swiss Funding Agencies (Switzerland); MST (Taipei); MHESI and NSTDA (Thailand); TUBITAK and TENMAK (Türkiye); NASU (Ukraine); STFC (United Kingdom); Individuals have received support from the Marie-Curie program and the European Research Council and Horizon 2020 Grant, Contracts No. 675440, No. 724704, No. 752730, No. 758316, No. 765710, No. 824093, No. 101115353, No. 101002207, No. 101001205, and COST Action CA16108 (European Union); the Leventis Foundation; the Alfred P. Sloan Foundation; the Alexander von Humboldt Foundation; the Science Committee, Project No. 22rl-037 (Armenia); the Fonds pour la Formation à la Recherche dans l'Industrie et dans l'Agriculture (FRIA-Belgium); the Beijing Municipal Science and Technology Commission,

No. Z191100007219010, the Fundamental Research Funds for the Central Universities, the Ministry of Science and Technology of China under Grant No. 2023YFA1605804, and the Natural Science Foundation of China under Grant No. 12061141002 (China); the Ministry of Education, Youth and Sports (MEYS) of the Czech Republic; the Shota Rustaveli National Science Foundation, Grant No. FR-22-985 (Georgia); the Deutsche Forschungsgemeinschaft (DFG), among others, under Germany's Excellence Strategy—EXC 2121 “Quantum Universe”—390833306, and under Project No. 400140256—GRK2497; the Hellenic Foundation for Research and Innovation (HFRI), Project No. 2288 (Greece); the Hungarian Academy of Sciences, the New National Excellence Program—ÚNKP, the NKFIH Research Grants No. K 131991, No. K 133046, No. K 138136, No. K 143460, No. K 143477, No. K 146913, No. K 146914, No. K 147048, No. 2020-2.2.1-ED-2021-00181, No. TKP2021-NKTA-64, and No. 2021-4.1.2-NEMZ_KI-2024-00036 (Hungary); the Council of Science and Industrial Research, India; ICSC—National Research Center for High Performance Computing, Big Data and Quantum Computing, FAIR—Future Artificial Intelligence Research, and CUP I53D23001070006 (Mission 4 Component 1), funded by the NextGenerationEU program (Italy); the Latvian Council of Science; the Ministry of Education and Science, Project No. 2022/WK/14, and the National Science Center, contracts Opus 2021/41/B/ST2/01369 and 2021/43/B/ST2/01552 (Poland); the Fundação para a Ciência e a Tecnologia, Grant No. CEECIND/01334/2018 (Portugal); MICIU/AEI/10.13039/501100011033, ERDF/EU, “European Union NextGenerationEU/PRTR,” and Programa Severo Ochoa del Principado de Asturias (Spain); the Chulalongkorn Academic into Its 2nd Century Project Advancement Project, and the National Science, Research and Innovation Fund via the Program Management Unit for Human Resources & Institutional Development, Research and Innovation, Grant No. B39G680009 (Thailand); the Kavli Foundation; the Nvidia Corporation; the SuperMicro Corporation; the Welch Foundation, Contract No. C-1845; and the Weston Havens Foundation (USA).

DATA AVAILABILITY

Release and preservation of data used by the CMS Collaboration as the basis for publications is guided by the CMS data preservation, re-use and open access policy [100].

- [1] Y. Fukuda *et al.* (Super-Kamiokande Collaboration), Measurement of the flux and zenith angle distribution of upward through going muons by Super-Kamiokande, *Phys. Rev. Lett.* **82**, 2644 (1999).
- [2] S. Fukuda *et al.* (Super-Kamiokande Collaboration), Solar B-8 and hep neutrino measurements from 1258 days of Super-Kamiokande data, *Phys. Rev. Lett.* **86**, 5651 (2001).
- [3] Q. R. Ahmad *et al.* (SNO Collaboration), Direct evidence for neutrino flavor transformation from neutral current interactions in the Sudbury Neutrino Observatory, *Phys. Rev. Lett.* **89**, 011301 (2002).
- [4] K. Eguchi *et al.* (KamLAND Collaboration), First results from KamLAND: Evidence for reactor anti-neutrino disappearance, *Phys. Rev. Lett.* **90**, 021802 (2003).
- [5] J. I. Illana and T. Riemann, Charged lepton flavor violation from massive neutrinos in Z decays, *Phys. Rev. D* **63**, 053004 (2001).
- [6] W. J. Marciano, T. Mori, and J. M. Roney, Charged lepton flavor violation experiments, *Annu. Rev. Nucl. Part. Sci.* **58**, 315 (2008).
- [7] L. Calibbi and G. Signorelli, Charged lepton flavour violation: An experimental and theoretical introduction, *Riv. Nuovo Cimento* **41**, 71 (2018).
- [8] P. Langacker, The physics of heavy Z' gauge bosons, *Rev. Mod. Phys.* **81**, 1199 (2009).
- [9] C. Cornella, P. Paradisi, and O. Sumensari, Hunting for ALPs with lepton flavor violation, *J. High Energy Phys.* **01** (2020) 158.
- [10] M. Bauer, M. Neubert, S. Renner, M. Schnubel, and A. Thamm, Flavor probes of axion-like particles, *J. High Energy Phys.* **09** (2022) 056.
- [11] K. Afanaciev *et al.* (MEG II Collaboration), A search for $\mu^+ \rightarrow e^+ \gamma$ with the first dataset of the MEG II experiment, *Eur. Phys. J. C* **84**, 216 (2024); **84**, 216(E) (2024).
- [12] U. Bellgardt *et al.* (SINDRUM Collaboration), Search for the decay $\mu^+ \rightarrow e^+ e^+ e^-$, *Nucl. Phys. B* **299**, 1 (1988).
- [13] W. H. Bertl *et al.* (SINDRUM II Collaboration), A search for muon to electron conversion in muonic gold, *Eur. Phys. J. C* **47**, 337 (2006).
- [14] W. Love *et al.* (CLEO Collaboration), Search for lepton flavor violation in Upsilon decays, *Phys. Rev. Lett.* **101**, 201601 (2008).
- [15] B. Aubert *et al.* (BABAR Collaboration), Search for the reactions $e^+ e^- \rightarrow \mu^+ \tau^-$ and $e^+ e^- \rightarrow e^+ \tau^-$, *Phys. Rev. D* **75**, 031103 (2007).
- [16] B. Aubert *et al.* (BABAR Collaboration), Searches for lepton flavor violation in the decays $\tau^\pm \rightarrow e^\pm \gamma$ and $\tau^\pm \rightarrow \mu^\pm \gamma$, *Phys. Rev. Lett.* **104**, 021802 (2010).
- [17] K. Hayasaka *et al.* (Belle Collaboration), Search for lepton flavor violating τ decays into three leptons with 719 million produced $\tau^+ \tau^-$ pairs, *Phys. Lett. B* **687**, 139 (2010).
- [18] S. Nussinov, R. D. Peccei, and X. M. Zhang, On unitarity based relations between various lepton family violating processes, *Phys. Rev. D* **63**, 016003 (2001).
- [19] R. Akers *et al.* (OPAL Collaboration), A search for lepton flavor violating Z^0 decays, *Z. Phys. C* **67**, 555 (1995).
- [20] P. Abreu *et al.* (DELPHI Collaboration), Search for lepton flavor number violating Z^0 decays, *Z. Phys. C* **73**, 243 (1997).
- [21] ATLAS Collaboration, Search for charged-lepton-flavour violation in Z-boson decays with the ATLAS detector, *Nat. Phys.* **17**, 819 (2021).
- [22] ATLAS Collaboration, Search for lepton-flavor-violation in Z-boson decays with τ leptons with the ATLAS detector, *Phys. Rev. Lett.* **127**, 271801 (2022).
- [23] ATLAS Collaboration, Search for the charged-lepton-flavor-violating decay $Z \rightarrow e\mu$ in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *Phys. Rev. D* **108**, 032015 (2023).
- [24] HEPData record for this analysis (2025), [10.17182/hepdata.161021](https://hepdata.net/record/10.17182/hepdata.161021).
- [25] CMS Collaboration, The CMS experiment at the CERN LHC, *J. Instrum.* **3**, S08004 (2008).
- [26] CMS Collaboration, Development of the CMS detector for the CERN LHC Run 3, *J. Instrum.* **19**, P05064 (2023).
- [27] CMS Tracker Collaboration, Description and performance of track and primary-vertex reconstruction with the CMS tracker, *J. Instrum.* **9**, P10009 (2014).
- [28] W. Adam *et al.* (CMS Tracker Group Collaboration), The CMS phase-1 pixel detector upgrade, *J. Instrum.* **16**, P02027 (2020).
- [29] CMS Collaboration, Track impact parameter resolution for the full pseudo rapidity coverage in the 2017 dataset with the CMS Phase-1 Pixel detector, CMS Detector Performance Summary Report No. CMS-DP-2020-049, 2020, <https://cds.cern.ch/record/2743740>.
- [30] CMS Collaboration, Technical proposal for the Phase-II upgrade of the Compact Muon Solenoid, CMS Technical Proposal Reports No. CERN-LHCC-2015-010, No. CMS-TDR-15-02, 2015, <http://cds.cern.ch/record/2020886>.
- [31] CMS Collaboration, Particle-flow reconstruction and global event description with the CMS detector, *J. Instrum.* **12**, P10003 (2017).
- [32] M. Cacciari, G. P. Salam, and G. Soyez, The anti- k_T jet clustering algorithm, *J. High Energy Phys.* **04** (2008) 063.
- [33] M. Cacciari, G. P. Salam, and G. Soyez, FastJet user manual, *Eur. Phys. J. C* **72**, 1896 (2012).
- [34] CMS Collaboration, Pileup mitigation at CMS in 13 TeV data, *J. Instrum.* **15**, P09018 (2020).
- [35] CMS Collaboration, Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV, *J. Instrum.* **12**, P02014 (2016).
- [36] CMS Collaboration, Identification of b-quark jets with the CMS experiment, *J. Instrum.* **8**, P04013 (2012).
- [37] CMS Collaboration, Performance of the DeepJet b tagging algorithm using 41.9/fb of data from proton-proton collisions at 13 TeV with Phase 1 CMS detector, CMS Detector Performance Summary Report No. CMS-DP-2018-058, 2018, <http://cds.cern.ch/record/2646773>.
- [38] E. Bols, J. Kieseler, M. Verzetti, M. Stoye, and A. Stakia, Jet flavour classification using DeepJet, *J. Instrum.* **15**, P12012 (2020).
- [39] CMS Collaboration, Performance of reconstruction and identification of τ leptons decaying to hadrons and ν_τ in pp collisions at $\sqrt{s} = 13$ TeV, *J. Instrum.* **13**, P10005 (2018).

- [40] CMS Collaboration, Identification of hadronic tau lepton decays using a deep neural network, *J. Instrum.* **17**, P07023 (2022).
- [41] CMS Collaboration, ECAL 2016 refined calibration and Run 2 summary plots, CMS Detector Performance Summary Report No. CMS-DP-2020-021, 2020, <https://cds.cern.ch/record/2717925>.
- [42] CMS Collaboration, Electron and photon reconstruction and identification with the CMS experiment at the CERN LHC, *J. Instrum.* **16**, P05014 (2020).
- [43] CMS Collaboration, Performance of the CMS muon detector and muon reconstruction with proton-proton collisions at $\sqrt{s} = 13$ TeV, *J. Instrum.* **13**, P06015 (2018).
- [44] D. Bertolini, P. Harris, M. Low, and N. Tran, Pileup per particle identification, *J. High Energy Phys.* **10** (2014) 059.
- [45] CMS Collaboration, Performance of missing transverse momentum reconstruction in proton-proton collisions at $\sqrt{s} = 13$ TeV using the CMS detector, *J. Instrum.* **14**, P07004 (2019).
- [46] CMS Collaboration, Performance of the CMS Level-1 trigger in proton-proton collisions at $\sqrt{s} = 13$ TeV, *J. Instrum.* **15**, P10017 (2020).
- [47] CMS Collaboration, The CMS trigger system, *J. Instrum.* **12**, P01020 (2016).
- [48] CMS Collaboration, Performance of the CMS high-level trigger during LHC Run 2, *J. Instrum.* **19**, P11021 (2024).
- [49] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H.-S. Shao, T. Stelzer, P. Torrielli, and M. Zaro, The automated computation of tree-level and next-to-leading order differential cross sections, and their matching to parton shower simulations, *J. High Energy Phys.* **07** (2014) 079.
- [50] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H.-S. Shao, T. Stelzer, P. Torrielli, and M. Zaro, Comparative study of various algorithms for the merging of parton showers and matrix elements in hadronic collisions, *Eur. Phys. J. C* **53**, 473 (2008).
- [51] P. Nason, A new method for combining NLO QCD with shower Monte Carlo algorithms, *J. High Energy Phys.* **11** (2004) 040.
- [52] S. Frixione, P. Nason, and C. Oleari, Matching NLO QCD computations with parton shower simulations: The POWHEG method, *J. High Energy Phys.* **11** (2007) 070.
- [53] S. Alioli, P. Nason, C. Oleari, and E. Re, A general framework for implementing NLO calculations in shower Monte Carlo programs: The POWHEG BOX, *J. High Energy Phys.* **06** (2010) 043.
- [54] S. Alioli, P. Nason, C. Oleari, and E. Re, NLO single-top production matched with shower in POWHEG: s - and t -channel contributions, *J. High Energy Phys.* **09** (2009) 111; **02** (2010) 011(E).
- [55] E. Re, Single-top Wt -channel production matched with parton showers using the POWHEG method, *Eur. Phys. J. C* **71**, 1547 (2011).
- [56] S. Alioli, P. Nason, C. Oleari, and E. Re, NLO Higgs boson production via gluon fusion matched with shower in POWHEG, *J. High Energy Phys.* **04** (2009) 002.
- [57] P. Nason and C. Oleari, NLO Higgs boson production via vector-boson fusion matched with shower in POWHEG, *J. High Energy Phys.* **02** (2010) 037.
- [58] T. Melia, P. Nason, R. Rontsch, and G. Zanderighi, W^+W^- , WZ and ZZ production in the POWHEG BOX, *J. High Energy Phys.* **11** (2011) 078.
- [59] P. Nason and G. Zanderighi, W^+W^- , WZ and ZZ production in the POWHEG-BOX-V2, *Eur. Phys. J. C* **74**, 2702 (2014).
- [60] G. Luisoni, P. Nason, C. Oleari, and F. Tramontano, $HW^\pm/HZ + 0$ and 1 jet at NLO with the POWHEG BOX interfaced to GoSam and their merging within MiNLO, *J. High Energy Phys.* **10** (2013) 083.
- [61] T. Sjöstrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C. O. Rasmussen, and P. Z. Skands, An introduction to PYTHIA 8.2, *Comput. Phys. Commun.* **191**, 159 (2015).
- [62] K. Melnikov and F. Petriello, Electroweak gauge boson production at hadron colliders through $\mathcal{O}(\alpha_s^2)$, *Phys. Rev. D* **74**, 114017 (2006).
- [63] M. Czakon and A. Mitov, Top++: A program for the calculation of the top-pair cross-section at hadron colliders, *Comput. Phys. Commun.* **185**, 2930 (2014).
- [64] N. Kidonakis, Top quark production, in *Helmholtz International Summer School on Physics of Heavy Quarks and Hadrons* (DESY, Hamburg, Germany, 2014), p. 139, 10.3204/DESY-PROC-2013-03/Kidonakis.
- [65] J. M. Campbell, R. K. Ellis, and C. Williams, Vector boson pair production at the LHC, *J. High Energy Phys.* **07** (2011) 018.
- [66] T. Gehrmann, M. Grazzini, S. Kallweit, P. Maierhöfer, A. von Manteuffel, S. Pozzorini, D. Rathlev, and L. Tancredi, W^+W^- production at hadron colliders in next to next to leading order QCD, *Phys. Rev. Lett.* **113**, 212001 (2014).
- [67] CMS Collaboration, Event generator tunes obtained from underlying event and multiparton scattering measurements, *Eur. Phys. J. C* **76**, 155 (2016).
- [68] CMS Collaboration, Extraction and validation of a new set of CMS PYTHIA8 tunes from underlying-event measurements, *Eur. Phys. J. C* **80**, 4 (2020).
- [69] R. Frederix and S. Frixione, Merging meets matching in MC@NLO, *J. High Energy Phys.* **12** (2012) 061.
- [70] R. D. Ball *et al.* (NNPDF Collaboration), Parton distributions for the LHC Run II, *J. High Energy Phys.* **04** (2015) 040.
- [71] R. D. Ball *et al.* (NNPDF Collaboration), Parton distributions from high-precision collider data, *Eur. Phys. J. C* **77**, 663 (2017).
- [72] R. D. Ball, V. Bertone, S. Carrazza, L. Del Debbio, S. Forte, A. Guffanti, N. P. Hartland, and J. Rojo (NNPDF Collaboration), Parton distributions with QED corrections, *Nucl. Phys.* **B877**, 290 (2013).
- [73] S. Agostinelli *et al.* (GEANT4 Collaboration), Geant4—a simulation toolkit, *Nucl. Instrum. Methods Phys. Res., Sect. A* **506**, 250 (2003).
- [74] S. Navas *et al.* (Particle Data Group), Review of particle physics, *Phys. Rev. D* **110**, 030001 (2024).
- [75] T. Chen and C. Guestrin, XGBoost: A scalable tree boosting system, in *Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and*

- Data Mining*, KDD '16 (Association for Computing Machinery, New York, NY, USA, 2016), p. 785.
- [76] M. J. Oreglia, A study of the reactions $\psi' \rightarrow \gamma\gamma\psi$, Master's thesis, Stanford University, 1980, SLAC Report SLAC-R-236.
- [77] J. E. Gaiser, Charmonium spectroscopy from radiative decays of the J/ψ and ψ' , Master's thesis, Stanford University, 1982, SLAC Report SLAC-0255.
- [78] P. D. Dauncey, M. Kenzie, N. Wardle, and G. J. Davies, Handling uncertainties in background shapes: The discrete profiling method, *J. Instrum.* **10**, P04015 (2014).
- [79] R. A. Fisher, On the mathematical foundations of theoretical statistics, *Phil. Trans. R. Soc.* **222**, 309 (1922).
- [80] CMS Collaboration, An embedding technique to determine $\tau\tau$ backgrounds in proton-proton collision data, *J. Instrum.* **14**, P06032 (2019).
- [81] CMS Collaboration, Measurements of Higgs boson production in the decay channel with a pair of τ leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV, *Eur. Phys. J. C* **83**, 562 (2023).
- [82] CMS Collaboration, Search for lepton-flavor violating decays of the Higgs boson in the $\mu\tau$ and $e\tau$ final states in proton-proton collisions at $\sqrt{s} = 13$ TeV, *Phys. Rev. D* **104**, 032013 (2021).
- [83] S. Davidson, S. Lacroix, and P. Verdier, LHC sensitivity to lepton flavour violating Z boson decays, *J. High Energy Phys.* **09** (2012) 092.
- [84] A. Hocker *et al.* (TMVA Collaboration), TMVA—Toolkit for multivariate data analysis, Technical Report CERN-OPEN-2007-007, 2007, <https://cds.cern.ch/record/1019880>.
- [85] CMS Collaboration, Precision luminosity measurement in proton-proton collisions at $\sqrt{s} = 13$ TeV in 2015 and 2016 at CMS, *Eur. Phys. J. C* **81**, 800 (2021).
- [86] CMS Collaboration, CMS luminosity measurement for the 2017 data-taking period at $\sqrt{s} = 13$ TeV, CMS Physics Analysis Summary, Report No. CMS-PAS-LUM-17-004, 2018, <https://cds.cern.ch/record/2621960/>.
- [87] CMS Collaboration, CMS luminosity measurement for the 2018 data-taking period at $\sqrt{s} = 13$ TeV, CMS Physics Analysis Summary, Report No. CMS-PAS-LUM-18-002, 2019, <https://cds.cern.ch/record/2676164/>.
- [88] CMS Collaboration, Measurement of the inelastic proton-proton cross section at $\sqrt{s} = 13$ TeV, *J. High Energy Phys.* **07** (2018) 161.
- [89] CMS Collaboration, The CMS statistical analysis and combination tool: COMBINE, *Comput. Softw. Big Sci.* **8**, 19 (2024).
- [90] J. S. Conway, Incorporating nuisance parameters in likelihoods for multisource spectra, in *Proceedings of the PHYSTAT 2011 Workshop* (CERN, Geneva, Switzerland, 2011), p. 115, [10.5170/CERN-2011-006.115](https://cds.cern.ch/record/105170/).
- [91] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, Asymptotic formulae for likelihood-based tests of new physics, *Eur. Phys. J. C* **71**, 1554 (2011); **73**, 2501(E) (2013).
- [92] T. Junk, Confidence level computation for combining searches with small statistics, *Nucl. Instrum. Methods Phys. Res., Sect. A* **434**, 435 (1999).
- [93] A. L. Read, Presentation of search results: The CL_s technique, *J. Phys. G* **28**, 2693 (2002).
- [94] F. E. James, *Statistical Methods in Experimental Physics 2nd Edition* (World Scientific Publishing Co. Pte. Ltd., Singapore, 2006).
- [95] G. Cowan, Statistics for searches at the LHC, in *LHC Phenomenology*, edited by E. Gardi, N. Glover, and R. Aidan, Scottish Graduate Series (Springer, Cham, 2015).
- [96] ATLAS Collaboration, Search for a heavy neutral particle decaying to $e\mu$, $e\tau$, or $\mu\tau$ in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector, *Phys. Rev. Lett.* **115**, 031801 (2015).
- [97] ATLAS Collaboration, Search for new phenomena in different-flavour high-mass dilepton final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *Eur. Phys. J. C* **76**, 541 (2016).
- [98] CMS Collaboration, Search for lepton flavour violating decays of heavy resonances and quantum black holes to an $e\mu$ pair in proton-proton collisions at $\sqrt{s} = 8$ TeV, *Eur. Phys. J. C* **76**, 317 (2016).
- [99] CMS Collaboration, Search for heavy resonances and quantum black holes in $e\mu$, $e\tau$, and $\mu\tau$ final states in proton-proton collisions at $\sqrt{s} = 13$ TeV, *J. High Energy Phys.* **05** (2023) 227.
- [100] CMS data availability statement, [10.7483/OPENDATA.CMS.1BNU.8V1W](https://cds.cern.ch/record/107483/).

A. Hayrapetyan,¹ V. Makarenko¹, A. Tumasyan^{1,b}, W. Adam², J. W. Andrejkovic,² L. Benato², T. Bergauer², M. Dragicevic², C. Giordano,² P. S. Hussain², M. Jeitler^{2,c}, N. Krammer², A. Li², D. Liko², I. Mikulec², J. Schieck^{2,c}, R. Schöfbeck^{2,c}, D. Schwarz², M. Shooshari,² M. Sonawane², W. Waltenberger², C.-E. Wulz^{2,c}, T. Janssen³, H. Kwon³, D. Ocampo Henao³, T. Van Laer³, P. Van Mechelen³, J. Bierkens⁴, N. Breugelmans,⁴ J. D'Hondt⁴, S. Dansana⁴, A. De Moor⁴, M. Delcourt⁴, F. Heyen⁴, Y. Hong⁴, P. Kashko⁴, S. Lowette⁴, I. Makarenko⁴, D. Müller⁴, J. Song⁴, S. Tavernier⁴, M. Tytgat^{4,d}, G. P. Van Onsem⁴, S. Van Putte⁴, D. Vannerom⁴, B. Bilin⁵, B. Clerbaux⁵, A. K. Das,⁵ I. De Bruyn⁵, G. De Lentdecker⁵, H. Evard⁵, L. Favart⁵, P. Gianneios⁵, A. Khalilzadeh,⁵ F. A. Khan⁵, A. Malara⁵, M. A. Shahzad,⁵ L. Thomas⁵, M. Vanden Bemden⁵, C. Vander Velde⁵, P. Vanlaer⁵, F. Zhang⁵, M. De Coen⁶, D. Dobur⁶, G. Gokbulut⁶, J. Knolle⁶, L. Lambrecht⁶, D. Marckx⁶, K. Skovpen⁶, N. Van Den Bossche⁶, J. van der Linden⁶, J. Vandenbroeck⁶, L. Wezenbeek⁶, S. Bein⁷, A. Benecke⁷, A. Bethani⁷, G. Bruno⁷, A. Cappati⁷, J. De Favereau De Jeneret⁷, C. Delaere⁷

A. Giammanco⁷, A. O. Guzel⁷, V. Lemaitre⁷, J. Lidrych⁷, P. Malek⁷, P. Mastrapasqua⁷, S. Turkcapar⁷,
 G. A. Alves⁸, M. Barroso Ferreira Filho⁸, E. Coelho⁸, C. Hensel⁸, T. Menezes De Oliveira⁸, C. Mora Herrera^{8,e},
 P. Rebello Teles⁸, M. Soeiro⁸, E. J. Tonelli Manganote^{8,f}, A. Vilela Pereira^{8,e}, W. L. Aldá Júnior⁹,
 H. Brandao Malbouisson⁹, W. Carvalho⁹, J. Chinellato^{9,g}, M. Costa Reis⁹, E. M. Da Costa⁹, G. G. Da Silva^{9,h},
 D. De Jesus Damiao⁹, S. Fonseca De Souza⁹, R. Gomes De Souza⁹, S. S. Jesus⁹, T. Laux Kuhn^{9,h}, M. Macedo⁹,
 K. Mota Amarilo⁹, L. Mundim⁹, H. Nogima⁹, J. P. Pinheiro⁹, A. Santoro⁹, A. Sznajder⁹, M. Thiel⁹,
 F. Torres Da Silva De Araujo^{9,i}, C. A. Bernardes^{10,h}, T. R. Fernandez Perez Tomei¹⁰, E. M. Gregores¹⁰,
 B. Lopes Da Costa¹⁰, I. Maletto Silverio¹⁰, P. G. Mercadante¹⁰, S. F. Novaes¹⁰, B. Orzari¹⁰, Sandra S. Padula¹⁰,
 V. Scheurer¹⁰, A. Aleksandrov¹¹, G. Antchev¹¹, P. Danev¹¹, R. Hadjiiska¹¹, P. Iaydjiev¹¹, M. Misheva¹¹,
 M. Shopova¹¹, G. Sultanov¹¹, A. Dimitrov¹², L. Litov¹², B. Pavlov¹², P. Petkov¹², A. Petrov¹², S. Keshri¹³,
 D. Laroze¹³, S. Thakur¹³, W. Brooks¹⁴, T. Cheng¹⁵, T. Javaid¹⁵, L. Wang¹⁵, L. Yuan¹⁵, Z. Hu¹⁶, Z. Liang¹⁶,
 J. Liu¹⁶, X. Wang¹⁶, G. M. Chen^{17,j}, H. S. Chen^{17,j}, M. Chen^{17,j}, Y. Chen¹⁷, Q. Hou¹⁷, X. Hou¹⁷, F. Iemmi¹⁷,
 C. H. Jiang¹⁷, A. Kapoor^{17,k}, H. Liao¹⁷, G. Liu¹⁷, Z.-A. Liu^{17,l}, J. N. Song^{17,l}, S. Song¹⁷, J. Tao¹⁷, C. Wang^{17,j},
 J. Wang¹⁷, H. Zhang¹⁷, J. Zhao¹⁷, A. Agapitos¹⁸, Y. Ban¹⁸, A. Carvalho Antunes De Oliveira¹⁸, S. Deng¹⁸,
 B. Guo¹⁸, Q. Guo¹⁸, C. Jiang¹⁸, A. Levin¹⁸, C. Li¹⁸, Q. Li¹⁸, Y. Mao¹⁸, S. Qian¹⁸, S. J. Qian¹⁸, X. Qin¹⁸, X. Sun¹⁸,
 D. Wang¹⁸, J. Wang¹⁸, H. Yang¹⁸, M. Zhang¹⁸, Y. Zhao¹⁸, C. Zhou¹⁸, S. Yang¹⁹, Z. You²⁰, K. Jaffel²¹, N. Lu²¹,
 G. Bauer^{22,m}, B. Li^{22,n}, H. Wang²², K. Yi^{22,o}, J. Zhang²², Y. Li²³, Z. Lin²⁴, C. Lu²⁴, M. Xiao^{24,p}, C. Avila²⁵,
 D. A. Barbosa Trujillo²⁵, A. Cabrera²⁵, C. Florez²⁵, J. Fraga²⁵, J. A. Reyes Vega²⁵, C. Rendón²⁶, M. Rodriguez²⁶,
 A. A. Ruales Barbosa²⁶, J. D. Ruiz Alvarez²⁶, N. Godinovic²⁷, D. Lelas²⁷, A. Sculac²⁷, M. Kovac²⁸,
 A. Petkovic²⁸, T. Sculac²⁸, P. Bargassa²⁹, V. Brigljevic²⁹, B. K. Chitroda²⁹, D. Ferencek²⁹, K. Jakovcic²⁹,
 A. Starodumov²⁹, T. Susa²⁹, A. Attikis³⁰, K. Christoforou³⁰, A. Hadjiagapiou³⁰, C. Leonidou³⁰, C. Nicolaou³⁰,
 L. Paizanos³⁰, F. Ptochos³⁰, P. A. Razis³⁰, H. Rykaczewski³⁰, H. Saka³⁰, A. Stepennov³⁰, M. Finger^{31,a},
 M. Finger Jr.³¹, A. Kveton³¹, E. Ayala³², E. Carrera Jarrin³³, H. Abdalla^{34,q}, Y. Assran^{34,r,s},
 M. Abdullah Al-Mashad³⁵, A. Hussein³⁵, H. Mohammed³⁵, K. Ehataht³⁶, M. Kadastik³⁶, T. Lange³⁶, C. Nielsen³⁶,
 J. Pata³⁶, M. Raidal³⁶, N. Seeba³⁶, L. Tani³⁶, A. Milieva³⁷, K. Osterberg³⁷, M. Voutilainen³⁷,
 N. Bin Norjoharuddeen³⁸, E. Brücken³⁸, F. Garcia³⁸, P. Inkaew³⁸, K. T. S. Kallonen³⁸, R. Kumar Verma³⁸,
 T. Lampén³⁸, K. Lassila-Perini³⁸, B. Lehtela³⁸, S. Lehti³⁸, T. Lindén³⁸, N. R. Mancilla Xinto³⁸,
 M. Myllymäki³⁸, M. m. Rantanen³⁸, S. Saariokari³⁸, N. T. Toikka³⁸, J. Tuominiemi³⁸, H. Kirschenmann³⁹,
 P. Luukka³⁹, H. Petrow³⁹, M. Besancon⁴⁰, F. Couderc⁴⁰, M. Dejardin⁴⁰, D. Denegri⁴⁰, P. Devouge⁴⁰, J. L. Faure⁴⁰,
 F. Ferri⁴⁰, S. Ganjour⁴⁰, P. Gras⁴⁰, G. Hamel de Monchenault⁴⁰, M. Kumar⁴⁰, V. Lohezic⁴⁰, J. Malcles⁴⁰,
 F. Orlandi⁴⁰, L. Portales⁴⁰, S. Ronchi⁴⁰, M. Ö. Sahin⁴⁰, A. Savoy-Navarro^{40,t}, P. Simkina⁴⁰, M. Titov⁴⁰,
 M. Tornago⁴⁰, F. Beaudette⁴¹, G. Boldrini⁴¹, P. Busson⁴¹, C. Charlot⁴¹, M. Chiusi⁴¹, T. D. Cuisset⁴¹,
 F. Damas⁴¹, O. Davignon⁴¹, A. De Wit⁴¹, T. Debnath⁴¹, I. T. Ehle⁴¹, B. A. Fontana Santos Alves⁴¹, S. Ghosh⁴¹,
 A. Gilbert⁴¹, R. Granier de Cassagnac⁴¹, L. Kalipoliti⁴¹, M. Manoni⁴¹, M. Nguyen⁴¹, S. Obraztsov⁴¹,
 C. Ochando⁴¹, R. Salerno⁴¹, J. B. Sauvan⁴¹, Y. Sirois⁴¹, G. Sokmen⁴¹, L. Urda Gómez⁴¹, A. Zabi⁴¹,
 A. Zghiche⁴¹, J.-L. Agram^{42,u}, J. Andrea⁴², D. Bloch⁴², J.-M. Brom⁴², E. C. Chabert⁴², C. Collard⁴²,
 G. Coulon⁴², S. Falke⁴², U. Goerlach⁴², R. Haeberle⁴², A.-C. Le Bihan⁴², M. Meena⁴², O. Poncet⁴², G. Saha⁴²,
 P. Vaucelle⁴², A. Di Florio⁴³, D. Amram⁴⁴, S. Beauceron⁴⁴, B. Blancon⁴⁴, G. Boudoul⁴⁴, N. Chanon⁴⁴,
 D. Contardo⁴⁴, P. Depasse⁴⁴, C. Dozen^{44,v}, H. El Mamouni⁴⁴, J. Fay⁴⁴, S. Gascon⁴⁴, M. Gouzevitch⁴⁴,
 C. Greenberg⁴⁴, G. Grenier⁴⁴, B. Ille⁴⁴, E. Jourdhuy⁴⁴, I. B. Laktineh⁴⁴, M. Lethuillier⁴⁴, B. Massoteau⁴⁴,
 L. Mirabito⁴⁴, A. Purohit⁴⁴, M. Vander Donckt⁴⁴, J. Xiao⁴⁴, G. Adamov⁴⁵, I. Lomidze⁴⁵, Z. Tsamalaidze^{45,w},
 V. Botta⁴⁶, S. Consuegra Rodríguez⁴⁶, L. Feld⁴⁶, K. Klein⁴⁶, M. Lipinski⁴⁶, D. Meuser⁴⁶, P. Nattland⁴⁶,
 V. Oppenländer⁴⁶, A. Pauls⁴⁶, D. Pérez Adán⁴⁶, N. Röwert⁴⁶, M. Teroerde⁴⁶, C. Daumann⁴⁷, S. Diekmann⁴⁷,
 A. Dodonova⁴⁷, N. Eich⁴⁷, D. Eliseev⁴⁷, F. Engelke⁴⁷, J. Erdmann⁴⁷, M. Erdmann⁴⁷, B. Fischer⁴⁷,
 T. Hebbeker⁴⁷, K. Hoepfner⁴⁷, F. Ivone⁴⁷, A. Jung⁴⁷, N. Kumar⁴⁷, M. y. Lee⁴⁷, F. Mausolf⁴⁷,
 M. Merschmeyer⁴⁷, A. Meyer⁴⁷, F. Nowotny⁴⁷, A. Pozdnyakov⁴⁷, W. Redjeb⁴⁷, H. Reithler⁴⁷, U. Sarkar⁴⁷,
 V. Sarkisovi⁴⁷, A. Schmidt⁴⁷, C. Seth⁴⁷, A. Sharma⁴⁷, J. L. Spah⁴⁷, V. Vaulin⁴⁷, S. Zaleski⁴⁷, M. R. Beckers⁴⁸,
 C. Dziwok⁴⁸, G. Flügge⁴⁸, N. Hoeflich⁴⁸, T. Kress⁴⁸, A. Nowack⁴⁸, O. Pooth⁴⁸, A. Stahl⁴⁸, A. Zotz⁴⁸,
 H. Aarup Petersen⁴⁹, A. Abel⁴⁹, M. Aldaya Martin⁴⁹, J. Alimena⁴⁹, S. Amoroso⁴⁹, Y. An⁴⁹, I. Andreev⁴⁹,

J. Bach⁴⁹, S. Baxter⁴⁹, M. Bayatmakou⁴⁹, H. Becerril Gonzalez⁴⁹, O. Behnke⁴⁹, A. Belvedere⁴⁹, F. Blekman^{49,x}, K. Borras^{49,y}, A. Campbell⁴⁹, S. Chatterjee⁴⁹, L. X. Coll Saravia⁴⁹, G. Eckerlin⁴⁹, D. Eckstein⁴⁹, E. Gallo^{49,x}, A. Geiser⁴⁹, V. Guglielmi⁴⁹, M. Guthoff⁴⁹, A. Hinzmann⁴⁹, L. Jeppe⁴⁹, M. Kasemann⁴⁹, C. Kleinwort⁴⁹, R. Kogler⁴⁹, M. Komm⁴⁹, D. Krücker⁴⁹, W. Lange⁴⁹, D. Leyva Pernia⁴⁹, K.-Y. Lin⁴⁹, K. Lipka^{49,z}, W. Lohmann^{49,aa}, J. Malvaso⁴⁹, R. Mankel⁴⁹, I.-A. Melzer-Pellmann⁴⁹, M. Mendizabal Morentin⁴⁹, A. B. Meyer⁴⁹, G. Milella⁴⁹, K. Moral Figueroa⁴⁹, A. Mussgiller⁴⁹, L. P. Nair⁴⁹, J. Niedziela⁴⁹, A. Nürnberg⁴⁹, J. Park⁴⁹, E. Ranken⁴⁹, A. Raspereza⁴⁹, D. Rastorguev⁴⁹, L. Rygaard⁴⁹, M. Scham^{49,bb,y}, S. Schnake^{49,y}, P. Schütze⁴⁹, C. Schwanenberger^{49,x}, D. Selivanova⁴⁹, K. Sharko⁴⁹, M. Shchedrolosiev⁴⁹, D. Stafford⁴⁹, M. Torkian⁴⁹, F. Vazzoler⁴⁹, A. Ventura Barroso⁴⁹, R. Walsh⁴⁹, D. Wang⁴⁹, Q. Wang⁴⁹, K. Wichmann⁴⁹, L. Wiens^{49,y}, C. Wissing⁴⁹, Y. Yang⁴⁹, S. Zakharov⁴⁹, A. Zimmermann Castro Santos⁴⁹, A. Albrecht⁵⁰, A. R. Alves Andrade⁵⁰, M. Antonello⁵⁰, S. Bollweg⁵⁰, M. Bonanomi⁵⁰, K. El Morabit⁵⁰, Y. Fischer⁵⁰, M. Frahm⁵⁰, E. Garutti⁵⁰, A. Grohsjean⁵⁰, J. Haller⁵⁰, D. Hundhausen⁵⁰, H. R. Jabusch⁵⁰, G. Kasieczka⁵⁰, P. Keicher⁵⁰, R. Klanner⁵⁰, W. Korcar⁵⁰, T. Kramer⁵⁰, C. c. Kuo⁵⁰, V. Kutzner⁵⁰, F. Labe⁵⁰, J. Lange⁵⁰, A. Lobanov⁵⁰, L. Moureaux⁵⁰, M. Mrowietz⁵⁰, A. Nigamova⁵⁰, K. Nikolopoulos⁵⁰, Y. Nissan⁵⁰, A. Paasch⁵⁰, K. J. Pena Rodriguez⁵⁰, N. Prouvost⁵⁰, T. Quadfasel⁵⁰, B. Raciti⁵⁰, M. Rieger⁵⁰, D. Savoie⁵⁰, J. Schindler⁵⁰, P. Schleper⁵⁰, M. Schröder⁵⁰, J. Schwandt⁵⁰, M. Sommerhalder⁵⁰, H. Stadie⁵⁰, G. Steinbrück⁵⁰, A. Tews⁵⁰, R. Ward⁵⁰, B. Wiederspan⁵⁰, M. Wolf⁵⁰, S. Brommer⁵¹, E. Butz⁵¹, Y. M. Chen⁵¹, T. Chwalek⁵¹, A. Dierlamm⁵¹, G. G. Dincer⁵¹, U. Elicabuk⁵¹, N. Faltermann⁵¹, M. Giffels⁵¹, A. Gottmann⁵¹, F. Hartmann^{51,cc}, R. Hofsaess⁵¹, M. Horzela⁵¹, U. Husemann⁵¹, J. Kieseler⁵¹, M. Klute⁵¹, R. Kunnilan Muhammed Rafeek⁵¹, O. Lavoryk⁵¹, J. M. Lawhorn⁵¹, A. Lintuluoto⁵¹, S. Maier⁵¹, M. Mormile⁵¹, Th. Müller⁵¹, E. Pfeffer⁵¹, M. Presilla⁵¹, G. Quast⁵¹, K. Rabbertz⁵¹, B. Regnery⁵¹, R. Schmieder⁵¹, N. Shadskiy⁵¹, I. Shvetsov⁵¹, H. J. Simonis⁵¹, L. Sowa⁵¹, L. Stockmeier⁵¹, K. Tauqeer⁵¹, M. Toms⁵¹, B. Topko⁵¹, N. Trevisani⁵¹, C. Verstege⁵¹, T. Voigtländer⁵¹, R. F. Von Cube⁵¹, J. Von Den Driesch⁵¹, M. Wassmer⁵¹, R. Wolf⁵¹, W. D. Zeuner⁵¹, X. Zuo⁵¹, G. Anagnostou⁵², G. Daskalakis⁵², A. Kyriakis⁵², A. Papadopoulos^{52,cc}, A. Stakia⁵², G. Melachroinos⁵³, Z. Painesis⁵³, I. Paraskevas⁵³, N. Saoulidou⁵³, K. Theofilatos⁵³, E. Tziaferi⁵³, K. Vellidis⁵³, I. Zisopoulos⁵³, T. Chatzistavrou⁵⁴, G. Karapostoli⁵⁴, K. Kousouris⁵⁴, E. Siamarkou⁵⁴, G. Tsiopolitis⁵⁴, I. Bestintzanos⁵⁵, I. Evangelou⁵⁵, C. Foudas⁵⁵, P. Katsoulis⁵⁵, P. Kokkas⁵⁵, P. G. Kosmoglou Kioseoglou⁵⁵, N. Manthos⁵⁵, I. Papadopoulos⁵⁵, J. Strologas⁵⁵, D. Druzhkin⁵⁶, C. Hajdu⁵⁶, D. Horvath^{56,dd,ee}, K. Márton⁵⁶, A. J. Rádl^{56,ff}, F. Sikler⁵⁶, V. Veszpremi⁵⁶, M. Csanád⁵⁷, K. Farkas⁵⁷, A. Fehérkúti^{57,gg}, M. M. A. Gadallah^{57,hh}, Á. Kadlecik⁵⁷, G. Pásztor⁵⁷, G. I. Veres⁵⁷, B. Ujvari⁵⁸, G. Zilizi⁵⁸, G. Bencze⁵⁹, S. Czellar⁵⁹, J. Molnar⁵⁹, Z. Szillasi⁵⁹, T. Csorgo^{60,gg}, F. Nemes^{60,gg}, T. Novak⁶⁰, I. Szanyi^{60,ii}, S. Bansal⁶¹, S. B. Beri⁶¹, V. Bhatnagar⁶¹, G. Chaudhary⁶¹, S. Chauhan⁶¹, N. Dhingra^{61,jj}, A. Kaur⁶¹, A. Kaur⁶¹, H. Kaur⁶¹, M. Kaur⁶¹, S. Kumar⁶¹, T. Sheokand⁶¹, J. B. Singh⁶¹, A. Singla⁶¹, A. Bhardwaj⁶², A. Chhetri⁶², B. C. Choudhary⁶², A. Kumar⁶², A. Kumar⁶², M. Naimuddin⁶², S. Phor⁶², K. Ranjan⁶², M. K. Saini⁶², S. Acharya^{63,kk}, B. Gumber^{63,kk}, B. Sahu^{63,kk}, S. Mukherjee⁶⁴, S. Baradia⁶⁵, S. Bhattacharya⁶⁵, S. Das Gupta⁶⁵, S. Dutta⁶⁵, S. Dutta⁶⁵, S. Sarkar⁶⁵, M. M. Ameen⁶⁶, P. K. Behera⁶⁶, S. Chatterjee⁶⁶, G. Dash⁶⁶, A. Dattamuni⁶⁶, P. Jana⁶⁶, P. Kalbhor⁶⁶, S. Kamble⁶⁶, J. R. Komaragiri^{66,ll}, T. Mishra⁶⁶, P. R. Pujahari⁶⁶, N. R. Saha⁶⁶, A. K. Sikdar⁶⁶, R. K. Singh⁶⁶, P. Verma⁶⁶, S. Verma⁶⁶, A. Vijay⁶⁶, B. K. Sirasva⁶⁷, L. Bhatt⁶⁸, S. Dugad⁶⁸, G. B. Mohanty⁶⁸, M. Shelake⁶⁸, P. Suryadevara⁶⁸, A. Bala⁶⁹, S. Banerjee⁶⁹, S. Barman^{69,mm}, R. M. Chatterjee⁶⁹, M. Guchait⁶⁹, Sh. Jain⁶⁹, A. Jaiswal⁶⁹, B. M. Joshi⁶⁹, S. Kumar⁶⁹, M. Maity^{69,mm}, G. Majumder⁶⁹, K. Mazumdar⁶⁹, S. Parolia⁶⁹, R. Saxena⁶⁹, A. Thachayath⁶⁹, S. Bahinipati^{70,nn}, D. Maity^{70,oo}, P. Mal⁷⁰, K. Naskar^{70,oo}, A. Nayak^{70,oo}, S. Nayak⁷⁰, K. Pal⁷⁰, R. Raturi⁷⁰, P. Sadangi⁷⁰, S. K. Swain⁷⁰, S. Varghese^{70,oo}, D. Vats^{70,oo}, A. Alpina⁷¹, S. Dube⁷¹, P. Hazarika⁷¹, B. Kansal⁷¹, A. Laha⁷¹, R. Sharma⁷¹, S. Sharma⁷¹, K. Y. Vaish⁷¹, S. Ghosh⁷², H. Bakhshiansohi^{73,pp}, A. Jafari^{73,qq}, V. Sedighzadeh Dalavi⁷³, M. Zeinali^{73,rr}, S. Bashiri⁷⁴, S. Chenarani^{74,ss}, S. M. Etesami⁷⁴, Y. Hosseini⁷⁴, M. Khakzad⁷⁴, E. Khazaie⁷⁴, M. Mohammadi Najafabadi⁷⁴, S. Tizchang^{74,tt}, M. Felcini⁷⁵, M. Grunewald⁷⁵, M. Abbrescia^{76a,76b}, M. Barbieri^{76a,76b}, M. Buonsante^{76a,76b}, A. Colaleo^{76a,76b}, D. Creanza^{76a,76c}, B. D'Anzi^{76a,76b}, N. De Filippis^{76a,76c}, M. De Palma^{76a,76b}, W. Elmetenawee^{76a,76b,uu}, N. Ferrara^{76a,76c}, L. Fiore^{76a}, L. Longo^{76a}, M. Louka^{76a,76b}, G. Maggi^{76a,76c}, M. Maggi^{76a}, I. Margjeka^{76a}, V. Mastrapasqua^{76a,76b}, S. My^{76a,76b}, F. Nenna^{76a,76b}, S. Nuzzo^{76a,76b}, A. Pellecchia^{76a,76b}, A. Pompili^{76a,76b}, G. Pugliese^{76a,76c}, R. Radogna^{76a,76b}

- D. Ramos^{76a}, A. Ranieri^{76a}, L. Silvestris^{76a}, F. M. Simone^{76a,76c}, Ü. Sözbilir^{76a}, A. Stamerra^{76a,76b},
D. Troiano^{76a,76b}, R. Venditti^{76a,76b}, P. Verwilligen^{76a}, A. Zaza^{76a,76b}, G. Abbiendi^{77a}, C. Battilana^{77a,77b},
D. Bonacorsi^{77a,77b}, P. Capiluppi^{77a,77b}, F. R. Cavallo^{77a}, M. Cuffiani^{77a,77b}, G. M. Dallavalle^{77a}, T. Diotallevi^{77a,77b},
F. Fabbri^{77a}, A. Fanfani^{77a,77b}, D. Fasanella^{77a}, P. Giacomelli^{77a}, C. Grandi^{77a}, L. Guiducci^{77a,77b},
S. Lo Meo^{77a,vv}, M. Lorusso^{77a,77b}, L. Lunerti^{77a}, S. Marcellini^{77a}, F. L. Navarria^{77a,77b}, G. Paggi^{77a,77b},
A. Perrotta^{77a}, F. Primavera^{77a,77b}, A. M. Rossi^{77a,77b}, S. Rossi Tisbeni^{77a,77b}, T. Rovelli^{77a,77b}, G. P. Siroli^{77a,77b},
S. Costa^{78a,78b,vv}, A. Di Mattia^{78a}, A. Lapertosa^{78a}, R. Potenza^{78a,78b}, A. Tricomi^{78a,78b,vv}, J. Altork^{79a,79b},
P. Assiouras^{79a}, G. Barbagli^{79a}, G. Bardelli^{79a}, M. Bartolini^{79a,79b}, A. Calandri^{79a,79b}, B. Camaiani^{79a,79b},
A. Cassese^{79a}, R. Ceccarelli^{79a}, V. Ciulli^{79a,79b}, C. Civinini^{79a}, R. D'Alessandro^{79a,79b}, L. Damenti^{79a,79b},
E. Focardi^{79a,79b}, T. Kello^{79a}, G. Latino^{79a,79b}, P. Lenzi^{79a,79b}, M. Lizzo^{79a}, M. Meschini^{79a}, S. Paoletti^{79a},
A. Papanastassiou^{79a,79b}, G. Sguazzoni^{79a}, L. Viliani^{79a}, L. Benussi⁸⁰, S. Colafranceschi⁸⁰, S. Meola^{80,xx},
D. Piccolo⁸⁰, M. Alves Gallo Pereira^{81a}, F. Ferro^{81a}, E. Robutti^{81a}, S. Tosi^{81a,81b}, A. Benaglia^{82a}, F. Brivio^{82a},
V. Camagni^{82a,82b}, F. Cetorelli^{82a,82b}, F. De Guio^{82a,82b}, M. E. Dinardo^{82a,82b}, P. Dini^{82a}, S. Gennai^{82a},
R. Gerosa^{82a,82b}, A. Ghezzi^{82a,82b}, P. Govoni^{82a,82b}, L. Guzzi^{82a}, G. Lavizzari^{82a,82b}, M. T. Lucchini^{82a,82b},
M. Malberti^{82a}, S. Malvezzi^{82a}, A. Massironi^{82a}, D. Menasce^{82a}, L. Moroni^{82a}, M. Paganoni^{82a,82b},
S. Palluotto^{82a,82b}, D. Pedrini^{82a}, A. Perego^{82a,82b}, B. S. Pinolini^{82a}, G. Pizzati^{82a,82b}, S. Ragazzi^{82a,82b},
T. Tabarelli de Fatis^{82a,82b}, S. Buontempo^{83a}, A. Cagnotta^{83a,83b}, C. Di Fraia^{83a,83b}, F. Fabozzi^{83a,83c},
L. Favilla^{83a,83d}, A. O. M. Iorio^{83a,83b}, L. Lista^{83a,83b,yy}, P. Paolucci^{83a,cc}, B. Rossi^{84a}, P. Azzi^{84a},
N. Bacchetta^{84a,zz}, A. Bergnoli^{84a}, P. Bortignon^{84a}, G. Bortolato^{84a,84b}, A. C. M. Bulla^{84a}, R. Carlin^{84a,84b},
P. Checchia^{84a}, T. Dorigo^{84a,aaa}, F. Gasparini^{84a,84b}, S. Giorgetti^{84a}, E. Lusiani^{84a}, M. Margoni^{84a,84b},
A. T. Meneguzzo^{84a,84b}, J. Pazzini^{84a,84b}, P. Ronchese^{84a,84b}, R. Rossin^{84a,84b}, F. Simonetto^{84a,84b}, M. Tosi^{84a,84b},
A. Triossi^{84a,84b}, S. Ventura^{84a}, M. Zanetti^{84a,84b}, P. Zotto^{84a,84b}, A. Zucchetta^{84a,84b}, G. Zumerle^{84a,84b},
A. Braghieri^{85a}, S. Calzaferri^{85a}, P. Montagna^{85a,85b}, M. Pelliccioni^{85a}, V. Re^{85a}, C. Riccardi^{85a,85b}, P. Salvini^{85a},
I. Vai^{85a,85b}, P. Vitulo^{85a,85b}, S. Ajmal^{86a,86b}, M. E. Ascoti^{86a,86b}, G. M. Bilei^{86a}, C. Carrivale^{86a,86b},
D. Ciangottini^{86a,86b}, L. Della Penna^{86a,86b}, L. Fanò^{86a,86b}, V. Mariani^{86a,86b}, M. Menichelli^{86a}, F. Moscatelli^{86a,bbb},
A. Rossi^{86a,86b}, A. Santocchia^{86a,86b}, D. Spiga^{86a}, T. Tedeschi^{86a,86b}, C. Aimè^{87a,87b}, C. A. Alexe^{87a,87c},
P. Asenov^{87a,87b}, P. Azzurri^{87a}, G. Bagliesi^{87a}, R. Bhattacharya^{87a}, L. Bianchini^{87a,87b}, T. Boccali^{87a},
E. Bossini^{87a}, D. Bruschini^{87a,87c}, L. Calligaris^{87a,87b}, R. Castaldi^{87a}, F. Cattafesta^{87a,87c}, M. A. Ciocci^{87a,87b},
M. Cipriani^{87a,87b}, V. D'Amante^{87a,87d}, R. Dell'Orso^{87a}, S. Donato^{87a,87b}, R. Forti^{87a,87b}, A. Giassi^{87a},
F. Ligabue^{87a,87c}, A. C. Marini^{87a,87b}, D. Matos Figueiredo^{87a}, A. Messineo^{87a,87b}, S. Mishra^{87a},
V. K. Muraleedharan Nair Bindhu^{87a,87b}, S. Nandan^{87a}, F. Palla^{87a}, M. Riggirello^{87a,87c}, A. Rizzi^{87a,87b},
G. Rolandi^{87a,87c}, S. Roy Chowdhury^{87a,ccc}, T. Sarkar^{87a}, A. Scribano^{87a}, P. Spagnolo^{87a}, F. Tenchini^{87a,87b},
R. Tenchini^{87a}, G. Tonelli^{87a,87b}, N. Turini^{87a,87d}, F. Vaselli^{87a,87c}, A. Venturi^{87a}, P. G. Verdini^{87a}, P. Akrap^{88a,88b},
C. Basile^{88a,88b}, S. C. Behera^{88a}, F. Cavallari^{88a}, L. Cunqueiro Mendez^{88a,88b}, F. De Ruggi^{88a,88b}, D. Del Re^{88a,88b},
E. Di Marco^{88a}, M. Diemoz^{88a}, F. Errico^{88a}, L. Frosina^{88a,88b}, R. Gargiulo^{88a,88b}, B. Harikrishnan^{88a,88b},
F. Lombardi^{88a,88b}, E. Longo^{88a,88b}, L. Martikainen^{88a,88b}, J. Mijuskovic^{88a,88b}, G. Organtini^{88a,88b}, N. Palmeri^{88a,88b},
R. Paramatti^{88a,88b}, C. Quaranta^{88a,88b}, S. Rahatlou^{88a,88b}, C. Rovelli^{88a}, F. Santanastasio^{88a,88b}, L. Soffi^{88a},
V. Vladimirov^{88a,88b}, N. Amapane^{89a,89b}, R. Arcidiacono^{89a,89c}, S. Argiro^{89a,89b}, M. Arneodo^{89a,89c}, N. Bartosik^{89a,89c},
R. Bellan^{89a,89b}, A. Bellora^{89a,89b}, C. Biino^{89a}, C. Borca^{89a,89b}, N. Cartiglia^{89a}, M. Costa^{89a,89b}, R. Covarelli^{89a,89b},
N. Demaria^{89a}, L. Finco^{89a}, M. Grippo^{89a,89b}, B. Kiani^{89a,89b}, L. Lanteri^{89a,89b}, F. Legger^{89a}, F. Luongo^{89a,89b},
C. Mariotti^{89a}, S. Maselli^{89a}, A. Mecca^{89a,89b}, L. Menzio^{89a,89b}, P. Meridiani^{89a}, E. Migliore^{89a,89b}, M. Monteno^{89a},
M. M. Obertino^{89a,89b}, G. Ortona^{89a}, L. Pacher^{89a,89b}, N. Pastrone^{89a}, M. Rupa^{89a,89c}, F. Siviero^{89a,89b},
V. Sola^{89a,89b}, A. Solano^{89a,89b}, A. Staiano^{89a}, C. Tarricone^{89a,89b}, D. Trocino^{89a}, G. Umoret^{89a,89b},
E. Vlasov^{89a,89b}, R. White^{89a,89b}, J. Babbar^{90a,90b}, S. Belforte^{90a}, V. Candelise^{90a,90b}, M. Casarsa^{90a}, F. Cossutti^{90a},
K. De Leo^{90a}, G. Della Ricca^{90a,90b}, R. Delli Gatti^{90a,90b}, S. Dogra⁹¹, J. Hong⁹¹, J. Kim⁹¹, T. Kim⁹¹, D. Lee⁹¹,
H. Lee⁹¹, J. Lee⁹¹, S. W. Lee⁹¹, C. S. Moon⁹¹, Y. D. Oh⁹¹, S. Sekmen⁹¹, B. Tae⁹¹, Y. C. Yang⁹¹, M. S. Kim⁹²,
G. Bak⁹³, P. Gwak⁹³, H. Kim⁹³, D. H. Moon⁹³, J. Seo⁹³, E. Asilar⁹⁴, F. Carnevali⁹⁴, J. Choi^{94,ddd}, T. J. Kim⁹⁴,
Y. Ryou⁹⁴, S. Ha⁹⁵, S. Han⁹⁵, B. Hong⁹⁵, K. Lee⁹⁵, K. S. Lee⁹⁵, S. Lee⁹⁵, J. Yoo⁹⁵, J. Goh⁹⁶, J. Shin⁹⁶,
S. Yang⁹⁶, Y. Kang⁹⁷, H. S. Kim⁹⁷, Y. Kim⁹⁷, S. Lee⁹⁷, J. Almond⁹⁸, J. H. Bhyun⁹⁸, J. Choi⁹⁸, J. Choi⁹⁸, W. Jun⁹⁸

J. Kim⁹⁸, T. Kim⁹⁸, Y. Kim⁹⁸, Y. W. Kim⁹⁸, S. Ko⁹⁸, H. Lee⁹⁸, J. Lee⁹⁸, J. Lee⁹⁸, B. H. Oh⁹⁸, S. B. Oh⁹⁸,
H. Seo⁹⁸, J. Shin⁹⁸, U. K. Yang⁹⁸, I. Yoon⁹⁸, W. Jang⁹⁹, D. Y. Kang⁹⁹, D. Kim⁹⁹, S. Kim⁹⁹, B. Ko⁹⁹,
J. S. H. Lee⁹⁹, Y. Lee⁹⁹, J. A. Merlin⁹⁹, I. C. Park⁹⁹, Y. Roh⁹⁹, I. J. Watson⁹⁹, G. Cho¹⁰⁰, K. Hwang¹⁰⁰, B. Kim¹⁰⁰,
S. Kim¹⁰⁰, K. Lee¹⁰⁰, H. D. Yoo¹⁰⁰, M. Choi¹⁰¹, M. R. Kim¹⁰¹, Y. Lee¹⁰¹, I. Yu¹⁰¹, T. Beyrouthy¹⁰²,
Y. Gharbia¹⁰², F. Alazemi¹⁰³, K. Dreimanis¹⁰⁴, O. M. Eberlins¹⁰⁴, A. Gaile¹⁰⁴, C. Munoz Diaz¹⁰⁴, D. Osite¹⁰⁴,
G. Pikurs¹⁰⁴, R. Plese¹⁰⁴, A. Potrebko¹⁰⁴, M. Seidel¹⁰⁴, D. Sidiropoulos Kontos¹⁰⁴, N. R. Strautnieks¹⁰⁵,
M. Ambrozias¹⁰⁶, A. Juodagalvis¹⁰⁶, S. Nargelas¹⁰⁶, A. Rinkevicius¹⁰⁶, G. Tamulaitis¹⁰⁶, I. Yusuff^{107,eee},
Z. Zolkapli¹⁰⁷, J. F. Benitez¹⁰⁸, A. Castaneda Hernandez¹⁰⁸, A. Cota Rodriguez¹⁰⁸, L. E. Cuevas Picos¹⁰⁸,
H. A. Encinas Acosta¹⁰⁸, L. G. Gallegos Maríñez¹⁰⁸, M. León Coello¹⁰⁸, J. A. Murillo Quijada¹⁰⁸, A. Sehwat¹⁰⁸,
L. Valencia Palomo¹⁰⁸, G. Ayala¹⁰⁹, H. Castilla-Valdez¹⁰⁹, H. Crotte Ledesma¹⁰⁹, R. Lopez-Fernandez¹⁰⁹,
J. Mejia Guisao¹⁰⁹, R. Reyes-Almanza¹⁰⁹, A. Sánchez Hernández¹⁰⁹, C. Oropeza Barrera¹¹⁰,
D. L. Ramirez Guadarrama¹¹⁰, M. Ramírez García¹¹⁰, I. Bautista¹¹¹, F. E. Neri Huerta¹¹¹, I. Pedraza¹¹¹,
H. A. Salazar Ibarguen¹¹¹, C. Uribe Estrada¹¹¹, I. Bujanja¹¹², N. Raicevic¹¹², P. H. Butler¹¹³, A. Ahmad¹¹⁴,
M. I. Asghar¹¹⁴, A. Awais¹¹⁴, M. I. M. Awan¹¹⁴, W. A. Khan¹¹⁴, V. Avati¹¹⁵, L. Forthomme¹¹⁵, L. Grzanka¹¹⁵,
M. Malawski¹¹⁵, K. Piotrkowski¹¹⁵, M. Bluj¹¹⁶, M. Górski¹¹⁶, M. Kazana¹¹⁶, M. Szleper¹¹⁶, P. Zalewski¹¹⁶,
K. Bunkowski¹¹⁷, K. Doroba¹¹⁷, A. Kalinowski¹¹⁷, M. Konecki¹¹⁷, J. Krolikowski¹¹⁷, A. Muhammad¹¹⁷,
P. Fokow¹¹⁸, K. Pozniak¹¹⁸, W. Zabolotny¹¹⁸, M. Araujo¹¹⁹, D. Bastos¹¹⁹, C. Beirão Da Cruz E Silva¹¹⁹,
A. Boletti¹¹⁹, M. Bozzo¹¹⁹, T. Camporesi¹¹⁹, G. Da Molin¹¹⁹, P. Faccioli¹¹⁹, M. Gallinaro¹¹⁹, J. Hollar¹¹⁹,
N. Leonardo¹¹⁹, G. B. Marozzo¹¹⁹, A. Petrilli¹¹⁹, M. Pisano¹¹⁹, J. Seixas¹¹⁹, J. Varela¹¹⁹, J. W. Wulff¹¹⁹,
P. Adzic¹²⁰, L. Markovic¹²⁰, P. Milenovic¹²⁰, V. Milosevic¹²⁰, D. Devetak¹²¹, M. Dordevic¹²¹, J. Milosevic¹²¹,
L. Nadder¹²¹, V. Rekovic¹²¹, M. Stojanovic¹²¹, M. Alcalde Martinez¹²², J. Alcaraz Maestre¹²²,
Cristina F. Bedoya¹²², J. A. Brochero Cifuentes¹²², Oliver M. Carretero¹²², M. Cepeda¹²², M. Cerrada¹²²,
N. Colino¹²², J. Cuchillo Ortega¹²², B. De La Cruz¹²², A. Delgado Peris¹²², A. Escalante Del Valle¹²²,
D. Fernández Del Val¹²², J. P. Fernández Ramos¹²², J. Flix¹²², M. C. Fouz¹²², M. Gonzalez Hernandez¹²²,
O. Gonzalez Lopez¹²², S. Goy Lopez¹²², J. M. Hernandez¹²², M. I. Josa¹²², J. Llorente Merino¹²²,
C. Martin Perez¹²², E. Martin Viscasillas¹²², D. Moran¹²², C. M. Morcillo Perez¹²², R. Paz Herrera¹²²,
C. Perez Dengra¹²², A. Pérez-Calero Yzquierdo¹²², J. Puerta Pelayo¹²², I. Redondo¹²², J. Vazquez Escobar¹²²,
J. F. de Trocóniz¹²³, B. Alvarez Gonzalez¹²⁴, J. Ayllon Torresano¹²⁴, A. Cardini¹²⁴, J. Cuevas¹²⁴,
J. Del Riego Badas¹²⁴, D. Estrada Acevedo¹²⁴, J. Fernandez Menendez¹²⁴, S. Folgueras¹²⁴,
I. Gonzalez Caballero¹²⁴, P. Leguina¹²⁴, M. Obeso Menendez¹²⁴, E. Palencia Cortezon¹²⁴, J. Prado Pico¹²⁴,
A. Soto Rodríguez¹²⁴, C. Vico Villalba¹²⁴, P. Vischia¹²⁴, S. Blanco Fernández¹²⁵, I. J. Cabrillo¹²⁵, A. Calderon¹²⁵,
J. Duarte Campderros¹²⁵, M. Fernandez¹²⁵, G. Gomez¹²⁵, C. Lasasa García¹²⁵, R. Lopez Ruiz¹²⁵,
C. Martinez Rivero¹²⁵, P. Martinez Ruiz del Arbol¹²⁵, F. Matorras¹²⁵, P. Matorras Cuevas¹²⁵,
E. Navarrete Ramos¹²⁵, J. Piedra Gomez¹²⁵, C. Quintana San Emeterio¹²⁵, L. Scodellaro¹²⁵, I. Vila¹²⁵,
R. Vilar Cortabitarte¹²⁵, J. M. Vizán García¹²⁵, B. Kailasapathy^{126,fff}, D. D. C. Wickramaratna¹²⁶,
W. G. D. Dharmaratna^{127,ggg}, K. Liyanage¹²⁷, N. Perera¹²⁷, D. Abbaneo¹²⁸, C. Amendola¹²⁸, R. Ardino¹²⁸,
E. Auffray¹²⁸, J. Baechler¹²⁸, D. Barney¹²⁸, M. Bianco¹²⁸, A. Bocci¹²⁸, L. Borgonovi¹²⁸, C. Botta¹²⁸,
A. Bragagnolo¹²⁸, C. E. Brown¹²⁸, C. Caillol¹²⁸, G. Cerminara¹²⁸, P. Connor¹²⁸, D. d'Enterria¹²⁸,
A. Dabrowski¹²⁸, A. David¹²⁸, A. De Roeck¹²⁸, M. M. DeFranchis¹²⁸, M. Deile¹²⁸, M. Dobson¹²⁸, W. Funk¹²⁸,
A. Gaddi¹²⁸, S. Giani¹²⁸, D. Gigi¹²⁸, K. Gill¹²⁸, F. Glege¹²⁸, M. Glowacki¹²⁸, A. Gruber¹²⁸, J. Hegeman¹²⁸,
J. K. Heikkilä¹²⁸, B. Huber¹²⁸, V. Innocente¹²⁸, T. James¹²⁸, P. Janot¹²⁸, O. Kaluzinska¹²⁸, O. Karacheban^{128,aa},
G. Karathanasis¹²⁸, S. Laurila¹²⁸, P. Lecoq¹²⁸, C. Lourenço¹²⁸, A.-M. Lyon¹²⁸, M. Magherini¹²⁸, L. Malgeri¹²⁸,
M. Mannelli¹²⁸, M. Matthewman¹²⁸, A. Mehta¹²⁸, F. Meijers¹²⁸, S. Mersi¹²⁸, E. Meschi¹²⁸, M. Migliorini¹²⁸,
F. Monti¹²⁸, F. Moortgat¹²⁸, M. Mulders¹²⁸, M. Musich¹²⁸, I. Neutelings¹²⁸, S. Orfanelli¹²⁸, F. Pantaleo¹²⁸,
M. Pari¹²⁸, G. Petrucciani¹²⁸, A. Pfeiffer¹²⁸, M. Pierini¹²⁸, M. Pitt¹²⁸, H. Qu¹²⁸, D. Rabady¹²⁸,
B. Ribeiro Lopes¹²⁸, F. Riti¹²⁸, P. Rosado¹²⁸, M. Rovere¹²⁸, H. Sakulin¹²⁸, R. Salvatico¹²⁸, S. Sanchez Cruz¹²⁸,
S. Scarfi¹²⁸, M. Selvaggi¹²⁸, A. Sharma¹²⁸, K. Shchelina¹²⁸, P. Silva¹²⁸, P. Sphicas^{128,hhh}, A. G. Stahl Leitner¹²⁸,
A. Steen¹²⁸, S. Summers¹²⁸, D. Treille¹²⁸, P. Tropea¹²⁸, E. Vernazza¹²⁸, J. Wanczyk^{128,iii}, J. Wang¹²⁸,
S. Wuchterl¹²⁸, M. Zarucki¹²⁸, P. Zehetner¹²⁸, P. Zejdl¹²⁸, G. Zevi Della Porta¹²⁸, T. Bevilacqua^{129,jjj}

- L. Caminada^{129,jjj} W. Erdmann¹²⁹ R. Horisberger¹²⁹ Q. Ingram¹²⁹ H. C. Kaestli¹²⁹ D. Kotlinski¹²⁹
 C. Lange¹²⁹ U. Langenegger¹²⁹ M. Missiroli^{129,jjj} L. Noehte^{129,jjj} T. Rohe¹²⁹ A. Samalan¹²⁹
 T. K. Aarrestad¹³⁰ M. Backhaus¹³⁰ G. Bonomelli¹³⁰ C. Cazzaniga¹³⁰ K. Datta¹³⁰
 P. De Bryas Dexmiers D'archiacchiac^{130,iii} A. De Cosa¹³⁰ G. Dissertori¹³⁰ M. Dittmar¹³⁰ M. Donegà¹³⁰
 F. Eble¹³⁰ K. Gedia¹³⁰ F. Glessgen¹³⁰ C. Grab¹³⁰ N. Härringer¹³⁰ T. G. Harte¹³⁰ W. Lustermann¹³⁰
 M. Malucchi¹³⁰ R. A. Manzoni¹³⁰ M. Marchegiani¹³⁰ L. Marchese¹³⁰ A. Mascellani^{130,iii} F. Nessi-Tedaldi¹³⁰
 F. Pauss¹³⁰ V. Perovic¹³⁰ B. Ristic¹³⁰ R. Seidita¹³⁰ J. Steggemann^{130,iii} A. Tarabini¹³⁰ D. Valsecchi¹³⁰
 R. Wallny¹³⁰ C. Amsler^{131,kkk} P. Bärtschi¹³¹ F. Bilandzija¹³¹ M. F. Canelli¹³¹ G. Celotto¹³¹ K. Cormier¹³¹
 M. Huwiler¹³¹ W. Jin¹³¹ A. Jofrehei¹³¹ B. Kilminster¹³¹ T. H. Kwok¹³¹ S. Leontsinis¹³¹ V. Lukashenko¹³¹
 A. Macchiolo¹³¹ F. Meng¹³¹ J. Motta¹³¹ A. Reimers¹³¹ P. Robmann¹³¹ M. Senger¹³¹ E. Shokr¹³¹
 F. Stäger¹³¹ R. Tramontano¹³¹ D. Bhowmik¹³² C. M. Kuo¹³² P. K. Rout¹³² S. Taj¹³² P. C. Tiwari^{132,II}
 L. Ceard¹³³ K. F. Chen¹³³ Z. g. Chen¹³³ A. De Iorio¹³³ W.-S. Hou¹³³ T. h. Hsu¹³³ Y. w. Kao¹³³ S. Karmakar¹³³
 G. Kole¹³³ Y. y. Li¹³³ R.-S. Lu¹³³ E. Paganis¹³³ X. f. Su¹³³ J. Thomas-Wilsker¹³³ L. s. Tsai¹³³ D. Tsiouou¹³³
 H. y. Wu¹³³ E. Yazgan¹³³ C. Asawatangtrakuldee¹³⁴ N. Srimanobhas¹³⁴ Y. Maghrbi¹³⁵ D. Agyel¹³⁶
 F. Boran¹³⁶ F. Dolek¹³⁶ I. Dumanoglu^{136,III} Y. Guler^{136,mmm} E. Gurpinar Guler^{136,mmm} C. Isik¹³⁶ O. Kara¹³⁶
 A. Kayis Topaksu¹³⁶ Y. Komurcu¹³⁶ G. Onengut¹³⁶ K. Ozdemir^{136,nnn} B. Tali^{136,ooo} U. G. Tok¹³⁶ E. Uslan¹³⁶
 I. S. Zorbakir¹³⁶ M. Yalvac^{137,ppp} B. Akgun¹³⁸ I. O. Atakisi^{138,qqq} E. Gülmez¹³⁸ M. Kaya^{138,rrr} O. Kaya^{138,sss}
 M. A. Sarkisla^{138,ttt} S. Tekten^{138,uuu} A. Cakir¹³⁹ K. Cankocak^{139,III,vvv} S. Sen^{139,www} O. Aydilek^{140,xxx}
 B. Hacisahinoglu¹⁴⁰ I. Hos^{140,yyy} B. Kaynak¹⁴⁰ S. Ozkorucuklu¹⁴⁰ O. Potok¹⁴⁰ H. Sert¹⁴⁰ C. Simsek¹⁴⁰
 C. Zorbilmez¹⁴⁰ S. Cerci¹⁴¹ A. A. Guvenli¹⁴¹ B. Isildak^{141,zzz} D. Sunar Cerci¹⁴¹ T. Yetkin^{141,v}
 A. Boyaryntsev¹⁴² O. Dadazhanova¹⁴² B. Grynyov¹⁴² L. Levchuk¹⁴³ J. J. Brooke¹⁴⁴ A. Bundock¹⁴⁴
 F. Bury¹⁴⁴ E. Clement¹⁴⁴ D. Cussans¹⁴⁴ D. Dharmender¹⁴⁴ H. Flacher¹⁴⁴ J. Goldstein¹⁴⁴ H. F. Heath¹⁴⁴
 M.-L. Holmberg¹⁴⁴ L. Kreczko¹⁴⁴ S. Paramesvaran¹⁴⁴ L. Robertshaw¹⁴⁴ M. S. Sanjrani¹⁴⁴ J. Segal¹⁴⁴
 V. J. Smith¹⁴⁴ A. H. Ball¹⁴⁵ K. W. Bell¹⁴⁵ A. Belyaev^{145,aaaa} C. Brew¹⁴⁵ R. M. Brown¹⁴⁵ D. J. A. Cockerill¹⁴⁵
 C. Cooke¹⁴⁵ A. Elliot¹⁴⁵ K. V. Ellis¹⁴⁵ J. Gajownik¹⁴⁵ K. Harder¹⁴⁵ S. Harper¹⁴⁵ J. Linacre¹⁴⁵
 K. Manolopoulos¹⁴⁵ M. Moallemi¹⁴⁵ D. M. Newbold¹⁴⁵ E. Olaiya¹⁴⁵ D. Petyt¹⁴⁵ T. Reis¹⁴⁵
 A. R. Sahasransu¹⁴⁵ G. Salvi¹⁴⁵ T. Schuh¹⁴⁵ C. H. Shepherd-Themistocleous¹⁴⁵ I. R. Tomalin¹⁴⁵
 K. C. Whalen¹⁴⁵ T. Williams¹⁴⁵ I. Andreou¹⁴⁶ R. Bainbridge¹⁴⁶ P. Bloch¹⁴⁶ O. Buchmuller¹⁴⁶
 C. A. Carrillo Montoya¹⁴⁶ D. Colling¹⁴⁶ J. S. Dancu¹⁴⁶ I. Das¹⁴⁶ P. Dauncey¹⁴⁶ G. Davies¹⁴⁶
 M. Della Negra¹⁴⁶ S. Fayer¹⁴⁶ G. Fedi¹⁴⁶ G. Hall¹⁴⁶ H. R. Hoorani¹⁴⁶ A. Howard¹⁴⁶ G. Iles¹⁴⁶ C. R. Knight¹⁴⁶
 P. Krueper¹⁴⁶ J. Langford¹⁴⁶ K. H. Law¹⁴⁶ J. León Holgado¹⁴⁶ E. Leutgeb¹⁴⁶ L. Lyons¹⁴⁶ A.-M. Magnan¹⁴⁶
 B. Maier¹⁴⁶ S. Mallios¹⁴⁶ A. Mastronikolis¹⁴⁶ M. Mieskolainen¹⁴⁶ J. Nash^{146,bbbb} M. Pesaresi¹⁴⁶
 P. B. Pradeep¹⁴⁶ B. C. Radburn-Smith¹⁴⁶ A. Richards¹⁴⁶ A. Rose¹⁴⁶ L. Russell¹⁴⁶ K. Savva¹⁴⁶ C. Seez¹⁴⁶
 R. Shukla¹⁴⁶ A. Tapper¹⁴⁶ K. Uchida¹⁴⁶ G. P. Uttley¹⁴⁶ T. Virdee^{146,cc} M. Vojinovic¹⁴⁶ N. Wardle¹⁴⁶
 D. Winterbottom¹⁴⁶ J. E. Cole¹⁴⁷ A. Khan¹⁴⁷ P. Kyberd¹⁴⁷ I. D. Reid¹⁴⁷ S. Abdullin¹⁴⁸ A. Brinkerhoff¹⁴⁸
 E. Collins¹⁴⁸ M. R. Darwish¹⁴⁸ J. Dittmann¹⁴⁸ K. Hatakeyama¹⁴⁸ V. Hegde¹⁴⁸ J. Hiltbrand¹⁴⁸
 B. McMaster¹⁴⁸ J. Samudio¹⁴⁸ S. Sawant¹⁴⁸ C. Sutantawibul¹⁴⁸ J. Wilson¹⁴⁸ J. M. Hogan^{149,cccc}
 R. Bartek¹⁵⁰ A. Dominguez¹⁵⁰ S. Raj¹⁵⁰ A. E. Simsek¹⁵⁰ S. S. Yu¹⁵⁰ B. Bam¹⁵¹ A. Buchot Perraguin¹⁵¹
 S. Campbell¹⁵¹ R. Chudasama¹⁵¹ S. I. Cooper¹⁵¹ C. Crovella¹⁵¹ G. Fidalgo¹⁵¹ S. V. Gleyzer¹⁵¹
 A. Khukhunaishvili¹⁵¹ K. Matchev¹⁵¹ E. Pearson¹⁵¹ C. U. Perez¹⁵¹ P. Rumerio^{151,dddd} E. Usai¹⁵¹ R. Yi¹⁵¹
 S. Cholak¹⁵² G. De Castro¹⁵² Z. Demiragli¹⁵² C. Erice¹⁵² C. Fangmeier¹⁵² C. Fernandez Madrazo¹⁵²
 E. Fontanesi¹⁵² J. Fulcher¹⁵² F. Golf¹⁵² S. Jeon¹⁵² J. O'Cain¹⁵² I. Reed¹⁵² J. Rohlf¹⁵² K. Salyer¹⁵²
 D. Sperka¹⁵² D. Spitzbart¹⁵² I. Suarez¹⁵² A. Tsatsos¹⁵² E. Wurtz¹⁵² A. G. Zecchinelli¹⁵² G. Barone¹⁵³
 G. Benelli¹⁵³ D. Cutts¹⁵³ S. Ellis¹⁵³ L. Gouskos¹⁵³ M. Hadley¹⁵³ U. Heintz¹⁵³ K. W. Ho¹⁵³ T. Kwon¹⁵³
 G. Landsberg¹⁵³ K. T. Lau¹⁵³ J. Luo¹⁵³ S. Mondal¹⁵³ J. Roloff¹⁵³ T. Russell¹⁵³ S. Sagir^{153,eeee} X. Shen¹⁵³
 M. Stamenkovic¹⁵³ N. Venkatasubramanian¹⁵³ S. Abbott¹⁵⁴ B. Barton¹⁵⁴ R. Breedon¹⁵⁴ H. Cai¹⁵⁴
 M. Calderon De La Barca Sanchez¹⁵⁴ M. Chertok¹⁵⁴ M. Citron¹⁵⁴ J. Conway¹⁵⁴ P. T. Cox¹⁵⁴ R. Erbacher¹⁵⁴
 O. Kukral¹⁵⁴ G. Mocellin¹⁵⁴ S. Ostrom¹⁵⁴ W. Wei¹⁵⁴ S. Yoo¹⁵⁴ K. Adamidis¹⁵⁵ M. Bachtis¹⁵⁵ D. Campos¹⁵⁵
 R. Cousins¹⁵⁵ A. Datta¹⁵⁵ G. Flores Avila¹⁵⁵ J. Hauser¹⁵⁵ M. Ignatenko¹⁵⁵ M. A. Iqbal¹⁵⁵ T. Lam¹⁵⁵

Y. f. Lo¹⁵⁵, E. Manca¹⁵⁵, A. Nunez Del Prado¹⁵⁵, D. Saltzberg¹⁵⁵, V. Valuev¹⁵⁵, R. Clare¹⁵⁶, J. W. Gary¹⁵⁶, G. Hanson¹⁵⁶, A. Aportela¹⁵⁷, A. Arora¹⁵⁷, J. G. Branson¹⁵⁷, S. Cittolin¹⁵⁷, S. Cooperstein¹⁵⁷, D. Diaz¹⁵⁷, J. Duarte¹⁵⁷, L. Giannini¹⁵⁷, Y. Gu¹⁵⁷, J. Guiang¹⁵⁷, V. Krutelyov¹⁵⁷, R. Lee¹⁵⁷, J. Letts¹⁵⁷, H. Li¹⁵⁷, M. Masciovecchio¹⁵⁷, F. Mokhtar¹⁵⁷, S. Mukherjee¹⁵⁷, M. Pieri¹⁵⁷, D. Primosch¹⁵⁷, M. Quinnan¹⁵⁷, V. Sharma¹⁵⁷, M. Tadel¹⁵⁷, E. Vourliotis¹⁵⁷, F. Würthwein¹⁵⁷, A. Yagil¹⁵⁷, Z. Zhao¹⁵⁷, A. Barzdukas¹⁵⁸, L. Brennan¹⁵⁸, C. Campagnari¹⁵⁸, S. Carron Montero^{158,ffff}, K. Downham¹⁵⁸, C. Grieco¹⁵⁸, M. M. Hussain¹⁵⁸, J. Incandela¹⁵⁸, J. Kim¹⁵⁸, M. W. K. Lai¹⁵⁸, A. J. Li¹⁵⁸, P. Masterson¹⁵⁸, J. Richman¹⁵⁸, S. N. Santpur¹⁵⁸, U. Sarica¹⁵⁸, R. Schmitz¹⁵⁸, F. Setti¹⁵⁸, J. Sheplock¹⁵⁸, D. Stuart¹⁵⁸, T. Á. Vami¹⁵⁸, X. Yan¹⁵⁸, D. Zhang¹⁵⁸, A. Albert¹⁵⁹, S. Bhattacharya¹⁵⁹, A. Bornheim¹⁵⁹, O. Cerri¹⁵⁹, R. Kansal¹⁵⁹, J. Mao¹⁵⁹, H. B. Newman¹⁵⁹, G. Reales Gutiérrez¹⁵⁹, T. Sievert¹⁵⁹, M. Spiropulu¹⁵⁹, J. R. Vlimant¹⁵⁹, R. A. Wynne¹⁵⁹, S. Xie¹⁵⁹, J. Alison¹⁶⁰, S. An¹⁶⁰, M. Cremonesi¹⁶⁰, V. Dutta¹⁶⁰, E. Y. Ertorer¹⁶⁰, T. Ferguson¹⁶⁰, T. A. Gómez Espinosa¹⁶⁰, A. Harilal¹⁶⁰, A. Kallil Tharayil¹⁶⁰, M. Kanemura¹⁶⁰, C. Liu¹⁶⁰, P. Meiring¹⁶⁰, T. Mudholkar¹⁶⁰, S. Murthy¹⁶⁰, P. Palit¹⁶⁰, K. Park¹⁶⁰, M. Paulini¹⁶⁰, A. Roberts¹⁶⁰, A. Sanchez¹⁶⁰, W. Terrill¹⁶⁰, J. P. Cumalat¹⁶¹, W. T. Ford¹⁶¹, A. Hart¹⁶¹, A. Hassani¹⁶¹, S. Kwan¹⁶¹, J. Pearkes¹⁶¹, C. Savard¹⁶¹, N. Schonbeck¹⁶¹, K. Stenson¹⁶¹, K. A. Ulmer¹⁶¹, S. R. Wagner¹⁶¹, N. Zipper¹⁶¹, D. Zuolo¹⁶¹, J. Alexander¹⁶², X. Chen¹⁶², D. J. Cranshaw¹⁶², J. Dickinson¹⁶², J. Fan¹⁶², X. Fan¹⁶², J. Grassi¹⁶², S. Hogan¹⁶², P. Kotamnives¹⁶², J. Monroy¹⁶², G. Niendorf¹⁶², M. Oshiro¹⁶², J. R. Patterson¹⁶², M. Reid¹⁶², A. Ryd¹⁶², J. Thom¹⁶², P. Wittich¹⁶², R. Zou¹⁶², L. Zygala¹⁶², M. Albrow¹⁶³, M. Alyari¹⁶³, O. Amram¹⁶³, G. Apollinari¹⁶³, A. Apresyan¹⁶³, L. A. T. Bauerdick¹⁶³, D. Berry¹⁶³, J. Berryhill¹⁶³, P. C. Bhat¹⁶³, K. Burkett¹⁶³, J. N. Butler¹⁶³, A. Canepa¹⁶³, G. B. Cerati¹⁶³, H. W. K. Cheung¹⁶³, F. Chlebana¹⁶³, C. Cosby¹⁶³, G. Cummings¹⁶³, I. Dutta¹⁶³, V. D. Elvira¹⁶³, J. Freeman¹⁶³, A. Gandrakota¹⁶³, Z. Gecse¹⁶³, L. Gray¹⁶³, D. Green¹⁶³, A. Grummer¹⁶³, S. Grünendahl¹⁶³, D. Guerrero¹⁶³, O. Gutsche¹⁶³, R. M. Harris¹⁶³, T. C. Herwig¹⁶³, J. Hirschauer¹⁶³, B. Jayatilaka¹⁶³, S. Jindariani¹⁶³, M. Johnson¹⁶³, U. Joshi¹⁶³, T. Klijnsma¹⁶³, B. Klima¹⁶³, K. H. M. Kwok¹⁶³, S. Lammel¹⁶³, C. Lee¹⁶³, D. Lincoln¹⁶³, R. Lipton¹⁶³, T. Liu¹⁶³, K. Maeshima¹⁶³, D. Mason¹⁶³, P. McBride¹⁶³, P. Merkel¹⁶³, S. Mrenna¹⁶³, S. Nahn¹⁶³, J. Ngadiuba¹⁶³, D. Noonan¹⁶³, S. Norberg¹⁶³, V. Papadimitriou¹⁶³, N. Pastika¹⁶³, K. Pedro¹⁶³, C. Pena^{163,gggg}, C. E. Perez Lara¹⁶³, F. Ravera¹⁶³, A. Reinsvold Hall^{163,hhhh}, L. Ristori¹⁶³, M. Safdari¹⁶³, E. Sexton-Kennedy¹⁶³, N. Smith¹⁶³, A. Soha¹⁶³, L. Spiegel¹⁶³, S. Stoynev¹⁶³, J. Strait¹⁶³, L. Taylor¹⁶³, S. Tkaczyk¹⁶³, N. V. Tran¹⁶³, L. Uplegger¹⁶³, E. W. Vaandering¹⁶³, C. Wang¹⁶³, I. Zoi¹⁶³, C. Aruta¹⁶⁴, P. Avery¹⁶⁴, D. Bourilkov¹⁶⁴, P. Chang¹⁶⁴, V. Cherepanov¹⁶⁴, R. D. Field¹⁶⁴, C. Huh¹⁶⁴, E. Koenig¹⁶⁴, M. Kolosova¹⁶⁴, J. Konigsberg¹⁶⁴, A. Korytov¹⁶⁴, N. Menendez¹⁶⁴, G. Mitselmakher¹⁶⁴, K. Mohrman¹⁶⁴, A. Muthirakalayil Madhu¹⁶⁴, N. Rawal¹⁶⁴, S. Rosenzweig¹⁶⁴, V. Sulimov¹⁶⁴, Y. Takahashi¹⁶⁴, J. Wang¹⁶⁴, T. Adams¹⁶⁵, A. Al Kadhimi¹⁶⁵, A. Askew¹⁶⁵, S. Bower¹⁶⁵, R. Hashmi¹⁶⁵, R. S. Kim¹⁶⁵, T. Kolberg¹⁶⁵, G. Martinez¹⁶⁵, M. Mazza¹⁶⁵, H. Prosper¹⁶⁵, P. R. Prova¹⁶⁵, M. Wulansatiti¹⁶⁵, R. Yohay¹⁶⁵, B. Alsufyani¹⁶⁶, S. Butalla¹⁶⁶, S. Das¹⁶⁶, M. Hohlmann¹⁶⁶, M. Lavinsky¹⁶⁶, E. Yanes¹⁶⁶, M. R. Adams¹⁶⁷, N. Barnett¹⁶⁷, A. Baty¹⁶⁷, C. Bennett¹⁶⁷, R. Cavanaugh¹⁶⁷, R. Escobar Franco¹⁶⁷, O. Evdokimov¹⁶⁷, C. E. Gerber¹⁶⁷, H. Gupta¹⁶⁷, M. Hawksworth¹⁶⁷, A. Hingrajiya¹⁶⁷, D. J. Hofman¹⁶⁷, J. h. Lee¹⁶⁷, D. S. Lemos¹⁶⁷, C. Mills¹⁶⁷, S. Nanda¹⁶⁷, G. Nigmatkulov¹⁶⁷, B. Ozek¹⁶⁷, T. Phan¹⁶⁷, D. Pilipovic¹⁶⁷, R. Pradhan¹⁶⁷, E. Prifti¹⁶⁷, P. Roy¹⁶⁷, T. Roy¹⁶⁷, N. Singh¹⁶⁷, M. B. Tonjes¹⁶⁷, N. Varelas¹⁶⁷, M. A. Wadud¹⁶⁷, J. Yoo¹⁶⁷, M. Alhusseini¹⁶⁸, D. Blend¹⁶⁸, K. Dilsiz^{168,iiii}, O. K. Köseyan¹⁶⁸, A. Mestvirishvili^{168,jjjj}, O. Neogi¹⁶⁸, H. Ogul^{168,kkkk}, Y. Onel¹⁶⁸, A. Penzo¹⁶⁸, C. Snyder¹⁶⁸, E. Tiras^{168,llll}, B. Blumenfeld¹⁶⁹, J. Davis¹⁶⁹, A. V. Gritsan¹⁶⁹, L. Kang¹⁶⁹, S. Kyriacou¹⁶⁹, P. Maksimovic¹⁶⁹, M. Roguljic¹⁶⁹, S. Sekhar¹⁶⁹, M. V. Srivastav¹⁶⁹, M. Swartz¹⁶⁹, A. Abreu¹⁷⁰, L. F. Alcerro Alcerro¹⁷⁰, J. Anguiano¹⁷⁰, S. Arteaga Escatel¹⁷⁰, P. Baringer¹⁷⁰, A. Bean¹⁷⁰, Z. Flowers¹⁷⁰, D. Grove¹⁷⁰, J. King¹⁷⁰, G. Krintiras¹⁷⁰, M. Lazarovits¹⁷⁰, C. Le Mahieu¹⁷⁰, J. Marquez¹⁷⁰, M. Murray¹⁷⁰, M. Nickel¹⁷⁰, S. Popescu^{170,mmmm}, C. Rogan¹⁷⁰, C. Royon¹⁷⁰, S. Rudrabhatla¹⁷⁰, S. Sanders¹⁷⁰, C. Smith¹⁷⁰, G. Wilson¹⁷⁰, B. Allmond¹⁷¹, R. Gujju Gurunadha¹⁷¹, N. Islam¹⁷¹, A. Ivanov¹⁷¹, K. Kaadze¹⁷¹, Y. Maravin¹⁷¹, J. Natoli¹⁷¹, D. Roy¹⁷¹, G. Sorrentino¹⁷¹, A. Baden¹⁷², A. Belloni¹⁷², J. Bistany-riebman¹⁷², S. C. Eno¹⁷², N. J. Hadley¹⁷², S. Jabeen¹⁷², R. G. Kellogg¹⁷², T. Koeth¹⁷², B. Kronheim¹⁷², S. Lascio¹⁷², P. Major¹⁷², A. C. Mignerey¹⁷², C. Palmer¹⁷², C. Papageorgakis¹⁷², M. M. Paranjpe¹⁷², E. Popova^{172,nnnn}, A. Shevelev¹⁷², L. Zhang¹⁷², C. Baldenegro Barrera¹⁷³, J. Bendavid¹⁷³, H. Bossi¹⁷³, S. Bright-Thonney¹⁷³, I. A. Cali¹⁷³

Y. c. Chen¹⁷³, P. c. Chou¹⁷³, M. D'Alfonso¹⁷³, J. Eysermans¹⁷³, C. Freer¹⁷³, G. Gomez-Ceballos¹⁷³,
M. Goncharov¹⁷³, G. Grosso¹⁷³, P. Harris¹⁷³, D. Hoang¹⁷³, G. M. Innocenti¹⁷³, D. Kovalskyi¹⁷³, J. Krupa¹⁷³,
L. Lavezzo¹⁷³, Y.-J. Lee¹⁷³, K. Long¹⁷³, C. McGinn¹⁷³, A. Novak¹⁷³, M. I. Park¹⁷³, C. Paus¹⁷³, C. Reissel¹⁷³,
C. Roland¹⁷³, G. Roland¹⁷³, S. Rothman¹⁷³, T. a. Sheng¹⁷³, G. S. F. Stephans¹⁷³, D. Walter¹⁷³, Z. Wang¹⁷³,
B. Wyslouch¹⁷³, T. J. Yang¹⁷³, B. Crossman¹⁷⁴, W. J. Jackson¹⁷⁴, C. Kapsiak¹⁷⁴, M. Krohn¹⁷⁴, D. Mahon¹⁷⁴,
J. Mans¹⁷⁴, B. Marzocchi¹⁷⁴, R. Rusack¹⁷⁴, O. Sancar¹⁷⁴, R. Saradhy¹⁷⁴, N. Strobbe¹⁷⁴, K. Bloom¹⁷⁵,
D. R. Claes¹⁷⁵, G. Haza¹⁷⁵, J. Hossain¹⁷⁵, C. Joo¹⁷⁵, I. Kravchenko¹⁷⁵, A. Rohilla¹⁷⁵, J. E. Siado¹⁷⁵,
W. Tabb¹⁷⁵, A. Vagnerini¹⁷⁵, A. Wightman¹⁷⁵, F. Yan¹⁷⁵, H. Bandyopadhyay¹⁷⁶, L. Hay¹⁷⁶, H. w. Hsia¹⁷⁶,
I. Iashvili¹⁷⁶, A. Kalogeropoulos¹⁷⁶, A. Kharchilava¹⁷⁶, A. Mandal¹⁷⁶, M. Morris¹⁷⁶, D. Nguyen¹⁷⁶,
S. Rappoccio¹⁷⁶, H. Rejeb Sfar¹⁷⁶, A. Williams¹⁷⁶, P. Young¹⁷⁶, D. Yu¹⁷⁶, G. Alverson¹⁷⁷, E. Barberis¹⁷⁷,
J. Bonilla¹⁷⁷, B. Bylsma¹⁷⁷, M. Campana¹⁷⁷, J. Dervan¹⁷⁷, Y. Haddad¹⁷⁷, Y. Han¹⁷⁷, I. Israr¹⁷⁷, A. Krishna¹⁷⁷,
M. Lu¹⁷⁷, N. Manganelli¹⁷⁷, R. McCarthy¹⁷⁷, D. M. Morse¹⁷⁷, T. Orimoto¹⁷⁷, A. Parker¹⁷⁷, L. Skinnari¹⁷⁷,
C. S. Thoreson¹⁷⁷, E. Tsai¹⁷⁷, D. Wood¹⁷⁷, S. Dittmer¹⁷⁸, K. A. Hahn¹⁷⁸, Y. Liu¹⁷⁸, M. Mackenzie¹⁷⁸,
M. McGinnis¹⁷⁸, Y. Miao¹⁷⁸, D. G. Monk¹⁷⁸, M. H. Schmitt¹⁷⁸, A. Taliencio¹⁷⁸, M. Velasco¹⁷⁸, J. Wang¹⁷⁸,
G. Agarwal¹⁷⁹, R. Band¹⁷⁹, R. Bucci¹⁷⁹, S. Castells¹⁷⁹, A. Das¹⁷⁹, A. Ehnis¹⁷⁹, R. Goldouzian¹⁷⁹, M. Hildreth¹⁷⁹,
K. Hurtado Anampa¹⁷⁹, T. Ivanov¹⁷⁹, C. Jessop¹⁷⁹, A. Karneyeu¹⁷⁹, K. Lannon¹⁷⁹, J. Lawrence¹⁷⁹,
N. Loukas¹⁷⁹, L. Lutton¹⁷⁹, J. Mariano¹⁷⁹, N. Marinelli¹⁷⁹, I. Mcalister¹⁷⁹, T. McCauley¹⁷⁹, C. Mcgrady¹⁷⁹,
C. Moore¹⁷⁹, Y. Musienko^{179,nnnn}, H. Nelson¹⁷⁹, M. Osherson¹⁷⁹, A. Piccinelli¹⁷⁹, R. Ruchti¹⁷⁹, A. Townsend¹⁷⁹,
Y. Wan¹⁷⁹, M. Wayne¹⁷⁹, H. Yockey¹⁷⁹, A. Basnet¹⁸⁰, M. Carrigan¹⁸⁰, R. De Los Santos¹⁸⁰, L. S. Durkin¹⁸⁰,
C. Hill¹⁸⁰, M. Joyce¹⁸⁰, M. Nunez Ornelas¹⁸⁰, D. A. Wenzl¹⁸⁰, B. L. Winer¹⁸⁰, B. R. Yates¹⁸⁰, H. Bouchamaoui¹⁸¹,
K. Coldham¹⁸¹, P. Das¹⁸¹, G. Dezoort¹⁸¹, P. Elmer¹⁸¹, A. Frankenthal¹⁸¹, M. Galli¹⁸¹, B. Greenberg¹⁸¹,
N. Haubrich¹⁸¹, K. Kennedy¹⁸¹, G. Kopp¹⁸¹, Y. Lai¹⁸¹, D. Lange¹⁸¹, A. Loeliger¹⁸¹, D. Marlow¹⁸¹, I. Ojalvo¹⁸¹,
J. Olsen¹⁸¹, F. Simpson¹⁸¹, D. Stickland¹⁸¹, C. Tully¹⁸¹, S. Malik¹⁸², R. Sharma¹⁸², A. S. Bakshi¹⁸³,
S. Chandra¹⁸³, R. Chawla¹⁸³, A. Gu¹⁸³, L. Gutay¹⁸³, M. Jones¹⁸³, A. W. Jung¹⁸³, D. Kondratyev¹⁸³, M. Liu¹⁸³,
G. Negro¹⁸³, N. Neumeister¹⁸³, G. Paspalaki¹⁸³, S. Piperov¹⁸³, J. F. Schulte¹⁸³, F. Wang¹⁸³, A. Wildridge¹⁸³,
W. Xie¹⁸³, Y. Yao¹⁸³, Y. Zhong¹⁸³, N. Parashar¹⁸⁴, A. Pathak¹⁸⁴, E. Shumka¹⁸⁴, D. Acosta¹⁸⁵, A. Agrawal¹⁸⁵,
C. Arbour¹⁸⁵, T. Carnahan¹⁸⁵, K. M. Ecklund¹⁸⁵, P. J. Fernández Manteca¹⁸⁵, S. Freed¹⁸⁵, P. Gardner¹⁸⁵,
F. J. M. Geurts¹⁸⁵, T. Huang¹⁸⁵, I. Krommydas¹⁸⁵, N. Lewis¹⁸⁵, W. Li¹⁸⁵, J. Lin¹⁸⁵, O. Miguel Colin¹⁸⁵,
B. P. Padley¹⁸⁵, R. Redjimi¹⁸⁵, J. Rotter¹⁸⁵, E. Yigitbasi¹⁸⁵, Y. Zhang¹⁸⁵, O. Bessidskaia Bylund¹⁸⁶, A. Bodek¹⁸⁶,
P. de Barbaro^{186,a}, R. Demina¹⁸⁶, J. L. Dulemba¹⁸⁶, A. Garcia-Bellido¹⁸⁶, H. S. Hare¹⁸⁶, O. Hindrichs¹⁸⁶,
N. Parmar¹⁸⁶, P. Parygin^{186,nnnn}, R. Taus¹⁸⁶, B. Chiarito¹⁸⁷, J. P. Chou¹⁸⁷, S. V. Clark¹⁸⁷, S. Donnelly¹⁸⁷,
D. Gadkari¹⁸⁷, Y. Gershtein¹⁸⁷, E. Halkiadakis¹⁸⁷, M. Heindl¹⁸⁷, C. Houghton¹⁸⁷, D. Jaroslawski¹⁸⁷,
S. Konstantinou¹⁸⁷, I. Laflotte¹⁸⁷, A. Lath¹⁸⁷, J. Martins¹⁸⁷, B. Rand¹⁸⁷, J. Reichert¹⁸⁷, P. Saha¹⁸⁷, S. Salur¹⁸⁷,
S. Schnetzer¹⁸⁷, S. Somalwar¹⁸⁷, R. Stone¹⁸⁷, S. A. Thayil¹⁸⁷, S. Thomas¹⁸⁷, J. Vora¹⁸⁷, D. Ally¹⁸⁸,
A. G. Delannoy¹⁸⁸, S. Fiorendi¹⁸⁸, J. Harris¹⁸⁸, S. Higginbotham¹⁸⁸, T. Holmes¹⁸⁸, A. R. Kanuganti¹⁸⁸,
N. Karunarathna¹⁸⁸, J. Lawless¹⁸⁸, L. Lee¹⁸⁸, E. Nibigira¹⁸⁸, B. Skipworth¹⁸⁸, S. Spanier¹⁸⁸, D. Aebi¹⁸⁹,
M. Ahmad¹⁸⁹, T. Akhter¹⁸⁹, K. Androsov¹⁸⁹, A. Bolshov¹⁸⁹, O. Bouhali^{189,oooo}, R. Eusebi¹⁸⁹, P. Flanagan¹⁸⁹,
J. Gilmore¹⁸⁹, Y. Guo¹⁸⁹, T. Kamon¹⁸⁹, H. Kim¹⁸⁹, S. Luo¹⁸⁹, R. Mueller¹⁸⁹, A. Safonov¹⁸⁹, N. Akchurin¹⁹⁰,
J. Damgov¹⁹⁰, Y. Feng¹⁹⁰, N. Gogate¹⁹⁰, Y. Kazhykarim¹⁹⁰, K. Lamichhane¹⁹⁰, S. W. Lee¹⁹⁰, C. Madrid¹⁹⁰,
A. Mankel¹⁹⁰, T. Peltola¹⁹⁰, I. Volobouev¹⁹⁰, E. Appelt¹⁹¹, Y. Chen¹⁹¹, S. Greene¹⁹¹, A. Gurrola¹⁹¹, W. Johns¹⁹¹,
R. Kunnawalkam Elayavalli¹⁹¹, A. Melo¹⁹¹, D. Rathjens¹⁹¹, F. Romeo¹⁹¹, P. Sheldon¹⁹¹, S. Tuo¹⁹¹,
J. Velkovska¹⁹¹, J. Viinikainen¹⁹¹, J. Zhang¹⁹¹, B. Cardwell¹⁹², H. Chung¹⁹², B. Cox¹⁹², J. Hakala¹⁹²,
R. Hirosky¹⁹², M. Jose¹⁹², A. Ledovskoy¹⁹², C. Mantilla¹⁹², C. Neu¹⁹², C. Ramón Álvarez¹⁹², S. Bhattacharya¹⁹³,
P. E. Karchin¹⁹³, A. Aravind¹⁹⁴, S. Banerjee¹⁹⁴, K. Black¹⁹⁴, T. Bose¹⁹⁴, E. Chavez¹⁹⁴, S. Dasu¹⁹⁴,
P. Everaerts¹⁹⁴, C. Galloni¹⁹⁴, H. He¹⁹⁴, M. Herndon¹⁹⁴, A. Herve¹⁹⁴, C. K. Koraka¹⁹⁴, S. Lomte¹⁹⁴,
R. Loveless¹⁹⁴, A. Mallampalli¹⁹⁴, A. Mohammadi¹⁹⁴, S. Mondal¹⁹⁴, T. Nelson¹⁹⁴, G. Parida¹⁹⁴, L. Pétré¹⁹⁴,
D. Pinna¹⁹⁴, A. Savin¹⁹⁴, V. Shang¹⁹⁴, V. Sharma¹⁹⁴, W. H. Smith¹⁹⁴, D. Teague¹⁹⁴, H. F. Tsoi¹⁹⁴, W. Vetens¹⁹⁴,
A. Warden¹⁹⁴, S. Afanasiev¹⁹⁵, V. Alexakhin¹⁹⁵, Yu. Andreev¹⁹⁵, T. Aushev¹⁹⁵, D. Budkouski¹⁹⁵,
R. Chistov^{195,nnnn}, M. Danilov^{195,nnnn}, T. Dimova^{195,nnnn}, A. Ershov^{195,nnnn}, S. Gninenko¹⁹⁵, I. Gorbunov¹⁹⁵

A. Gribushin^{195,nnnn}, A. Kamenev¹⁹⁵, V. Karjavine¹⁹⁵, M. Kirsanov¹⁹⁵, V. Klyukhin^{195,nnnn}, O. Kodolova^{195,pppp},
 V. Korenkov¹⁹⁵, A. Kozyrev^{195,nnnn}, N. Krasnikov¹⁹⁵, A. Lanev¹⁹⁵, A. Malakhov¹⁹⁵, V. Matveev^{195,nnnn},
 A. Nikitenko^{195,qqqq,pppp}, V. Palichik¹⁹⁵, V. Perelygin¹⁹⁵, S. Petrushanko^{195,nnnn}, S. Polikarpov^{195,nnnn},
 O. Radchenko^{195,nnnn}, M. Savina¹⁹⁵, V. Shalaev¹⁹⁵, S. Shmatov¹⁹⁵, S. Shulha¹⁹⁵, Y. Skovpen^{195,nnnn},
 V. Smirnov¹⁹⁵, O. Teryaev¹⁹⁵, I. Tlisova^{195,nnnn}, A. Toropin¹⁹⁵, N. Voytishin¹⁹⁵, B. S. Yuldashev^{195,a,rrrr},
 A. Zarubin¹⁹⁵, I. Zhizhin¹⁹⁵, L. Dudko¹⁹⁶, K. Ivanov¹⁹⁶, V. Kim^{196,nnnn}, V. Murzin¹⁹⁶, V. Oreshkin¹⁹⁶,
 D. Sosnov¹⁹⁶, E. Boos¹⁹⁶, V. Bunichev¹⁹⁶, M. Dubinin^{196,gggg}, V. Savrin¹⁹⁶ and A. Snigirev¹⁹⁶

(CMS Collaboration)

¹*Yerevan Physics Institute, Yerevan, Armenia*

²*Institut für Hochenergiephysik, Vienna, Austria*

³*Universiteit Antwerpen, Antwerpen, Belgium*

⁴*Vrije Universiteit Brussel, Brussel, Belgium*

⁵*Université Libre de Bruxelles, Bruxelles, Belgium*

⁶*Ghent University, Ghent, Belgium*

⁷*Université Catholique de Louvain, Louvain-la-Neuve, Belgium*

⁸*Centro Brasileiro de Pesquisas Físicas, Rio de Janeiro, Brazil*

⁹*Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil*

¹⁰*Universidade Estadual Paulista, Universidade Federal do ABC, São Paulo, Brazil*

¹¹*Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Sofia, Bulgaria*

¹²*University of Sofia, Sofia, Bulgaria*

¹³*Instituto De Alta Investigación, Universidad de Tarapacá, Casilla 7 D, Arica, Chile*

¹⁴*Universidad Técnica Federico Santa María, Valparaíso, Chile*

¹⁵*Beihang University, Beijing, China*

¹⁶*Department of Physics, Tsinghua University, Beijing, China*

¹⁷*Institute of High Energy Physics, Beijing, China*

¹⁸*State Key Laboratory of Nuclear Physics and Technology, Peking University, Beijing, China*

¹⁹*State Key Laboratory of Nuclear Physics and Technology, Institute of Quantum Matter, South China Normal University, Guangzhou, China*

²⁰*Sun Yat-Sen University, Guangzhou, China*

²¹*University of Science and Technology of China, Hefei, China*

²²*Nanjing Normal University, Nanjing, China*

²³*Institute of Modern Physics and Key Laboratory of Nuclear Physics and Ion-beam Application (MOE)—Fudan University, Shanghai, China*

²⁴*Zhejiang University, Hangzhou, Zhejiang, China*

²⁵*Universidad de Los Andes, Bogota, Colombia*

²⁶*Universidad de Antioquia, Medellin, Colombia*

²⁷*University of Split, Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, Split, Croatia*

²⁸*University of Split, Faculty of Science, Split, Croatia*

²⁹*Institute Rudjer Boskovic, Zagreb, Croatia*

³⁰*University of Cyprus, Nicosia, Cyprus*

³¹*Charles University, Prague, Czech Republic*

³²*Escuela Politécnica Nacional, Quito, Ecuador*

³³*Universidad San Francisco de Quito, Quito, Ecuador*

³⁴*Academy of Scientific Research and Technology of the Arab Republic of Egypt, Egyptian Network of High Energy Physics, Cairo, Egypt*

³⁵*Center for High Energy Physics (CHEP-FU), Fayoum University, El-Fayoum, Egypt*

³⁶*National Institute of Chemical Physics and Biophysics, Tallinn, Estonia*

³⁷*Department of Physics, University of Helsinki, Helsinki, Finland*

³⁸*Helsinki Institute of Physics, Helsinki, Finland*

³⁹*Lappeenranta-Lahti University of Technology, Lappeenranta, Finland*

⁴⁰*IRFU, CEA, Université Paris-Saclay, Gif-sur-Yvette, France*

⁴¹*Laboratoire Leprince-Ringuet, CNRS/IN2P3, Ecole Polytechnique, Institut Polytechnique de Paris, Palaiseau, France*

⁴²*Université de Strasbourg, CNRS, IPHC UMR 7178, Strasbourg, France*

- ⁴³*Centre de Calcul de l'Institut National de Physique Nucleaire et de Physique des Particules, CNRS/IN2P3, Villeurbanne, France*
- ⁴⁴*Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France*
- ⁴⁵*Georgian Technical University, Tbilisi, Georgia*
- ⁴⁶*RWTH Aachen University, I. Physikalisches Institut, Aachen, Germany*
- ⁴⁷*RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany*
- ⁴⁸*RWTH Aachen University, III. Physikalisches Institut B, Aachen, Germany*
- ⁴⁹*Deutsches Elektronen-Synchrotron, Hamburg, Germany*
- ⁵⁰*University of Hamburg, Hamburg, Germany*
- ⁵¹*Karlsruher Institut fuer Technologie, Karlsruhe, Germany*
- ⁵²*Institute of Nuclear and Particle Physics (INPP), NCSR Demokritos, Aghia Paraskevi, Greece*
- ⁵³*National and Kapodistrian University of Athens, Athens, Greece*
- ⁵⁴*National Technical University of Athens, Athens, Greece*
- ⁵⁵*University of Ioánnina, Ioánnina, Greece*
- ⁵⁶*HUN-REN Wigner Research Centre for Physics, Budapest, Hungary*
- ⁵⁷*MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary*
- ⁵⁸*Faculty of Informatics, University of Debrecen, Debrecen, Hungary*
- ⁵⁹*HUN-REN ATOMKI—Institute of Nuclear Research, Debrecen, Hungary*
- ⁶⁰*Karoly Robert Campus, MATE Institute of Technology, Gyongyos, Hungary*
- ⁶¹*Panjab University, Chandigarh, India*
- ⁶²*University of Delhi, Delhi, India*
- ⁶³*University of Hyderabad, Hyderabad, India*
- ⁶⁴*Indian Institute of Technology Kanpur, Kanpur, India*
- ⁶⁵*Saha Institute of Nuclear Physics, HBNI, Kolkata, India*
- ⁶⁶*Indian Institute of Technology Madras, Madras, India*
- ⁶⁷*IISER Mohali, India, Mohali, India*
- ⁶⁸*Tata Institute of Fundamental Research-A, Mumbai, India*
- ⁶⁹*Tata Institute of Fundamental Research-B, Mumbai, India*
- ⁷⁰*National Institute of Science Education and Research, An OCC of Homi Bhabha National Institute, Bhubaneswar, Odisha, India*
- ⁷¹*Indian Institute of Science Education and Research (IISER), Pune, India*
- ⁷²*Indian Institute of Technology Hyderabad, Telangana, India*
- ⁷³*Isfahan University of Technology, Isfahan, Iran*
- ⁷⁴*Institute for Research in Fundamental Sciences (IPM), Tehran, Iran*
- ⁷⁵*University College Dublin, Dublin, Ireland*
- ^{76a}*INFN Sezione di Bari, Bari, Italy*
- ^{76b}*Università di Bari, Bari, Italy*
- ^{76c}*Politecnico di Bari, Bari, Italy*
- ^{77a}*INFN Sezione di Bologna, Bologna, Italy*
- ^{77b}*Università di Bologna, Bologna, Italy*
- ^{78a}*INFN Sezione di Catania, Catania, Italy*
- ^{78b}*Università di Catania, Catania, Italy*
- ^{79a}*INFN Sezione di Firenze, Firenze, Italy*
- ^{79b}*Università di Firenze, Firenze, Italy*
- ⁸⁰*INFN Laboratori Nazionali di Frascati, Frascati, Italy*
- ^{81a}*INFN Sezione di Genova, Genova, Italy*
- ^{81b}*Università di Genova, Genova, Italy*
- ^{82a}*INFN Sezione di Milano-Bicocca, Milano, Italy*
- ^{82b}*Università di Milano-Bicocca, Milano, Italy*
- ^{83a}*INFN Sezione di Napoli, Napoli, Italy*
- ^{83b}*Università di Napoli 'Federico II', Napoli, Italy*
- ^{83c}*Università della Basilicata, Potenza, Italy*
- ^{83d}*Scuola Superiore Meridionale (SSM), Napoli, Italy*
- ^{84a}*INFN Sezione di Padova, Padova, Italy*
- ^{84b}*Università di Padova, Padova, Italy*
- ^{84c}*Università degli Studi di Cagliari, Cagliari, Italy*
- ^{85a}*INFN Sezione di Pavia, Pavia, Italy*
- ^{85b}*Università di Pavia, Pavia, Italy*
- ^{86a}*INFN Sezione di Perugia, Perugia, Italy*

- ^{86b}Università di Perugia, Perugia, Italy
^{87a}INFN Sezione di Pisa, Pisa, Italy
^{87b}Università di Pisa, Pisa, Italy
^{87c}Scuola Normale Superiore di Pisa, Pisa, Italy
^{87d}Università di Siena, Siena, Italy
^{88a}INFN Sezione di Roma, Roma, Italy
^{88b}Sapienza Università di Roma, Roma, Italy
^{89a}INFN Sezione di Torino, Torino, Italy
^{89b}Università di Torino, Torino, Italy
^{89c}Università del Piemonte Orientale, Novara, Italy
^{90a}INFN Sezione di Trieste, Trieste, Italy
^{90b}Università di Trieste, Trieste, Italy
⁹¹Kyungpook National University, Daegu, Korea
⁹²Department of Mathematics and Physics—GWNu, Gangneung, Korea
⁹³Chonnam National University, Institute for Universe and Elementary Particles, Kwangju, Korea
⁹⁴Hanyang University, Seoul, Korea
⁹⁵Korea University, Seoul, Korea
⁹⁶Kyung Hee University, Department of Physics, Seoul, Korea
⁹⁷Sejong University, Seoul, Korea
⁹⁸Seoul National University, Seoul, Korea
⁹⁹University of Seoul, Seoul, Korea
¹⁰⁰Yonsei University, Department of Physics, Seoul, Korea
¹⁰¹Sungkyunkwan University, Suwon, Korea
¹⁰²College of Engineering and Technology, American University of the Middle East (AUM), Dasman, Kuwait
¹⁰³Kuwait University—College of Science—Department of Physics, Safat, Kuwait
¹⁰⁴Riga Technical University, Riga, Latvia
¹⁰⁵University of Latvia (LU), Riga, Latvia
¹⁰⁶Vilnius University, Vilnius, Lithuania
¹⁰⁷National Centre for Particle Physics, Universiti Malaya, Kuala Lumpur, Malaysia
¹⁰⁸Universidad de Sonora (UNISON), Hermosillo, Mexico
¹⁰⁹Centro de Investigación y de Estudios Avanzados del IPN, Mexico City, Mexico
¹¹⁰Universidad Iberoamericana, Mexico City, Mexico
¹¹¹Benemerita Universidad Autónoma de Puebla, Puebla, Mexico
¹¹²University of Montenegro, Podgorica, Montenegro
¹¹³University of Canterbury, Christchurch, New Zealand
¹¹⁴National Centre for Physics, Quaid-I-Azam University, Islamabad, Pakistan
¹¹⁵AGH University of Krakow, Krakow, Poland
¹¹⁶National Centre for Nuclear Research, Swierk, Poland
¹¹⁷Institute of Experimental Physics, Faculty of Physics, University of Warsaw, Warsaw, Poland
¹¹⁸Warsaw University of Technology, Warsaw, Poland
¹¹⁹Laboratório de Instrumentação e Física Experimental de Partículas, Lisboa, Portugal
¹²⁰Faculty of Physics, University of Belgrade, Belgrade, Serbia
¹²¹VINCA Institute of Nuclear Sciences, University of Belgrade, Belgrade, Serbia
¹²²Centro de Investigaciones Energéticas Medioambientales y Tecnológicas (CIEMAT), Madrid, Spain
¹²³Universidad Autónoma de Madrid, Madrid, Spain
¹²⁴Universidad de Oviedo, Instituto Universitario de Ciencias y Tecnologías Espaciales de Asturias (ICTEA), Oviedo, Spain
¹²⁵Instituto de Física de Cantabria (IFCA), CSIC-Universidad de Cantabria, Santander, Spain
¹²⁶University of Colombo, Colombo, Sri Lanka
¹²⁷University of Ruhuna, Department of Physics, Matara, Sri Lanka
¹²⁸CERN, European Organization for Nuclear Research, Geneva, Switzerland
¹²⁹PSI Center for Neutron and Muon Sciences, Villigen, Switzerland
¹³⁰ETH Zurich—Institute for Particle Physics and Astrophysics (IPA), Zurich, Switzerland
¹³¹Universität Zürich, Zurich, Switzerland
¹³²National Central University, Chung-Li, Taiwan
¹³³National Taiwan University (NTU), Taipei, Taiwan
¹³⁴High Energy Physics Research Unit, Department of Physics, Faculty of Science, Chulalongkorn University, Bangkok, Thailand
¹³⁵Tunis El Manar University, Tunis, Tunisia

- ¹³⁶*Çukurova University, Physics Department, Science and Art Faculty, Adana, Turkey*
¹³⁷*Middle East Technical University, Physics Department, Ankara, Turkey*
¹³⁸*Bogazici University, Istanbul, Turkey*
¹³⁹*Istanbul Technical University, Istanbul, Turkey*
¹⁴⁰*Istanbul University, Istanbul, Turkey*
¹⁴¹*Yildiz Technical University, Istanbul, Turkey*
¹⁴²*Institute for Scintillation Materials of National Academy of Science of Ukraine, Kharkiv, Ukraine*
¹⁴³*National Science Centre, Kharkiv Institute of Physics and Technology, Kharkiv, Ukraine*
¹⁴⁴*University of Bristol, Bristol, United Kingdom*
¹⁴⁵*Rutherford Appleton Laboratory, Didcot, United Kingdom*
¹⁴⁶*Imperial College, London, United Kingdom*
¹⁴⁷*Brunel University, Uxbridge, United Kingdom*
¹⁴⁸*Baylor University, Waco, Texas, USA*
¹⁴⁹*Bethel University, St. Paul, Minnesota, USA*
¹⁵⁰*Catholic University of America, Washington, DC, USA*
¹⁵¹*The University of Alabama, Tuscaloosa, Alabama, USA*
¹⁵²*Boston University, Boston, Massachusetts, USA*
¹⁵³*Brown University, Providence, Rhode Island, USA*
¹⁵⁴*University of California, Davis, Davis, California, USA*
¹⁵⁵*University of California, Los Angeles, California, USA*
¹⁵⁶*University of California, Riverside, Riverside, California, USA*
¹⁵⁷*University of California, San Diego, La Jolla, California, USA*
¹⁵⁸*University of California, Santa Barbara—Department of Physics, Santa Barbara, California, USA*
¹⁵⁹*California Institute of Technology, Pasadena, California, USA*
¹⁶⁰*Carnegie Mellon University, Pittsburgh, Pennsylvania, USA*
¹⁶¹*University of Colorado Boulder, Boulder, Colorado, USA*
¹⁶²*Cornell University, Ithaca, New York, USA*
¹⁶³*Fermi National Accelerator Laboratory, Batavia, Illinois, USA*
¹⁶⁴*University of Florida, Gainesville, Florida, USA*
¹⁶⁵*Florida State University, Tallahassee, Florida, USA*
¹⁶⁶*Florida Institute of Technology, Melbourne, Florida, USA*
¹⁶⁷*University of Illinois Chicago, Chicago, Illinois, USA*
¹⁶⁸*The University of Iowa, Iowa City, Iowa, USA*
¹⁶⁹*Johns Hopkins University, Baltimore, Maryland, USA*
¹⁷⁰*The University of Kansas, Lawrence, Kansas, USA*
¹⁷¹*Kansas State University, Manhattan, Kansas, USA*
¹⁷²*University of Maryland, College Park, Maryland, USA*
¹⁷³*Massachusetts Institute of Technology, Cambridge, Massachusetts, USA*
¹⁷⁴*University of Minnesota, Minneapolis, Minnesota, USA*
¹⁷⁵*University of Nebraska-Lincoln, Lincoln, Nebraska, USA*
¹⁷⁶*State University of New York at Buffalo, Buffalo, New York, USA*
¹⁷⁷*Northeastern University, Boston, Massachusetts, USA*
¹⁷⁸*Northwestern University, Evanston, Illinois, USA*
¹⁷⁹*University of Notre Dame, Notre Dame, Indiana, USA*
¹⁸⁰*The Ohio State University, Columbus, Ohio, USA*
¹⁸¹*Princeton University, Princeton, New Jersey, USA*
¹⁸²*University of Puerto Rico, Mayaguez, Puerto Rico, USA*
¹⁸³*Purdue University, West Lafayette, Indiana, USA*
¹⁸⁴*Purdue University Northwest, Hammond, Indiana, USA*
¹⁸⁵*Rice University, Houston, Texas, USA*
¹⁸⁶*University of Rochester, Rochester, New York, USA*
¹⁸⁷*Rutgers, The State University of New Jersey, Piscataway, New Jersey, USA*
¹⁸⁸*University of Tennessee, Knoxville, Tennessee, USA*
¹⁸⁹*Texas A&M University, College Station, Texas, USA*
¹⁹⁰*Texas Tech University, Lubbock, Texas, USA*
¹⁹¹*Vanderbilt University, Nashville, Tennessee, USA*
¹⁹²*University of Virginia, Charlottesville, Virginia, USA*
¹⁹³*Wayne State University, Detroit, Michigan, USA*
¹⁹⁴*University of Wisconsin—Madison, Madison, Wisconsin, USA*
¹⁹⁵*An institute or international laboratory covered by a cooperation agreement with CERN*

¹⁹⁶*An institute formerly covered by a cooperation agreement with CERN*

^aDeceased.

^bAlso at Yerevan State University, Yerevan, Armenia.

^cAlso at TU Wien, Vienna, Austria.

^dAlso at Ghent University, Ghent, Belgium.

^eAlso at Universidade do Estado do Rio de Janeiro, Rio de Janeiro, Brazil.

^fAlso at FACAMP—Faculdades de Campinas, Sao Paulo, Brazil.

^gAlso at Universidade Estadual de Campinas, Campinas, Brazil.

^hAlso at Federal University of Rio Grande do Sul, Porto Alegre, Brazil.

ⁱAlso at The University of the State of Amazonas, Manaus, Brazil.

^jAlso at University of Chinese Academy of Sciences, Beijing, China.

^kAlso at China Center of Advanced Science and Technology, Beijing, China.

^lAlso at University of Chinese Academy of Sciences, Beijing, China.

^mAlso at Henan Normal University, Xinxiang, China.

ⁿAlso at University of Shanghai for Science and Technology, Shanghai, China.

^oAlso at The University of Iowa, Iowa City, Iowa, USA.

^pAlso at Center for High Energy Physics, Peking University, Beijing, China.

^qAlso at Cairo University, Cairo, Egypt.

^rAlso at Suez University, Suez, Egypt.

^sAlso at British University in Egypt, Cairo, Egypt.

^tAlso at Purdue University, West Lafayette, Indiana, USA.

^uAlso at Université de Haute Alsace, Mulhouse, France.

^vAlso at Istinye University, Istanbul, Turkey.

^wAlso at Another institute or international laboratory covered by a cooperation agreement with CERN.

^xAlso at University of Hamburg, Hamburg, Germany.

^yAlso at RWTH Aachen University, III. Physikalisches Institut A, Aachen, Germany.

^zAlso at Bergische University Wuppertal (BUW), Wuppertal, Germany.

^{aa}Also at Brandenburg University of Technology, Cottbus, Germany.

^{bb}Also at Forschungszentrum Jülich, Juelich, Germany.

^{cc}Also at CERN, European Organization for Nuclear Research, Geneva, Switzerland.

^{dd}Also at HUN-REN ATOMKI—Institute of Nuclear Research, Debrecen, Hungary.

^{ee}Also at Universitatea Babes-Bolyai—Facultatea de Fizica, Cluj-Napoca, Romania.

^{ff}Also at MTA-ELTE Lendület CMS Particle and Nuclear Physics Group, Eötvös Loránd University, Budapest, Hungary.

^{gg}Also at HUN-REN Wigner Research Centre for Physics, Budapest, Hungary.

^{hh}Also at Physics Department, Faculty of Science, Assiut University, Assiut, Egypt.

ⁱⁱAlso at The University of Kansas, Lawrence, Kansas, USA.

^{jj}Also at Punjab Agricultural University, Ludhiana, India.

^{kk}Also at University of Hyderabad, Hyderabad, India.

^{ll}Also at Indian Institute of Science (IISc), Bangalore, India.

^{mm}Also at University of Visva-Bharati, Santiniketan, India.

ⁿⁿAlso at IIT Bhubaneswar, Bhubaneswar, India.

^{oo}Also at Institute of Physics, Bhubaneswar, India.

^{pp}Also at Deutsches Elektronen-Synchrotron, Hamburg, Germany.

^{qq}Also at Isfahan University of Technology, Isfahan, Iran.

^{rr}Also at Sharif University of Technology, Tehran, Iran.

^{ss}Also at Department of Physics, University of Science and Technology of Mazandaran, Behshahr, Iran.

^{tt}Also at Department of Physics, Faculty of Science, Arak University, ARAK, Iran.

^{uu}Also at Helwan University, Cairo, Egypt.

^{vv}Also at Italian National Agency for New Technologies, Energy and Sustainable Economic Development, Bologna, Italy.

^{ww}Also at Centro Siciliano di Fisica Nucleare e di Struttura Della Materia, Catania, Italy.

^{xx}Also at Università degli Studi Guglielmo Marconi, Roma, Italy.

^{yy}Also at Scuola Superiore Meridionale, Università di Napoli 'Federico II', Napoli, Italy.

^{zz}Also at Fermi National Accelerator Laboratory, Batavia, Illinois, USA.

^{aaa}Also at Lulea University of Technology, Lulea, Sweden.

^{bbb}Also at Consiglio Nazionale delle Ricerche—Istituto Officina dei Materiali, Perugia, Italy.

^{ccc}Also at UPES—University of Petroleum and Energy Studies, Dehradun, India.

^{ddd}Also at Institut de Physique des 2 Infinis de Lyon (IP2I), Villeurbanne, France.

^{eee}Also at Department of Applied Physics, Faculty of Science and Technology, Universiti Kebangsaan Malaysia, Bangi, Malaysia.

^{fff}Also at Trincomalee Campus, Eastern University, Sri Lanka, Nilaveli, Sri Lanka.

- ggg Also at Saegis Campus, Nugegoda, Sri Lanka.
- hhh Also at National and Kapodistrian University of Athens, Athens, Greece.
- iii Also at Ecole Polytechnique Fédérale Lausanne, Lausanne, Switzerland.
- jjj Also at Universität Zürich, Zurich, Switzerland.
- kkk Also at Stefan Meyer Institute for Subatomic Physics, Vienna, Austria.
- lll Also at Near East University, Research Center of Experimental Health Science, Mersin, Turkey.
- mmm Also at Konya Technical University, Konya, Turkey.
- nnn Also at Izmir Bakircay University, Izmir, Turkey.
- ooo Also at Adiyaman University, Adiyaman, Turkey.
- ppp Also at Bozok Universitetesi Rektörlüğü, Yozgat, Turkey.
- qqq Also at Istanbul Sabahattin Zaim University, Istanbul, Turkey.
- rrr Also at Marmara University, Istanbul, Turkey.
- sss Also at Milli Savunma University, Istanbul, Turkey.
- ttt Also at Informatics and Information Security Research Center, Gebze/Kocaeli, Turkey.
- uuu Also at Kafkas University, Kars, Turkey.
- vvv Also at Istanbul Okan University, Istanbul, Turkey.
- www Also at Hacettepe University, Ankara, Turkey.
- xxx Also at Erzincan Binali Yildirim University, Erzincan, Turkey.
- yyy Also at Istanbul University—Cerrahpasa, Faculty of Engineering, Istanbul, Turkey.
- zzz Also at Yildiz Technical University, Istanbul, Turkey.
- aaa Also at School of Physics and Astronomy, University of Southampton, Southampton, United Kingdom.
- bbb Also at Monash University, Faculty of Science, Clayton, Australia.
- ccc Also at Bethel University, St. Paul, Minnesota, USA.
- ddd Also at Università di Torino, Torino, Italy.
- eee Also at Karamanoğlu Mehmetbey University, Karaman, Turkey.
- fff Also at California Lutheran University, Thousand Oaks, California, USA.
- ggg Also at California Institute of Technology, Pasadena, California, USA.
- hhh Also at United States Naval Academy, Annapolis, Maryland, USA.
- iii Also at Bingol University, Bingol, Turkey.
- jjj Also at Georgian Technical University, Tbilisi, Georgia.
- kkk Also at Sinop University, Sinop, Turkey.
- lll Also at Erciyes University, Kayseri, Turkey.
- mmm Also at Horia Hulubei National Institute of Physics and Nuclear Engineering (IFIN-HH), Bucharest, Romania.
- nnn Also at Another institute formerly covered by a cooperation agreement with CERN.
- ooo Also at Hamad Bin Khalifa University (HBKU), Doha, Qatar.
- ppp Also at Yerevan Physics Institute, Yerevan, Armenia.
- qqq Also at Imperial College, London, United Kingdom.
- rrr Also at Institute of Nuclear Physics of the Uzbekistan Academy of Sciences, Tashkent, Uzbekistan.