Comparative dosimetric analysis of deep inspiration breath-hold versus free breathing radiotherapy in lung cancer: special emphasis on cardiac and subcardiac structure protection

Von der Medizinischen Fakultät
der Rheinisch-Westfälischen Technischen Hochschule Aachen
zur Erlangung des akademischen Grades
eines Doktors der Medizin
genehmigte Dissertation

vorgelegt von

Biney Pal Singh

aus

Solingen

Berichter: Univ.-Prof. Dr. med. Michael J. Eble
Univ.-Prof. Dr. med. Philipp Bruners

Tag der mündlichen Prüfung: 28.07.2026

Diese Dissertation ist auf den Internetseiten der Universitätsbibliothek online verfügbar.

Meinem Bruder und meinen Eltern – in Liebe und Dankbarkeit.

Auflistung der eigenen Publikationen → *Dissertation darstellend*

Biney Pal Singh, Ahmed Ali Chughtai, Jennifer Stock, Philipp Bruners, Michael J. Eble, Ahmed Allam Mohamed,
Comparative dosimetric analysis of deep inspiration breath-hold versus free breathing radiotherapy in lung cancer: special emphasis on cardiac and subcardiac structure protection, Technical Innovations & Patient Support in Radiation Oncology, Volume 37, March 2026, 100371

Technical Innovations & Patient Support in Radiation Oncology 37 (2026) 100371

Contents lists available at ScienceDirect

Technical Innovations & Patient Support in Radiation Oncology

journal homepage: www.sciencedirect.com/journal/technical-innovations-and-patient-support-in-radiation-oncology

Research Article

Comparative dosimetric analysis of deep inspiration breath-hold versus free breathing radiotherapy in lung cancer: special emphasis on cardiac and subcardiac structure protection

Biney Pal Singh [a,c,*], Ahmed Ali Chughtai [a,c], Jennifer Stock [a,c], Philipp Bruners [b,c], Michael J. Eble [a,c], Ahmed Allam Mohamed [a,c]

[a] Department of Radiation Oncology, Medical Faculty, RWTH Aachen University Hospital, Aachen, Germany
[b] Department of Diagnostic and Interventional Radiology, RWTH Aachen University Hospital, Aachen, Germany
[c] Center for Integrated Oncology Aachen, Bonn, Cologne and Duesseldorf (CIO ABCD), Aachen, Germany

ARTICLE INFO

Keywords:
Deep inspiration breath-hold
Lung cancer
Cardiac substructures
Respiratory gating
Dose sparing

ABSTRACT

Background: Radiation therapy (RT) plays a central role in the treatment of lung cancer. However, cardiac and pulmonary toxicities remain major concerns. Deep inspiration breath-hold (DIBH) has been shown to improve target stability and reduce organ-at-risk (OAR) exposure, but data in lung cancer, particularly regarding cardiac substructures, remain limited.

Materials and methods: We retrospectively analysed 32 patients with lung cancer treated with RT at our institution between 2020 and 2021. Each patient underwent planning CTs in both DIBH and free breathing (FB), generating a total of 64 treatment plans using VMAT (60 Gy in 30 fractions). Contouring followed ESTRO ACROP guidelines, including detailed segmentation of cardiac chambers and coronary arteries. Dosimetric endpoints included Dmean, D2%, and VxGy for lung, heart, and subcardiac structures. Statistical comparisons between DIBH and FB plans were performed.

Results: DIBH significantly increased lung volume and improved all lung dosimetric parameters, including Dmean (13.13 vs. 14.49 Gy, $p < 0.001$) and V20Gy (24.37 % vs. 27.28 %, $p < 0.001$). Cardiac sparing was evident, with lower heart D2% (39.91 vs. 41.67 Gy, $p = 0.012$) and V45Gy (2.23 % vs. 2.53 %, $p = 0.005$). Notably, significant reductions were observed in the LAD (Dmean: 5.59 vs. 6.57 Gy, $p = 0.004$), LCX, and LV. Subgroup analyses demonstrated consistent dosimetric advantages across tumor laterality and level 7 nodal involvement.

Conclusions: DIBH offers substantial and consistent dosimetric benefits in lung cancer RT, reducing radiation burden to pulmonary, cardiac, and coronary structures. Given the known association between cardiac dose and long-term morbidity, these findings support the broader implementation of DIBH, especially in anatomically challenging cases involving subcarinal nodes or left-sided tumors.

Introduction

With ongoing demographic shifts, namely population aging, growth, and rising socioeconomic disparities, the global incidence and morbidity of cancer continue to increase.[1] Among these, with an estimated 2.5 million cases, lung cancer is the most common tumour, and the most common tumour in terms of deaths, with 1.8 million, in 2022, showing a global increase in the last decade.[1,2] In this context, not only is a well-structured healthcare system essential, but also the optimized implementation of therapeutic modalities. For patients with locally advanced or inoperable lung cancer, radiotherapy (RT) plays a pivotal role, underscoring the need for continual evaluation and refinement of its application.

Over the past decades, advances such as intensity-modulated radiotherapy (IMRT), volumetric modulated arc therapy (VMAT), and stereotactic ablative radiotherapy (SABR) have significantly enhanced the precision and conformality of RT [3]. Nevertheless, one of the persistent challenges in thoracic irradiation lies in managing breathing-induced motion, which affects both target coverage and sparing of organs at risk (OARs). To mitigate these uncertainties, techniques such as deep

* Corresponding author at: Universitätsklinikum RWTH Aachen, Pauwelsstraße 30, 52074 Aachen, Germany.
E-mail address: bsingh@ukaachen.de (B.P. Singh).

https://doi.org/10.1016/j.tipsro.2025.100371
Received 30 August 2025; Received in revised form 11 December 2025; Accepted 14 December 2025
Available online 15 December 2025
2405-6324/© 2025 The Author(s). Published by Elsevier B.V. on behalf of European Society for Radiotherapy & Oncology. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

inspiration breath-hold (DIBH) have been developed [4]. As early as the 1990 s, studies demonstrated the clinical feasibility of breath-hold strategies, showing that tumor immobilization is both tolerable and reproducible for patients, while also enabling improved tumor coverage and reduced lung dose [5,6]. Since then, a growing body of literature has confirmed the advantages of DIBH over free breathing (FB) in thoracic RT, including significant reductions in lung and cardiac doses, and associated improvements in clinical outcomes, most notably, reductions in rates of radiation pneumonitis and overall mortality [7–10].

Due to its anatomical and functional centrality, the heart and its substructures are frequently exposed during lung RT, making them vulnerable to both acute and late toxicities. Multiple studies have identified dose–response relationships between irradiation of cardiac and subcardiac structures and subsequent cardiovascular events [11–14]. These effects not only limit the potential for safe dose escalation in lung cancer but may also adversely impact overall survival through increased risks of ischemia, arrhythmias, and heart failure. The reported incidence of severe (grade $\geq$ 3) cardiac events following chemoradiotherapy ranges from 10 % to 33 % [11,12,15,16].

With improving survival outcomes in lung cancer, there is a growing imperative to assess pre-treatment cardiovascular risk better and to integrate strategies for effective cardiac sparing within RT planning [13,16]. Against this backdrop, the present study aims to investigate the potential benefits of DIBH in RT planning for patients with lung cancer, particularly in comparison to FB with respect to dosimetric exposure of subcardiac structures, including heart chambers and coronary arteries.

Materials and methods

This retrospective study included 32 patients with lung cancer who received radiotherapy at our institution between January 1, 2020, and December 31, 2021. Each patient underwent planning computed tomography (P-CT) using both deep inspiration breath-hold (DIBH) and free-breathing four-dimensional CT (4D-CT) techniques, resulting in a total of 64P-CT scans.

Imaging and treatment planning

P-CTs were acquired using a 16-slice CT scanner (Brilliance CT Big Bore Oncology, Philips Medical Systems, Cleveland, OH, USA) with settings: 120 kV, 146 mAs, pitch of 0.813, and slice thickness of 3 mm. DIBH CT scans utilized Catalyst™ (CRAD, Uppsala, Sweden), an optical surface imaging system monitoring the sternal area movement.

Positron emission tomography (PET-CT) scans were available for selected patients and helped with contour delineation. Data were anonymized and imported into Pinnacle3 treatment planning software (version 14.0; Philips Healthcare, Amsterdam, Netherlands).

Volume delineation

Target volumes were delineated following ESTRO ACROP guidelines. Gross Tumor Volume (GTVp) was defined on lung window CT images; internal Gross Tumor Volume (iGTVp) was delineated for FB-CT using respiratory phases (0 %, 50 %, 90 %). A margin of 6–8 mm (depending on histology: 6 mm for squamous cell carcinoma, 8 mm for adenocarcinoma and small cell lung carcinoma) around GTV within lung tissue formed Clinical Target Volumes (CTVp/iCTVp). Enlarged or metabolically active lymph nodes on contrast-enhanced CT or PET-CT were delineated as nodal GTV (GTVn). Mediastinal lymph node stations were delineated as nodal CTV (CTVn), excluding uninvolved critical structures (heart, great vessels, trachea, bronchial tree, esophagus). The Planning Target Volume (PTV) was created by expanding the combined CTV (CTVp + CTVn) by 5 mm isotropically.

Lungs were delineated separately and then combined to form the total lung. Additionally, the heart with its four chambers −the left atrium (LA), right atrium (RA), left ventricle (LV), and right ventricle (RV) − was defined. The right and left coronary arteries (RCA and LCA), along with their main branches, namely the posterior descending artery (PDA), left anterior descending (LAD), and left circumflex artery (LCX), were outlined as subcardial structures based on delineation atlases from Duane F et al. [17].

Initial delineations were performed by two radiation oncology trainees (BS and AC) and then reviewed by two experienced radiation oncologists (ME and AAM). In cases of discrepancy, contours were revised through consensus with a radiologist (PB).

Treatment planning and evaluation

VMAT plans, consisting of two full coplanar arcs with 6 MV photon energy, were generated for both FB and DIBH CT scans, totaling 64 plans. Dose calculations used a collapsed cone algorithm with a $3 \times 3 \times 3$ mm^3 dose grid. The prescribed dose was 60 Gy delivered in 30 fractions. Treatment planning was optimized to ensure a uniform dose distribution across the PTV, aiming to deliver at least 95 % of the prescribed dose to 100 % of the CTV and more than 95 % of the PTV. Plan quality was evaluated using the following dose–volume metrics: D2% (approximate maximum dose to the target/OAR), D98% (approximate minimum dose to the target), Dmean (average dose to the target or OAR), Dx (dose received by x% of the target/OAR volume), and VxGy (percentage of volume receiving at least x Gy). Organ-at-risk constraints included: spinal canal D2% $<$ 45 Gy, mean liver dose $<$ 32 Gy, mean lung dose $<$ 17.5 Gy with V20Gy $<$ 20 %, and mean heart dose $<$ 26 Gy.

Since the primary aim of this study was to examine the intrinsic dose reduction achieved with DIBH, no explicit dose constraints or tolerance thresholds for subcardiac structures were adhered to while treatment planning. Nevertheless, our findings are later discussed in the light of reported constraints – succinctly summarized by Walls G et. al. [16].

Dosimetric analysis

For each DIBH and FB plan a dose-volume histogram (DVH) was generated. From these various dosimetric and volumetric parameters were obtained:

- PTV volume, as well as D2%, D50% and D98% of the PTV
- VRI, i.e. the volume of PTV enclosed by certain – in our case the 95 % − isodose line

Using these parameters following indices were calculated:

- conformity index (CI) = VRI/PTV
- homogeneity index (HI) = (D2% − D98%)/IR * 100, where RI represents the reference isodose

In addition to that, for comparing analysis of DIBH and FB plans following dosimetric data was recorded:

Lung, total:	Dmean (Gy), V20Gy (%), V40Gy (%)
Lung, ipsilateral and contralateral:	Dmean (Gy), V5Gy (%), V10Gy (%), V20 (%), V40 (%)
Heart:	D2% (Gy), V20 Gy (%), V45 Gy (%), V60 Gy (%), Dmean (Gy)
Cardiac chambers: • left atrium and ventricle • right atrium and ventricle	D2% (Gy), V10 Gy (%), V20 Gy (%), V30 Gy (%), V40 Gy (%), Dmean (Gy)
Coronary arteries: • left coronary artery • LAD • LCX • right coronary artery • PDA	D2% (Gy), V5 Gy (%), V10 Gy (%), V20 Gy (%), V30 Gy (%), V40 Gy (%), Dmean (Gy)

Statistical analysis

Collected data was further analyzed using Microsoft Excel (Microsoft Corporation, Redmond, Washington, USA) and Python (Version 3.13.2); graphs were generated with GraphPad Prism (GraphPad Software, Boston, Massachusetts, USA) and Excel.

First, the normal distribution of the data was assessed with the Shapiro-Wilk-Test, using a p-value threshold of 0.05. Datasets with a p-value $>$ 0.05 were considered normally distributed and were compared using a parametric, paired *t*-test. If one of the datasets was classified as

Table 1
Volumetric and dosimetric characteristics of investigated radiation plans in deep inspiration breath-hold (DIBH) and free breathing (FB).

Volumetric parameter/ dosimetric index	FB	DIBH	Test	p-value
PTV volume	511.32 cm^3	496.24 cm^2	a	0.061
CI	1.02	1.01	a	0.36
HI	8.68	9.23	a	0.09
MU	656.23	695.79	a	0.12

non-parametric, the Wilcoxon signed-rank test was applied. In both tests, a p-value < 0.05 indicated a significant difference between the compared data groups.

Ethical statement/considerations:

The study was approved by the local ethics committee of RWTH Aachen University (Approval No.: EK 25–232).

Results

Patient Characteristics

The study cohort comprised 32 patients with a median age of 69 years. Histologically, patients presented predominantly with squamous cell carcinoma (n = 12), small cell lung carcinoma (n = 9), adenocarcinoma (n = 8), and mixed or large cell neuroendocrine carcinoma (n = 3). Detailed staging and tumor localization are summarized in Supplementary Table 1.

Target volume and dosimetric indices

The mean PTV was slightly reduced in DIBH; 496.24 cm^3 compared to FB; 511.32 cm^3, though this difference did not reach statistical significance (p = 0.061). Both CI and HI demonstrated comparable results between techniques (CI: DIBH = 1.01, FB = 1.02, p = 0.36; HI: DIBH = 9.23, FB = 8.68, p = 0.09) (Table 1).

Lung dosimetry

Relative to FB 4D-CT, DIBH scan produced markedly larger lung volumes, expanding the whole lung by an average of 1,602.7 cm^3, the ipsilateral lung by 722.11 cm^3, and the contralateral lung by 873.25 cm^3.

This volumetric gain translated into clear dosimetric advantages: the lung Dmean fell from 14.49 Gy with FB to 13.13 Gy with DIBH (p < 0.001), as well as in volume parameters V20Gy (24.37 % vs. 27.28 %, p < 0.001) and V40Gy (6.67 % vs. 8.40 %, p < 0.001) (Fig. 1, 2Table 2).

Focusing on the ipsilateral lung, DIBH consistently reduced every evaluated parameter, including Dmean (18.11 vs. 20.46 Gy, p < 0.001), V5Gy, V10Gy, V20Gy, and V40Gy. The contralateral lung demonstrated significant improvements, primarily in Dmean (9.08 vs. 9.62 Gy, p = 0.009) and V5Gy (54.57 % vs. 58.28 %, p = 0.001) (Fig. 1, Table 2).

Cardiac and subcardiac structure dosimetry

The DIBH approach resulted in a significant reduction in heart volume (516.4 vs 547.3 cm^3, p < 0.001). Additionally, significant dosimetric improvements were observed in the heart's D2% (39.91 Gy in




Fig. 1. Boxplots showing dosimetric outcome of lung total (A), lung ipsilateral (B), lung contralateral (C) and heart (D) comparing results of DIBH plan (colourful) and FB plan (grey).

Table 2
All significantly different dosimetric parameters in planning in deep inspiration breath-hold (DIBH) and free breathing (FB), a = paired *t*-test, b = Wilcoxon signed-rank test.

	FB	DIBH		Test	p-value
Lung total					
Dmean (Gy)	14.49	13.13	−9.39	a	**2.59365E-05**
V20Gy (%)	27.28	24.37	−10.67	a	**0.000430897**
V40Gy / %	8.4	6.67	−20.60	a	**1.27E-05**
Lung ipsilateral					
Dmean / Gy	20.46	18.11	−11.49	a	**8.28E-06**
V5Gy / %	69.47	65.27	−6.05	a	**8.65E-05**
V10Gy / %	60.36	55.79	−7.57	a	**9.56E-05**
V20Gy / %	44.77	39.43	−11.93	a	**4.04E-05**
V40Gy / %	17.23	13.53	−21.47	a	**3.85E-06**
Lung contralateral					
Dmean / Gy	9.62	9.08	−5.61	a	**0.009350664**
V5Gy / %	58.28	54.57	−6.37	a	**0.001208776**
Heart					
D2% / Gy	41.67	39.91	−4.22	b	**0.012457265**
V45 Gy / %	2.53	2.23	−11.86	b	**0.00548476**
Left ventricle					
D2% / Gy	12.38	10.53	−14.94	b	**0.00968003**
Dmean / Gy	4.15	3.76	−9.40	b	**0.008160867**
Right atrium					
D2% / Gy	26.79	23.84	−11.01	b	**0.031105267**
V30 Gy / %	4.68	3.27	−30.13	b	**0.012407092**
V40 Gy / %	2.13	1.39	−34.74	b	**0.000843013**
Left coronary artery					
D2% / Gy	15.23	13.05	−14.31	b	**0.020216676**
Left circumflex coronary					
V5 Gy / %	49.33	43.55	−11.72	b	**0.002699796**
Dmean / Gy	9.6	8.85	−7.81	b	**0.014803257**
Left anterior descending					
D2% / Gy	15.21	12.32	−19.00	b	**0.00154797**
V5 Gy / %	49.68	39.39	−20.71	b	**0.001435352**
Dmean / Gy	6.57	5.59	−14.92	b	**0.00438754**
Posterior descending artery					
D2% / Gy	2.2	2.08	−5.45	b	**0.044224421**
Dmean / Gy	1.63	1.6	−1.84	b	**0.026056039**

DIBH vs. 41.67 Gy in FB, p = 0.012) and V45 Gy (2.23 % in DIBH vs. 2.53 % in FB, p = 0.005). Heart Dmean and V20Gy showed reductions in DIBH but did not reach statistical significance (p = 0.078 and p = 0.110, respectively) (Fig. 1, Table 2). No significant difference was observed for V60G.

Subcardiac analysis showed that DIBH significantly reduced radiation exposure to the LV, with decreases in D2% (10.53 vs. 12.38 Gy, p = 0.009) and Dmean (3.76 vs. 4.15 Gy, p = 0.008). No significant dosimetric differences were found for the RV (Fig. 2, Table 2).

For atrial structures, significant dose reductions with DIBH were limited to the RA, including D2% (23.84 vs. 26.79 Gy, p = 0.031), V30Gy (3.27 % vs. 4.68 %, p = 0.012), and V40Gy (1.39 % vs. 2.13 %, p < 0.001).

In the coronary arteries, DIBH significantly reduced D2% for the LCA (13.05 vs. 15.23 Gy, p = 0.020), LAD (12.32 vs. 15.21 Gy, p = 0.002), and PDA (2.08 vs. 2.20 Gy, p = 0.044). Additionally, Dmean was significantly lower for LAD (5.59 vs. 6.57 Gy, p = 0.004), PDA (1.60 vs. 1.63 Gy, p = 0.026), and LCX (8.85 vs. 9.60 Gy, p = 0.015). No significant differences were observed for the RCA (Fig. 2, Table 2).

Subgroup analyses

A subgroup analysis was performed to assess the effect of tumor laterality (left versus right lung tumors) and the presence or absence of subcarinal lymph node metastases (thoracic level station seven, "LS 7) on the observed dosimetric advantages of DIBH. Laterality of tumors was evenly distributed – 17 patients had tumors of the left and 15 patients of the right lung. While 22 patients demonstrated involvement of lymph node station 7, it was absent in 10.

For tumors of the left lung, DIBH significantly enhanced pulmonary and cardiac dosimetry. For the total lung, Dmean decreased from 14.04 Gy to 12.86 Gy (p = 0.002), with notable reductions in V20Gy (25.72 % to 23.72 %, p = 0.032) and V40Gy (7.60 % to 6.31 %, p = 0.002). Ipsilateral lung metrics consistently showed significant improvements, with decreases in Dmean (from 20.20 Gy to 17.94 Gy, p < 0.001) and V40Gy (from 16.56 % to 13.23 %, p < 0.001). For the contralateral lung, both Dmean (9.88 Gy to 9.30 Gy, p = 0.042) and V5Gy (58.89 % to 54.60 %, p = 0.006) significantly declined. Cardiac exposure was significantly reduced in both the LCX (D2%: 29.83 Gy to 25.46 Gy, p = 0.005; Dmean: 13.48 Gy to 11.99 Gy, p = 0.008) and LAD (D2%: 18.66 Gy to 14.41 Gy, p = 0.015; Dmean: 8.34 Gy to 7.09 Gy, p = 0.043) (Table 3).

For right-lung tumors, significant dosimetric benefits were observed in the total lung, including reductions in Dmean (from 15 Gy to 13.44 Gy, p = 0.005), V20Gy (from 29.05 % to 25.11 %, p = 0.006), and V40Gy (from 9.31 % to 7.09 %, p = 0.002). The ipsilateral lung showed significant decreases in Dmean (20.75 Gy to 18.30 Gy, p = 0.007), V20Gy (46.07 % to 39.80 %, p = 0.004), and V40Gy (18.00 % to 13.86 %, p < 0.001). Cardiac sparing was also prominent, as demonstrated by reductions in heart V45Gy (2.83 % to 2.47 %, p = 0.009), LV Dmean (2.42 Gy to 2.23 Gy, p = 0.012), and substantial decreases in RA- dose metrics, particularly D2% (39.85 Gy to 32.89 Gy, p = 0.005) and V40Gy (4.51 % to 2.93 %, p = 0.001). Additionally, V5Gy of LAD decreased from 39.73 % to 26.45 % (p = 0.021) (Table 3).

Further, patients with positive subcarinal lymph nodes demonstrated particularly pronounced dosimetric advantages with DIBH. Cardiac dosimetry showed improvements, including heart V45Gy (3.08 % to 2.72 %, p = 0.012), and significant reductions in LV D2% (13.51 Gy to 11.01 Gy, p = 0.017). Coronary artery exposure was also reduced, particularly in LAD D2% (16.30 Gy to 13.67 Gy, p = 0.007), Dmean (7.14 Gy to 6.09 Gy, p = 0.016), and PDA's Dmean (1.87 Gy to 1.87 Gy, p = 0.040) (Table 4).

In patients without subcarinal lymph node involvement, significant improvements in coronary artery exposure were limited to the LCA D2% (11.12 Gy to 9.61 Gy, p = 0.014) and LCX V5Gy (32.59 % to 26.70 %, p = 0.028) (Table 4).

Overall, across all evaluated subgroups, DIBH consistently maintained or enhanced the favorable dosimetric profile compared to FB, without statistically significant deterioration of any evaluated parameter.

Discussion

The pivotal role of radiotherapy in lung cancer management is well-established, as is its association with potential adverse cardiac effects. Evidence from recent studies has emphasized that radiation exposure to cardiac structures, particularly the coronary arteries and cardiac chambers, significantly increases the risk of major adverse cardiac events (MACE), arrhythmias, and mortality [18,19].

Atkins et al. highlighted that an increased mean cardiac dose and LAD V15Gy ≥ 10 % are independent predictors of MACE and all-cause mortality. [18] Similar findings were reinforced by systematic reviews, underscoring the necessity of dose optimization to cardiac and subcardiac structures to mitigate cardiovascular morbidity and mortality following RT for lung cancer [14,16].

Beyond immediate dosimetric considerations, the long-term implications of cardiac exposure during thoracic RT have gained increasing attention. Accelerated coronary artery disease, often termed RICAD, is now recognized as a distinct clinical entity, often affecting proximal coronary segments such as the LAD and LCX, with a latency period that may extend beyond a decade post-RT. As highlighted by Kirresh et al., even moderate doses to these vessels can accelerate atherosclerotic changes, leading to clinically significant coronary events, particularly in patients with pre-existing cardiovascular risk factors.[20] Importantly, RICAD is frequently underdiagnosed due to its overlap with traditional ischemic heart disease and the lack of routine cardiac surveillance in long-term cancer survivors.

Also, Wiedemann et al. showed that thoracic radiotherapy is




Fig. 2. Linear diagrams showing relative volumes in relation to dose for the left ventricle (E), right ventricle (F), left atrium (G), right atrium (H), left coronary artery (I), left circumflex coronary (J), left anterior descending (K), right coronary artery (L), posterior descending artery (M).

associated with remodelling processes of the pulmonary vessels, which can be associated with cardiac diseases.[21] The incidences of severe cardiac events, i.e. grade 3 and higher, in patients with lung cancer after chemoradiotherapy range from 10 to 33 %, depending on the observed study [11,12,15,16], making prudent radiotherapy planning with consistent protection of the pulmonary and cardiac OARs essential. The use of DIBH can contribute to that goal.

The positive effects of DIBH use on the dosimetric outcome for pulmonary, cardiac, and subcardiac structures have been recorded in various studies. General effects have been described in all thoracic malignancies. Benefits for cardiac substructures are most particularly demonstrated in breast cancer patients [22–26]. A comparative study for a group of oesophageal cancer patients was conducted, emphasising the importance of DIBH for dose reduction in cardiac substructures [27]. To our knowledge, our now presented work is the first to give a detailed overview of dosimetric changes in cardiac substructures in DIBH compared to FB in lung cancer.

Analysing the entire group, a significant improvement was observed in all dosimetric parameters of the total and ipsilateral lung. Additionally, parameters of the contralateral lung improved with DIBH, but only Dmean and V5Gy showed significant changes. The underlying mechanisms, namely, an increase in lung volume and a reduction of target volume mobility and lung tissue density, have been described, and their benefits have been repeatedly demonstrated [5–7,9,10,28].

The observed cardiac sparing effects, notably significant reductions in heart's D2% and V45Gy, echo the results of previously published literature, which reported improved dosimetric outcomes with the implementation of DIBH [29]. Recent Systematic reviews have consistently highlighted a dose–response relationship between cardiac irradiation and subsequent cardiovascular events, reinforcing the importance of precise cardiac dose management [14,16].

In evaluating specific cardiac substructures, we observed significant dosimetric improvements in the LV and RA, results that should be considered alongside the findings by Jang et al., who identified the LV as especially sensitive to radiation dose and strongly linked to increased cardiac event risks [12]. Our findings on coronary artery sparing, particularly the LAD, LCX, and PDA, provide important evidence supporting the adoption of DIBH in clinical practice as possible to potentially reduce ischemic cardiac events [30,31].

A notable finding in our analysis was the varied impact of DIBH

Table 3
All significantly different dosimetric parameters in planning in deep inspiration breath-hold (DIBH) and free breathing (FB) of patients with left or right lungs tumours, a = paired *t*-test, b = Wilcoxon signed-rank test.

Tumor in the LEFT Lung	**FB**	**DIBH**		**Test**	**p-value**
Lung total					
Dmean (Gy)	14.04	12.86	−8.40	a	**0.001714318**
V20Gy (%)	25.72	23.72	−7.78	a	**0.031620242**
V40Gy / %	7.6	6.31	−16.97	a	**0.002472702**
Lung ipsilateral					
Dmean / Gy	20.2	17.94	−11.19	a	**0.000261375**
V5Gy / %	70.38	65.35	−7.15	a	**6.25577E-05**
V10Gy / %	60.21	55.38	−8.02	a	**0.000407702**
V20Gy / %	43.61	39.11	−10.32	a	**0.004194853**
V40Gy / %	16.56	13.23	−20.11	a	**0.000332789**
Lung contralateral					
Dmean / Gy	9.88	9.3	−5.87	a	**0.042332474**
V5Gy / %	58.89	54.6	−7.28	a	**0.006134563**
Left circumflex coronary					
D2% / Gy	29.83	25.46	−14.65	b	**0.004638672**
V5 Gy / %	56.32	50.19	−10.88	b	**0.009925486**
Dmean / Gy	13.48	11.99	−11.05	b	**0.007904053**
Left anterior descending					
D2% / Gy	18.66	14.41	−22.78	a	**0.015299659**
V5 Gy / %	58.46	50.81	−13.09	a	**0.017054667**
Dmean / Gy	8.34	7.09	−14.99	a	**0.042789071**
Tumor in the RIGHT Lung					
Lung total					
Dmean (Gy)	15	13.44	−10.40	a	**0.005285044**
V20Gy (%)	29.05	25.11	−13.56	a	**0.006200701**
V40Gy / %	9.31	7.09	−23.85	a	**1.59E-03**
Lung ipsilateral					
Dmean / Gy	20.75	18.3	−11.81	a	**7.35E-03**
V10Gy / %	60.53	56.25	−7.07	a	**3.55E-02**
V20Gy / %	46.07	39.8	−13.61	a	**4.30E-03**
V40Gy / %	18	13.86	−23.00	b	**6.10E-04**
Heart					
V45 Gy / %	2.83	2.47	−12.72	a	**0.008692475**
Dmean / Gy	7.17	6.53	−8.93	a	**0.034038065**
Left ventricle					
Dmean / Gy	2.42	2.23	−7.85	b	**0.012451172**
Right atrium					
D2% / Gy	39.85	32.89	−17.47	a	**0.005337672**
V20 Gy / %	16.79	12.99	−22.63	b	**0.033046944**
V30 Gy / %	9.56	6.49	−32.11	b	**0.003701749**
V40 Gy / %	4.51	2.93	−35.03	b	**0.00146885**
Dmean / Gy	10.09	8.55	−15.26	b	**0.008361816**
Left anterior descending					
V5 Gy / %	39.73	26.45	−33.43	a	**0.020891279**

Table 4
All significantly different dosimetric parameters in planning in deep inspiration breath-hold (DIBH) and free breathing (FB) of patients with or without involvement of lymph nodes station 7, a = paired *t*-test, b = Wilcoxon signed-rank test.

LS 7 positive	**FB**	**DIBH**		**Test**	p-value
Lung total					
Dmean (Gy)	16.11	14.52	−9.87	a	**0.00020377**
V20Gy (%)	31.13	27.62	−11.28	a	**0.001696284**
V40Gy / %	8.77	6.94	−20.87	a	**2.98E-04**
Lung ipsilateral					
Dmean / Gy	21.27	18.89	−11.19	a	**1.95E-04**
V5Gy / %	73.92	69.72	−5.68	a	**7.30E-04**
V10Gy / %	64.44	60.16	−6.64	a	**1.95E-03**
V20Gy / %	46.77	41.28	−11.74	a	**9.25E-04**
V40Gy / %	17.06	13.36	−21.69	a	**4.77E-07**
Lung contralateral					
Dmean / Gy	11.64	10.83	−6.96	a	**0.003460223**
V5Gy / %	68.14	62.86	−7.75	a	**0.000397692**
V10Gy / %	49.53	46.47	−6.18	a	**0.018867291**
Heart					
V45 Gy / %	3.08	2.72	−11.69	b	**0.01163652**
Left ventricle					
D2% / Gy	13.51	11.01	−18.50	b	**0.017245169**
Dmean / Gy	4.42	3.94	−10.86	b	**0.036495955**
Right atrium					
V40 Gy / %	2.22	1.54	−30.63	b	**0.003494192**
Left circumflex coronary					
V5 Gy / %	56.94	51.21	−10.06	b	**0.02490692**
Dmean / Gy	10.51	9.75	−7.23	b	**0.03587532**
Left anterior descending					
D2% / Gy	16.3	13.67	−16.13	b	**0.007443428**
V5 Gy / %	54.3	44.05	−18.88	a	**0.005592619**
Dmean / Gy	7.14	6.09	−14.71	b	**0.015698997**
Posterior descending artery					
Dmean / Gy	1.87	1.87	0.00	b	**0.039833215**
LS 7 negative					
Lung total					
V40Gy / %	7.59	6.09	−19.76	a	**2.23E-02**
Lung ipsilateral					
Dmean / Gy	18.66	16.38	−12.22	a	**2.38E-02**
V10Gy / %	51.39	46.16	−10.18	a	**2.71E-02**
V20Gy / %	40.35	35.38	−12.32	a	**2.21E-02**
V40Gy / %	17.63	13.89	−21.21	a	**2.08E-02**
Left coronary artery					
D2% / Gy	11.12	9.61	−13.58	a	**0.014399122**
Left circumflex coronary					
V5 Gy / %	32.59	26.7	−18.07	b	**0.027707849**

based on tumor laterality. Patients with left-sided tumors showed significant reductions in coronary artery doses. Specifically, V5Gy, D2%, and Dmean of the LCX and LAD were significantly lowered by DIBH, highlighting the benefit of sparing the LAD in reducing cardiac events after RT, which is well documented in breast cancer. However, unlike the entire cohort, there was no significant dose reduction in heart parameters overall or specifically in the left ventricle. The most noteworthy improvement in the group with right lung tumors was observed in parameters related to the right atrium. Patients experienced dose reductions across nearly all parameters, including V5 Gy of the LAD. The varying degrees of dose reduction can be explained by the different radiation exposure of the respective structures and the movement and deformation of the heart during DIBH. The LCX and LAD which due to their left-sided location are more affected while irradiation of same-sided tumors they experience a dose improvement because of the compression of the heart and the shift of caudal parts of heart from the left to the center. In contrast, the burden on the right atrium is considerably higher in radiation of right-sided carcinomas. Therefore, the narrowing of the heart while DIBH shows more pronounced dose sparing in these tumors [8].

Given these findings, we further explored whether the benefit of DIBH is maintained in anatomically complex cases, specifically when RT fields include LS 7 due to nodal involvement in the inferior mediastinum and proximity to the heart, we specifically analysed the impact of DIBH on RT dose distribution in relation to whether this nodal station required inclusion in the target volume. In patients with LS 7-positive disease, DIBH significantly reduced radiation exposure to both pulmonary and cardiac OARs compared to FB. Lung sparing was consistent across all evaluated parameters, including total lung Dmean, V20, and V40, as well as ipsilateral and contralateral lung sub-volumes, indicating that the benefits of DIBH are maintained even when target volumes extend to the inferior mediastinum. Moreover, cardiac dose was also significantly reduced, with lower V45, Dmean and D2% of the LV, and V5/Dmean of both the LCX and LAD, underscoring the cardioprotective potential of DIBH in this setting. In general, it should be noted that in our cohort the baseline dose exposure of pulmonary and cardiac structures is lower in LS 7-negative carcinomas. Previous studies have already shown, for example, significantly higher cardiac doses due to multistation nodal involvement [32]. With considerably higher exposure in subcarinal involvement, there is greater potential for dose reduction in this subgroup when DIBH is used.

The overall cardioprotective potential is underlined even more when above mentioned findings are placed in context of published dose relationships and thresholds for cardiac events, for example compiled by

Walls G et al. [16] Among others the Dmean of both ventricles and the LAD is listed as a parameter correlating with ischemia and heart failure in radiated patients with lung cancer. DIBH usage reduced – even though not significantly – the Dmean of the right ventricle in the total, the right tumor cohort, and both LS 7 subgroups. Same metric was bettered significantly for the left ventricle in total cohort, for right-sided and LS 7 positive tumors. LAD's Dmean was improved significantly in the complete cohort, and the left tumour and LS 7 positive subgroups. The V30 of the left ventricle is also associated with the development of ischemia; DIBH decreased the value in the total cohort, and in left sided and LS 7 positive tumors.

Since neither all volumetric and dosimetric values were assessed, nor all structures were contoured, our data cannot completely be positioned within existing explicit thresholds and constraints for all adverse cardiac events. Nevertheless, the parameters which are collected already emphasize the benefit of DIBH for patients' clinical outcome after lung cancer therapy in the light of published literature.

Despite these promising findings, the implementation of DIBH in clinical practice is not without challenges. Effective DIBH requires consistent patient cooperation and breath-hold reproducibility, which can be difficult for individuals with compromised pulmonary function or limited compliance. Thus, although all patients in our cohort were initially able to perform DIBH-CT, only eleven patients' radiation plans were ultimately planned on those. Motion management technologies such as surface-guided RT or respiratory gating systems are often necessary to monitor breath-hold accuracy but add technical complexity and require additional resources. Moreover, integrating DIBH into routine clinical workflows demands dedicated staff training and patient preparation, which may impact efficiency in high-volume centers [33,34]. These considerations must be weighed when evaluating the feasibility and scalability of DIBH in broader clinical practice.

Study limitations

Our study has several limitations that should be acknowledged. Firstly, the retrospective design and relatively small sample size may limit the generalizability of our findings and the statistical power to detect subtle differences. Secondly, the study lacks clinical follow-up data to correlate dosimetric advantages with actual clinical outcomes such as reduced cardiac and pulmonary toxicity. Additionally, interobserver variability in contour delineation, despite standardized guidelines and consensus reviews, could have introduced minor inconsistencies. Furthermore, patient compliance and reproducibility of the DIBH technique could vary, potentially affecting dosimetric accuracy. Lastly, due to the retrospective nature of the study, potential confounding variables such as patient comorbidities and baseline cardiac function were not fully controlled, necessitating future prospective studies to validate our results comprehensively.

Conclusion

Overall, our study robustly supports the implementation of DIBH in lung cancer radiotherapy planning, demonstrating dosimetric benefits in the majority of cardiac and pulmonary parameters, and in a multitude of these even significant improvements. Aligning with recommendations from recent systematic reviews, our results reinforce the importance of individualized radiation planning, meticulous cardiac structure delineation, and proactive dose-sparing strategies to optimize therapeutic outcomes and patient quality of life.

Waiver of patient consent

This is a retrospective case study. Patient consent has been waived by Ethic committee.

Author Contributions:

Manuscript drafting and editing (BPS), revision of Manuscript (AMM, MJE), Conception and design of the study (BPS, AAC, MJE), Segmentation of target volumes and OAR (BPS, AAC), Revision of the segmentation (PB, AAM, MJE), Radiation treatment planning (JS) Data collection (BPS), statistical analysis (BPS, AAM), Revision of the analysis (AAM, MJE).

Funding

This research received no specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Appendix A. Supplementary material

Supplementary data to this article can be found online at https://doi.org/10.1016/j.tipsro.2025.100371.

References

[1] Bray F, Laversanne M, Sung H, et al. Global cancer statistics 2022: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. CA Can J Clin 2024;74(3):229–63.
[2] Collaborators GBDRTC. Global, regional, and national burden of respiratory tract cancers and associated risk factors from 1990 to 2019: a systematic analysis for the Global Burden of Disease Study 2019. Lancet Respir Med 2021; 9(9): 1030-49.
[3] Brown S, Banfill K, Aznar MC, Whitehurst P, Faivre FC. The evolving role of radiotherapy in non-small cell lung cancer. Br J Radiol 2019;92(1104):20190524.
[4] Wong JW, Sharpe MB, Jaffray DA, et al. The use of active breathing control (ABC) to reduce margin for breathing motion. Int J Radiat Oncol 1999;44(4):911–9.
[5] Hanley J, Debois MM, Mah D, et al. Deep inspiration breath-hold technique for lung tumors: the potential value of target immobilization and reduced lung density in dose escalation. Int J Radiat Oncol Biol Phys 1999;45(3):603–11.
[6] Mah D, Hanley J, Rosenzweig KE, et al. Technical aspects of the deep inspiration breath-hold technique in the treatment of thoracic cancer. Int J Radiat Oncol Biol Phys 2000;48(4):1175–85.
[7] Barnes EIA, Murray BR, Robinson DM, Underwood LJ, Hanson J, Roa WHY. Dosimetric evaluation of lung tumor immobilization using breath hold at deep inspiration. Int J Radiat Oncol 2001;50(4):1091–8.
[8] Fjellanger K, Rossi L, Heijmen BJM, et al. Patient selection, inter-fraction plan robustness and reduction of toxicity risk with deep inspiration breath hold in intensity-modulated radiotherapy of locally advanced non-small cell lung cancer. Front Oncol 2022;12:966134.
[9] Giraud P, Morvan E, Claude L, et al. Respiratory gating techniques for optimization of lung cancer radiotherapy. J Thorac Oncol 2011;6(12):2058–68.
[10] Marchand V, Zefkili S, Desrousseaux J, Simon L, Dauphinot C, Giraud P. Dosimetric comparison of free-breathing and deep inspiration breath-hold radiotherapy for lung cancer. Strahlenther Onkol 2012;188(7):582–9.
[11] Dess RT, Sun Y, Matuszak MM, et al. Cardiac events after radiation therapy: combined analysis of prospective multicenter trials for locally advanced non-small-cell lung cancer. J Clin Oncol 2017;35(13):1395–402.
[12] Jang BS, Cha MJ, Kim HJ, et al. Heart substructural dosimetric parameters and risk of cardiac events after definitive chemoradiotherapy for stage III non-small cell lung cancer. Radiother Oncol 2020;152:126–32.
[13] Vojtisek R. Cardiac toxicity of lung cancer radiotherapy. Rep Pract Oncol Radiother 2020;25(1):13–9.
[14] Mitchell JD, Cehic DA, Morgia M, et al. Cardiovascular manifestations from therapeutic radiation: a multidisciplinary expert consensus statement from the international cardio-oncology society. JACC CardioOncol 2021;3(3):360–80.
[15] Ming X, Feng Y, Yang C, Wang W, Wang P, Deng J. Radiation-induced heart disease in lung cancer radiotherapy: a dosimetric update. Medicine (Baltimore) 2016;95 (41):e5051.
[16] Walls GM, Bergom C, Mitchell JD, et al. Cardiotoxicity following thoracic radiotherapy for lung cancer. Br J Can 2025;132(4):311–25.
[17] Duane F, Aznar MC, Bartlett F, et al. A cardiac contouring atlas for radiotherapy. Radiother Oncol 2017;122(3):416–22.
[18] Atkins KM, Chaunzwa TL, Lamba N, et al. Association of left anterior descending coronary artery radiation dose with major adverse cardiac events and mortality in patients with non-small cell lung cancer. JAMA Oncol 2021;7(2):206–19.
[19] Atkins KM, Rawal B, Chaunzwa TL, et al. Cardiac radiation dose, cardiac disease, and mortality in patients with lung cancer. J Am Coll Cardiol 2019;73(23): 2976–87.
[20] Kirresh A, White L, Mitchell A, et al. Radiation-induced coronary artery disease: a difficult clinical conundrum. Clin Med (Lond) 2022;22(3):251–6.
[21] Wiedemann J, Coppes RP, van Luijk P. Radiation-induced cardiac side-effects: the lung as target for interacting damage and intervention. Front Oncol 2022;12: 931023.

[22] Joo JH, Kim SS, Ahn SD, et al. Cardiac dose reduction during tangential breast irradiation using deep inspiration breath hold: a dose comparison study based on deformable image registration. Radiat Oncol 2015;10:264.
[23] Bruzzaniti V, Abate A, Pinnaro P, et al. Dosimetric and clinical advantages of deep inspiration breath-hold (DIBH) during radiotherapy of breast cancer. J Exp Clin Can Res 2013;32(1):88.
[24] Wolf J, Stoller S, Lubke J, et al. Deep inspiration breath-hold radiation therapy in left-sided breast cancer patients: a single-institution retrospective dosimetric analysis of organs at risk doses. Strahlenther Onkol 2023;199(4):379–88.
[25] Pandeli C, Smyth LML, David S, See AW. Dose reduction to organs at risk with deep-inspiration breath-hold during right breast radiotherapy: a treatment planning study. Radiat Oncol 2019;14(1):223.
[26] Schonecker S, Heinz C, Sohn M, et al. Reduction of cardiac and coronary artery doses in irradiation of left-sided breast cancer during inspiration breath hold : a planning study. Strahlenther Onkol 2016;192(11):750–8.
[27] Mohamed AA, Douglas MN, Bruners P, Eble MJ. Dosimetric advantages for cardiac substructures in radiotherapy of esophageal cancer in deep-inspiration breath hold. Strahlenther Onkol 2024;200(7):624–32.
[28] Berson AM, Emery R, Rodriguez L, et al. Clinical experience using respiratory gated radiation therapy: comparison of free-breathing and breath-hold techniques. Int J Radiat Oncol Biol Phys 2004;60(2):419–26.
[29] Yan D, Ning L, Chen Y, et al. Analysis of deep inspiration breath-hold technique to improve dosimetric and clinical advantages in postoperative intensity-modulated radiation therapy for thymomas. Quant Imaging Med Surg 2022;12(8):4239–47.
[30] Daher J, Rizza A, Tonacci A, Borghini A. A comprehensive review of radiotherapy-induced coronary artery disease-epidemiology, biological mechanisms, and preventive strategies. Int J Mol Sci 2025;26(11).
[31] Milo MLH, Thorsen LBJ, Johnsen SP, et al. Risk of coronary artery disease after adjuvant radiotherapy in 29,662 early breast cancer patients: a population-based Danish Breast Cancer Group study. Radiother Oncol 2021;157:106–13.
[32] McNew LK, Bowen SR, Gopan O, et al. The relationship between cardiac radiation dose and mediastinal lymph node involvement in stage III non-small cell lung cancer patients. Adv Radiat Oncol 2017;2(2):192–6.
[33] Reitz D, Walter F, Schonecker S, et al. Stability and reproducibility of 6013 deep inspiration breath-holds in left-sided breast cancer. Radiat Oncol 2020;15(1):121.
[34] Gaál S, Kahán Z, Paczona V, et al. Deep-inspirational breath-hold (DIBH) technique in left-sided breast cancer: various aspects of clinical utility. Radiat Oncol 2021;16(1).

Patient number	Age	Sex	Histology	Tumour stage	Tumour localisation
1	64	m	adenocarcinoma	cT4, N2	right upper lobe
2	73	f	squamous cell carcinoma	cT3, N0	right upper lobe
3	78	m	squamous cell carcinoma	cT4, N2	left lower lobe
4	78	m	squamous cell carcinoma	cT3, N2	left lower lobe
5	58	m	small cell lung carcinoma	cT1b, N2	left upper lobe
6	65	f	adenocarcinoma	cT2b, N2	right upper lobe
7	76	m	adenocarcinoma	cT3, N3	left lower lobe
8	75	m	squamous cell carcinoma	cT3, N1	left upper lobe
9	69	f	small cell lung carcinoma	cT3, N2	right lower lobe
10	79	m	squamous cell carcinoma	cT3, N1	left lower lobe
11	53	m	adenocarcinoma	cT4, N3	left lower lobe
12	68	f	small cell lung carcinoma	cT4, N3	left upper lobe
13	68	f	squamous cell carcinoma	cT3, N0	right lower lobe
14	79	m	small cell lung carcinoma	cT3, N3	right upper lobe
15	71	f	adenocarcinoma	cT1c, N3	left upper lobe
16	74	m	squamous cell carcinoma	cT2a, N3	lingula
17	59	f	small cell lung carcinoma	cT2, N3	right upper lobe
18	69	m	squamous cell carcinoma	cT2a, N1	right upper lobe
19	56	f	adenocarcinoma	cT1c, N3	right upper lobe
20	79	m	squamous cell carcinoma	cT3, N3	left upper lobe
21	64	m	squamous cell carcinoma	cT4, N3	left upper lobe
22	55	m	large cell neuroendocrine carcinoma	cT4, N2	right lung, beyond lobes
23	85	m	squamous cell carcinoma	cT3, N2	right middle lobe
24	54	m	large cell neuroendocrine carcinoma	cT1a, N3	right upper lobe
25	72	F	small cell lung carcinoma	cT3, N1	left lower lobe

26	86	m	small cell lung carcinoma	cT4, N2	right upper lobe
27	63	m	adenocarcinoma	cT1b, cN2	left upper lobe
28	62	m	small cell lung carcinoma	cT3, N2	right lung, beyond lobes
29	64	m	small cell lung carcinoma	cT2a, N3	left upper lobe
30	71	m	adenocarcinoma	cT2a, N2	left lower lobe
31	66	m	squamous cell carcinoma	cT2a, N2	right upper lobe
32	80	f	large cell neuroendocrine carcinoma	cT4, N0	left lower lobe

Supplementary Table 1: Patients characteristics.

	FB	DIBH	Mean relative difference (%)	Test	p-value
Lung total					
Dmean (Gy)	14.49	13.13	-9.39	a	**2.59365E-05**
V20Gy (%)	27.28	24.37	-10.67	a	**0.000430897**
V40Gy / %	8.4	6.67	-20.60	a	**1.27E-05**
Lung ipsilateral					
Dmean / Gy	20.46	18.11	-11.49	a	**8.28E-06**
V5Gy / %	69.47	65.27	-6.05	a	**8.65E-05**
V10Gy / %	60.36	55.79	-7.57	a	**9.56E-05**
V20Gy / %	44.77	39.43	-11.93	a	**4.04E-05**
V40Gy / %	17.23	13.53	-21.47	a	**3.85E-06**
Lung contralateral					
Dmean / Gy	9.62	9.08	-5.61	a	**0.009350664**
V5Gy / %	58.28	54.57	-6.37	a	**0.001208776**
V10Gy / %	39.27	37.72	-3.95	a	0.125608751
V20Gy / %	12.98	12.01	-7.47	b	0.643191592
V40Gy / %	1.18	1.16	-1.69	b	0.887979046
Heart					
D2% / Gy	41.67	39.91	-4.22	b	**0.012457265**
V20 Gy / %	9.07	8.15	-10.14	a	0.110277863
V45 Gy / %	2.53	2.23	-11.86	b	**0.00548476**
V60 Gy/ %	0.4	0.38	-5.00	b	0.696346664
Dmean / Gy	7.27	6.91	-4.95	a	0.077763842
Left ventricle					
D2% / Gy	12.38	10.53	-14.94	b	**0.00968003**
V10 Gy / %	10.68	9.33	-12.64	b	0.395696234
V20 Gy / %	2.51	1.78	-29.08	b	0.202328308
V30 Gy / %	0.76	0.51	-32.89	b	0.32698935
V40 Gy / %	0.3	0.19	-36.67	b	0.345231072
Dmean / Gy	4.15	3.76	-9.40	b	**0.008160867**
Right ventricle					
D2% / Gy	7.44	6.68	-10.22	b	0.268159443
V10 Gy / %	3.73	2.65	-28.95	b	0.57598703
V20 Gy / %	0.23	0.33	43.48	b	1
V30 Gy / %	0	0	0.00		
V40 Gy / %	0	0	0.00		
Dmean / Gy	2.88	2.69	-6.60	b	0.245303636
Left atrium					
D2% / Gy	48.22	47.44	-1.62	b	0.295070265
V10 Gy / %	48.03	47.81	-0.46	b	0.991794348

V20 Gy / %	27.81	28.76	3.42	a	0.560752212
V30 Gy / %	18.21	18.86	3.57	a	0.585957439
V40 Gy / %	12.26	12.8	4.40	b	0.478455666
Dmean / Gy	15.94	16.1	1.00	a	0.794055103
Right atrium					
D2% / Gy	26.79	23.84	-11.01	b	**0.031105267**
V10 Gy / %	18.47	17.62	-4.60	b	0.374043408
V20 Gy / %	8.6	7.21	-16.16	b	0.326027571
V30 Gy / %	4.68	3.27	-30.13	b	**0.012407092**
V40 Gy / %	2.13	1.39	-34.74	b	**0.000843013**
Dmean / Gy	7.18	6.58	-8.36	b	0.112336656
Left coronary artery					
D2% / Gy	15.23	13.05	-14.31	b	**0.020216676**
V5 Gy / %	87.66	84.01	-4.16	b	0.172954918
V10 Gy / %	51.45	46.8	-9.04	b	0.454888845
V20 Gy / %	12.53	8.98	-28.33	b	0.423596318
V30 Gy / %	3.23	2.72	-15.79	b	0.179712495
V40 Gy / %	0	0			
Dmean / Gy	11.63	10.51	-9.63	b	0.06784779
Left circumflex coronary					
D2% / Gy	21.09	19.43	-7.87	b	0.09773239
V5 Gy / %	49.33	43.55	-11.72	b	**0.002699796**
V10 Gy / %	29.1	26.96	-7.35	b	0.469162525
V20 Gy / %	11.48	11.58	0.87	b	0.876701302
V30 Gy / %	7.75	6.83	-11.87	b	0.498962299
V40 Gy / %	5.23	4.22	-19.31	b	0.128190174
Dmean / Gy	9.6	8.85	-7.81	b	**0.014803257**
Left anterior descending					
D2% / Gy	15.21	12.32	-19.00	b	**0.00154797**
V5 Gy / %	49.68	39.39	-20.71	b	**0.001435352**
V10 Gy / %	20.98	17.54	-16.40	b	
V20 Gy / %	4.29	3.08	-28.21	b	0.151956186
V30 Gy / %	1.53	0.7	-54.25	b	0.179712495
V40 Gy / %	0.46	0.06	-86.96	b	0.179712495
Dmean / Gy	6.57	5.59	-14.92	b	**0.00438754**
Right coronary artery					
D2% / Gy	7.17	6.87	-4.18	b	0.845056152
V5 Gy / %	19.05	20.25	6.30	b	0.497914888
V10 Gy / %	5.21	4.69	-9.98	b	0.514669723
V20 Gy / %	1.21	1.63	34.71	b	1
V30 Gy / %	0	0			
V40 Gy / %	0	0			

Dmean / Gy	3.4	3.27	-3.82	b	0.270850908
Posterior descending artery					
D2% / Gy	2.2	2.08	-5.45	b	**0.044224421**
V5 Gy / %	5.32	7.39	38.91	b	0.465208819
V10 Gy / %	0	0.58		b	0.317310508
V20 Gy / %	0	0			
V30 Gy / %	0	0			
V40 Gy / %	0	0			
Dmean / Gy	1.63	1.6	-1.84	b	**0.026056039**

Supplementary Table 2: Dosimetric outcomes of planning in deep inspiration breath-hold (DIBH) and free breathing (FB) for the patients of the complete cohort.

LEFT (n = 17)	FB	DIBH	Mean relative difference (%)	Test	p-value
Lung total					
Dmean (Gy)	14.04	12.86	-8.40	a	**0.001714318**
V20Gy (%)	25.72	23.72	-7.78	a	**0.031620242**
V40Gy / %	7.6	6.31	-16.97	a	**2.47E-03**
Lung ipsilateral					
Dmean / Gy	20.2	17.94	-11.19	a	**2.61E-04**
V5Gy / %	70.38	65.35	-7.15	a	**6.26E-05**
V10Gy / %	60.21	55.38	-8.02	a	**4.08E-04**
V20Gy / %	43.61	39.11	-10.32	a	**4.19E-03**
V40Gy / %	16.56	13.23	-20.11	a	**3.33E-04**
Lung contralateral					
Dmean / Gy	9.88	9.3	-5.87	a	**0.042332474**
V5Gy / %	58.89	54.6	-7.28	a	**0.006134563**
V10Gy / %	40.23	37.62	-6.49	a	0.059191333
V20Gy / %	13.45	12.83	-4.61	a	0.425869854
V40Gy / %	1.49	1.56	4.70	b	0.530284597
Heart					
D2% / Gy	38.68	37.18	-3.88	b	0.127030095
V20 Gy / %	8.89	8.43	-5.17	a	0.54134466
V45 Gy / %	2.26	2.01	-11.06	b	0.157708987
V60 Gy/ %	0.3	0.3	0.00	b	0.811892227
Dmean / Gy	7.35	7.25	-1.36	a	0.703084138
Left ventricle					
D2% / Gy	17.62	14.54	-17.48	b	0.055719324
V10 Gy / %	18.93	15.85	-16.27	b	0.332879755
V20 Gy / %	4.73	3.35	-29.18	b	0.213122469
V30 Gy / %	1.43	0.96	-32.87	b	0.32698935
V40 Gy / %	0.57	0.36	-36.84	b	0.345231072
Dmean / Gy	5.67	5.11	-9.88	b	0.124772736
Right ventricle					
D2% / Gy	6.82	6.88	0.88	a	0.877284229
V10 Gy / %	1.91	1.9	-0.52	b	0.475245832
V20 Gy / %	0	0			
V30 Gy / %	0	0			
V40 Gy / %	0	0			
Dmean / Gy	2.82	2.87	1.77	a	0.743237702
Left atrium					
D2% / Gy	44.27	43.63	-1.45	b	0.352874756
V10 Gy / %	50.48	51.69	2.40	b	0.325870024

V20 Gy / %	30.43	30.62	0.62	a	0.934782077
V30 Gy / %	19.41	19.78	1.91	a	0.838995175
V40 Gy / %	12.86	13.2	2.64	b	0.861304257
Dmean / Gy	16.46	16.62	0.97	a	0.84938169
Right atrium					
D2% / Gy	15.27	15.86	3.86	a	0.616057924
V10 Gy / %	9.85	10.65	8.12	b	0.753152365
V20 Gy / %	1.38	2.1	52.17	b	0.239135782
V30 Gy / %	0.38	0.42	10.53	b	1
V40 Gy / %	0.03	0.03	0.00	b	0.712701857
Dmean / Gy	4.62	4.83	4.55	a	0.397485277
Left coronary artery					
D2% / Gy	16.38	14.4	-12.09	a	0.147839759
V5 Gy / %	86.74	81.86	-5.63	b	0.179712495
V10 Gy / %	59.31	54.25	-8.53	b	0.284502698
V20 Gy / %	18.28	12.08	-33.92	b	0.463071015
V30 Gy / %	5.84	5.1	-12.67	b	0.317310508
V40 Gy / %	0	0			
Dmean / Gy	12.98	11.68	-10.02	a	0.173949769
Left circumflex coronary					
D2% / Gy	29.83	25.46	-14.65	b	**0.004638672**
V5 Gy / %	56.32	50.19	-10.88	b	**0.009925486**
V10 Gy / %	43.02	38.22	-11.16	b	0.099539688
V20 Gy / %	21.48	18.75	-12.71	b	0.332879755
V30 Gy / %	14.59	12.85	-11.93	b	0.498962299
V40 Gy / %	9.84	7.95	-19.21	b	0.128190174
Dmean / Gy	13.48	11.99	-11.05	b	**0.007904053**
Left anterior descending					
D2% / Gy	18.66	14.41	-22.78	a	**0.015299659**
V5 Gy / %	58.46	50.81	-13.09	a	**0.017054667**
V10 Gy / %	32.59	27.89	-14.42	b	
V20 Gy / %	7.91	5.42	-31.48	b	0.085830958
V30 Gy / %	2.87	1.31	-54.36	b	0.179712495
V40 Gy / %	0.87	0.12	-86.21	b	0.179712495
Dmean / Gy	8.34	7.09	-14.99	a	**0.042789071**
Right coronary artery					
D2% / Gy	6	6.26	4.33	a	0.543565372
V5 Gy / %	18.21	19.26	5.77	b	0.929152765
V10 Gy / %	2.06	2.27	10.19	b	1
V20 Gy / %	0	0			
V30 Gy / %	0	0			
V40 Gy / %	0	0			

Dmean / Gy	2.92	2.85	-2.40	b	0.690885645
Posterior descending artery					
D2% / Gy	2.44	2.18	-10.66	b	0.221854908
V5 Gy / %	7.83	8.12	3.70	b	1
V10 Gy / %	0	0			
V20 Gy / %	0	0			
V30 Gy / %	0	0			
V40 Gy / %	0	0			
Dmean / Gy	1.86	1.81	-2.69	b	0.132064599

Supplementary Table 3: Dosimetric outcomes of planning in deep inspiration breath-hold (DIBH) and free breathing (FB) for the patients with left lung tumors.

RIGHT (n = 15)	FB	DIBH	Mean relative difference (%)	Test	p-value
Lung total					
Dmean (Gy)	15	13.44	-10.40	a	**0.005285044**
V20Gy (%)	29.05	25.11	-13.56	a	**0.006200701**
V40Gy / %	9.31	7.09	-23.85	a	**1.59E-03**
Lung ipsilateral					
Dmean / Gy	20.75	18.3	-11.81	a	**7.35E-03**
V5Gy / %	68.43	65.17	-4.76	a	7.32E-02
V10Gy / %	60.53	56.25	-7.07	a	**3.55E-02**
V20Gy / %	46.07	39.8	-13.61	a	**4.30E-03**
V40Gy / %	18	13.86	-23.00	b	**6.10E-04**
Lung contralateral					
Dmean / Gy	9.32	8.84	-5.15	a	0.121498134
V5Gy / %	57.57	54.53	-5.28	a	0.082958541
V10Gy / %	38.19	37.83	-0.94	a	0.814826393
V20Gy / %	12.43	11.08	-10.86	b	0.761535645
V40Gy / %	0.82	0.7	-14.63	b	0.446059549
Heart					
D2% / Gy	45.05	43.01	-4.53	b	0.055358887
V20 Gy / %	9.27	7.83	-15.53	a	0.116039212
V45 Gy / %	2.83	2.47	-12.72	a	**0.008692475**
V60 Gy/ %	0.51	0.47	-7.84	b	0.575636914
Dmean / Gy	7.17	6.53	-8.93	a	**0.034038065**
Left ventricle					
D2% / Gy	6.44	5.98	-7.14	a	0.271960361
V10 Gy / %	1.33	1.95	46.62	b	0.400236144
V20 Gy / %	0	0.01		b	0.317310508
V30 Gy / %	0	0			
V40 Gy / %	0	0			
Dmean / Gy	2.42	2.23	-7.85	b	**0.012451172**
Right ventricle					
D2% / Gy	8.14	6.54	-19.66	b	0.157811369
V10 Gy / %	5.79	3.51	-39.38	b	0.10880943
V20 Gy / %	0.48	0.71	47.92	b	1
V30 Gy / %	0	0			
V40 Gy / %	0	0			
Dmean / Gy	2.96	2.49	-15.88	b	0.107758046
Left atrium					
D2% / Gy	52.69	51.76	-1.77	b	0.638671875
V10 Gy / %	45.25	43.41	-4.07	a	0.528540526
V20 Gy / %	24.84	26.64	7.25	a	0.438117752

V30 Gy / %	16.85	17.82	5.76	a	0.548408513
V40 Gy / %	11.57	12.35	6.74	a	0.505890251
Dmean / Gy	15.36	15.51	0.98	a	0.86526113
Right atrium					
D2% / Gy	39.85	32.89	-17.47	a	**0.005337672**
V10 Gy / %	28.24	25.53	-9.60	b	0.177114041
V20 Gy / %	16.79	12.99	-22.63	b	**0.033046944**
V30 Gy / %	9.56	6.49	-32.11	b	**0.003701749**
V40 Gy / %	4.51	2.93	-35.03	b	**0.00146885**
Dmean / Gy	10.09	8.55	-15.26	b	**0.008361816**
Left coronary artery					
D2% / Gy	13.92	11.53	-17.17	a	0.141633743
V5 Gy / %	88.7	86.44	-2.55	b	0.465208819
V10 Gy / %	42.55	38.36	-9.85	b	0.858863013
V20 Gy / %	6.01	5.47	-8.99	b	0.500184257
V30 Gy / %	0.27	0.03	-88.89	b	0.317310508
V40 Gy / %	0	0			
Dmean / Gy	10.11	9.18	-9.20	a	0.464578818
Left circumflex coronary					
D2% / Gy	11.19	12.6	12.60	a	0.345286999
V5 Gy / %	41.4	36.02	-13.00	a	0.063961121
V10 Gy / %	13.33	14.19	6.45	b	0.656641934
V20 Gy / %	0.13	3.45	2553.85	b	0.115851498
V30 Gy / %	0	0			
V40 Gy / %	0	0			
Dmean / Gy	5.21	5.28	1.34	b	0.638671875
Left anterior descending					
D2% / Gy	11.3	9.95	-11.95	a	0.328278434
V5 Gy / %	39.73	26.45	-33.43	a	**0.020891279**
V10 Gy / %	7.82	5.81	-25.70	b	
V20 Gy / %	0.18	0.43	138.89	b	0.715000655
V30 Gy / %	0	0			
V40 Gy / %	0	0			
Dmean / Gy	4.57	3.89	-14.88	a	0.146398018
Right coronary artery					
D2% / Gy	8.49	7.56	-10.95	b	0.846923828
V5 Gy / %	20.01	21.37	6.80	b	0.38627072
V10 Gy / %	8.77	7.44	-15.17	b	0.345231072
V20 Gy / %	2.59	3.47	33.98	b	1
V30 Gy / %	0	0			
V40 Gy / %	0	0			
Dmean / Gy	3.94	3.73	-5.33	b	0.30279541
Posterior descending artery					

D2% / Gy	1.93	1.97	2.07	b	0.078054604
V5 Gy / %	2.48	6.57	164.92	b	0.317310508
V10 Gy / %	0	1.24		b	0.317310508
V20 Gy / %	0	0			
V30 Gy / %	0	0			
V40 Gy / %	0	0			
Dmean / Gy	1.36	1.36	0.00	b	0.082200168

Supplementary Table 4: Dosimetric outcomes of planning in deep inspiration breath-hold (DIBH) and free breathing (FB) for the patients with right lung tumors.

LS 7 positive (n = 22)	FB	DIBH	Mean relative difference (%)	Test	p-value
Lung total					
Dmean (Gy)	16.11	14.52	-9.87	a	**0.00020377**
V20Gy (%)	31.13	27.62	-11.28	a	**0.001696284**
V40Gy / %	8.77	6.94	-20.87	a	**2.98E-04**
Lung ipsilateral					
Dmean / Gy	21.27	18.89	-11.19	a	**1.95E-04**
V5Gy / %	73.92	69.72	-5.68	a	**7.30E-04**
V10Gy / %	64.44	60.16	-6.64	a	**1.95E-03**
V20Gy / %	46.77	41.28	-11.74	a	**9.25E-04**
V40Gy / %	17.06	13.36	-21.69	a	**4.77E-07**
Lung contralateral					
Dmean / Gy	11.64	10.83	-6.96	a	**0.003460223**
V5Gy / %	68.14	62.86	-7.75	a	**0.000397692**
V10Gy / %	49.53	46.47	-6.18	a	**0.018867291**
V20Gy / %	17.53	15.9	-9.30	b	0.236556235
V40Gy / %	1.61	1.57	-2.48	b	0.849795819
Heart					
D2% / Gy	48.45	46.93	-3.14	b	0.053652394
V20 Gy / %	10.13	9.29	-8.29	a	0.193778284
V45 Gy / %	3.08	2.72	-11.69	b	**0.01163652**
V60 Gy/ %	0.53	0.51	-3.77	b	0.647236743
Dmean / Gy	8.02	7.69	-4.11	a	0.140922252
Left ventricle					
D2% / Gy	13.51	11.01	-18.50	b	**0.017245169**
V10 Gy / %	10.34	8.82	-14.70	b	0.306526583
V20 Gy / %	2.62	1.53	-41.60	b	0.123025194
V30 Gy / %	0.85	0.26	-69.41	b	0.115851498
V40 Gy / %	0.31	0.03	-90.32	b	0.067889155
Dmean / Gy	4.42	3.94	-10.86	b	**0.036495955**
Right ventricle					
D2% / Gy	7.9	7.34	-7.09	b	0.726183891
V10 Gy / %	4.33	2.98	-31.18	b	0.414519853
V20 Gy / %	0.21	0.48	128.57	b	0.654720846
V30 Gy / %	0	0			
V40 Gy / %	0	0			
Dmean / Gy	3.04	2.93	-3.62	b	0.489442076
Left atrium					
D2% / Gy	55.96	56.25	0.52	b	0.137489319
V10 Gy / %	55.37	55.3	-0.13	a	0.980078445

V20 Gy / %	33.05	33.88	2.51	a	0.69185051
V30 Gy / %	22.55	23.25	3.10	a	0.655909616
V40 Gy / %	15.54	16.24	4.50	a	0.559713698
Dmean / Gy	18.6	18.78	0.97	a	0.822724616
Right atrium					
D2% / Gy	29.91	27.04	-9.60	b	0.05007267
V10 Gy / %	20.05	20.35	1.50	b	0.77452898
V20 Gy / %	8.58	8	-6.76	b	0.687373616
V30 Gy / %	4.38	3.6	-17.81	b	0.055179277
V40 Gy / %	2.22	1.54	-30.63	b	**0.003494192**
Dmean / Gy	7.7	7.35	-4.55	b	0.176191807
Left coronary artery					
D2% / Gy	17.09	14.62	-14.45	b	0.112871647
V5 Gy / %	95.68	97.33	1.72	b	0.715000655
V10 Gy / %	61.03	53.84	-11.78	b	0.349101945
V20 Gy / %	16.64	10	-39.90	b	0.260392944
V30 Gy / %	4.7	3.96	-15.74	b	0.179712495
V40 Gy / %	0	0			
Dmean / Gy	13.07	11.7	-10.48	b	0.1465168
Left circumflex coronary					
D2% / Gy	23.72	22.77	-4.01	b	0.424488545
V5 Gy / %	56.94	51.21	-10.06	b	**0.02490692**
V10 Gy / %	33.24	30	-9.75	b	0.414040003
V20 Gy / %	11.49	12.71	10.62	b	0.826048864
V30 Gy / %	8.08	7.45	-7.80	b	0.345231072
V40 Gy / %	5.76	4.47	-22.40	b	0.138010738
Dmean / Gy	10.51	9.75	-7.23	b	**0.03587532**
Left anterior descending					
D2% / Gy	16.3	13.67	-16.13	b	**0.007443428**
V5 Gy / %	54.3	44.05	-18.88	a	**0.005592619**
V10 Gy / %	23.5	19.33	-17.74	b	0.054484128
V20 Gy / %	5.22	3.54	-32.18	b	0.138640634
V30 Gy / %	2.03	1.01	-50.25	b	0.317310508
V40 Gy / %	0.6	0.09	-85.00	b	0.317310508
Dmean / Gy	7.14	6.09	-14.71	b	**0.015698997**
Right coronary artery					
D2% / Gy	7.56	7.32	-3.17	b	0.923965931
V5 Gy / %	22.55	23.99	6.39	b	0.569494278
V10 Gy / %	5.53	5.46	-1.27	b	0.916511908
V20 Gy / %	0.51	2.37	364.71	b	0.654720846
V30 Gy / %	0	0			
V40 Gy / %	0	0			

Dmean / Gy	3.67	6.71	82.83	b	0.482591152
Posterior descending artery					
D2% / Gy	2.6	2.46	-5.38	b	0.054245472
V5 Gy / %	7.74	10.75	38.89	b	0.465208819
V10 Gy / %	0	0.85		b	0.317310508
V20 Gy / %	0	0			
V30 Gy / %	0	0			
V40 Gy / %	0	0			
Dmean / Gy	1.87	1.87	0.00	b	**0.039833215**

Supplementary Table 5: Dosimetric outcomes of planning in deep inspiration breath-hold (DIBH) and free breathing (FB) for the patients with involvement of lymph node station 7.

LS 7 negative (n = 10)	FB	DIBH	Mean relative difference (%)	Test	p-value
Lung total					
Dmean (Gy)	10.93	10.08	-7.78	a	0.055415081
V20Gy (%)	18.81	17.22	-8.45	a	0.11702495
V40Gy / %	7.59	6.09	-19.76	a	**2.23E-02**
Lung ipsilateral					
Dmean / Gy	18.66	16.38	-12.22	a	**2.38E-02**
V5Gy / %	59.68	55.45	-7.09	a	5.78E-02
V10Gy / %	51.39	46.16	-10.18	a	**2.71E-02**
V20Gy / %	40.35	35.38	-12.32	a	**2.21E-02**
V40Gy / %	17.63	13.89	-21.21	a	**2.08E-02**
Lung contralateral					
Dmean / Gy	5.17	5.25	1.55	a	0.678309742
V5Gy / %	36.58	36.32	-0.71	a	0.855865983
V10Gy / %	16.71	18.47	10.53	a	0.189344033
V20Gy / %	2.69	3.44	27.88	b	0.235725651
V40Gy / %	0.21	0.25	19.05	b	0.179712495
Heart					
D2% / Gy	26.74	24.47	-8.49	b	0.16015625
V20 Gy / %	6.74	5.65	-16.17	b	0.176296374
V45 Gy / %	1.31	1.14	-12.98	b	0.345447531
V60 Gy/ %	0.11	0.1	-9.09	b	0.705456986
Dmean / Gy	5.61	5.19	-7.49	b	0.10546875
Left ventricle					
D2% / Gy	9.89	9.47	-4.25	b	0.431640625
V10 Gy / %	11.42	10.45	-8.49	b	1
V20 Gy / %	2.27	3.34	47.14	b	0.654720846
V30 Gy / %	0.56	1.06	89.29	b	0.654720846
V40 Gy / %	0.28	0.53	89.29	b	0.317310508
Dmean / Gy	3.54	3.37	-4.80	b	0.10546875
Right ventricle					
D2% / Gy	6.41	5.23	-18.41	b	0.173070921
V10 Gy / %	2.41	1.93	-19.92	b	1
V20 Gy / %	0.25	0	-100.00	b	0.317310508
V30 Gy / %	0	0			
V40 Gy / %	0	0			
Dmean / Gy	2.54	2.17	-14.57	b	0.26261829
Left atrium					
D2% / Gy	31.18	28.07	-9.97	b	0.845703125
V10 Gy / %	31.88	31.33	-1.73	b	0.888637861

V20 Gy / %	16.27	17.49	7.50	b	0.865772375
V30 Gy / %	8.66	9.19	6.12	b	0.685830435
V40 Gy / %	5.03	5.25	4.37	b	0.892738401
Dmean / Gy	10.1	10.21	1.09	b	1
Right atrium					
D2% / Gy	19.93	16.81	-15.65	b	0.514669723
V10 Gy / %	14.98	11.63	-22.36	b	0.224915884
V20 Gy / %	8.64	5.45	-36.92	b	0.248863875
V30 Gy / %	5.35	2.55	-52.34	b	0.138010738
V40 Gy / %	1.94	1.07	-44.85	b	0.10880943
Dmean / Gy	6.05	4.86	-19.67	b	0.440867035
Left coronary artery					
D2% / Gy	11.12	9.61	-13.58	a	**0.014399122**
V5 Gy / %	70	54.7	-21.86	b	0.179712495
V10 Gy / %	30.39	31.31	3.03	b	0.592980098
V20 Gy / %	3.48	6.75	93.97	b	0.654720846
V30 Gy / %	0	0			
V40 Gy / %	0	0			
Dmean / Gy	8.47	7.9	-6.73	a	0.298325803
Left circumflex coronary					
D2% / Gy	15.31	12.09	-21.03	b	0.085830958
V5 Gy / %	32.59	26.7	-18.07	b	**0.027707849**
V10 Gy / %	20.01	20.25	1.20	b	0.892738401
V20 Gy / %	11.45	9.09	-20.61	b	0.654720846
V30 Gy / %	7.02	5.46	-22.22	b	0.654720846
V40 Gy / %	4.04	3.68	-8.91	b	0.654720846
Dmean / Gy	7.61	6.85	-9.99	b	0.16015625
Left anterior descending					
D2% / Gy	12.81	9.34	-27.09	a	0.234049702
V5 Gy / %	39.51	29.15	-26.22	b	0.128190174
V10 Gy / %	15.44	13.61	-11.85	b	
V20 Gy / %	2.23	2.08	-6.73	b	0.715000655
V30 Gy / %	0.42	0	-100.00	b	0.317310508
V40 Gy / %	0.15	0	-100.00	b	0.317310508
Dmean / Gy	5.33	4.5	-15.57	b	0.138640634
Right coronary artery					
D2% / Gy	6.29	5.88	-6.52	b	0.888637861
V5 Gy / %	11.35	12.01	5.81	b	0.685830435
V10 Gy / %	4.49	3	-33.18	b	0.285049407
V20 Gy / %	2.75	0	-100.00	b	0.317310508
V30 Gy / %	0	0			
V40 Gy / %	0	0			

Dmean / Gy	2.8	2.29	-18.21	b	0.575403023
Posterior descending artery					
D2% / Gy	1.33	1.26	-5.26	b	0.498962299
V5 Gy / %	0	0			
V10 Gy / %	0	0			
V20 Gy / %	0	0			
V30 Gy / %	0	0			
V40 Gy / %	0	0			
Dmean / Gy	1.08	1	-7.41	b	0.39802472

Supplementary Table 6: Dosimetric outcomes of planning in deep inspiration breath-hold (DIBH) and free breathing (FB) for the patients without involvement of lymph node station 7.

Danksagung

An dieser Stelle möchte ich Alldenjenigen meinen Dank aussprechen, ohne die diese Dissertation nicht möglich gewesen wäre:

Ein besonderer Dank gilt meinem Doktorvater, Herrn Prof. Dr. med. Michael J. Eble, für die Ermöglichung eines so klinisch relevanten Themas – für die Betreuung, Unterstützung und Supervision während der gesamten Promotion.

Besonders möchte ich meinem betreuenden Oberarzt, Herrn PD Dr. med. Ahmed Allam Mohamed, danken, mit dessen Rat, Korrekturen und Wegweisung das Projekt umgesetzt werden konnte.

Danken möchte ich zudem meinem Kollegen, Herrn MUDr. Ahmed Ali Chughtai, für seine vielen Hilfestellungen.

Ein weiterer Dank geht an Frau Jennifer Heinrichs für die physikalischen Planung, wie auch an Herrn Prof. Dr. med. Philipp Bruners für die radiologische Begleitung der Arbeit.

Ganz besonders möchte ich meinen Eltern und meinem Bruder danken, dich mich auf meinem gesamten Weg bedingungslos unterstützt haben.

Erklärung § 5 Abs. 1 zur Datenaufbewahrung

Hiermit erkläre ich, dass die dieser Dissertation zu Grunde liegenden Originaldaten in der ***Klinik für Radioonkologie und Strahlentherapie*** des Universitätsklinikums Aachen hinterlegt sind.

Muster 5

Erklärung gemäß § 5 Abs. (1) und (2), und § 11 Abs. (3) 12. der Promotionsordnung

Hiermit erkläre ich, **Biney Pal Singh,** an Eides statt, dass ich den wesentlichen Anteil an der Publikation:
Biney Pal Singh, Ahmed Ali Chughtai, Jennifer Stock, Philipp Bruners, Michael J. Eble and Ahmed Allam Mohamed**:** Comparative dosimetric analysis of deep inspiration breath-hold versus free breathing radiotherapy in lung cancer: special emphasis on cardiac and subcardiac structure protection; Technical Innovations & Patient Support in Radiation Oncology (tipsRO).; Volume 37, 100371, März 2026

Die Anteile an der Arbeit waren wie folgt:

Bitte Namen angeben ----------->	Doktor-and, Biney Pal Singh	Facharzt, Herr Dr. Ahmed Ali Chughtai	Medizinphysikexpertin, Frau Jennifer Heinrichs (geb. Stock)	Radiologe, Herr Prof. Dr. Philipp Bruners	Betreuender Oberarzt, Herr PD Dr. Ahmed Allam Mohamed	Doktorvater, Herr Prof. Dr. Michael J. Eble	Arbeiten aus früheren Prüfungen gelten als Fremdleistung *	Summe (%)
Studienüberwachung					30	70		100
Studiendesign/Konzeption	60	20				20		100
Erstellung Segmentation medizinisch	70	30						100
Revision Segmentation				50	40	10		100
Erstellung Bestrahlungspläne physikalisch			100					100
Datenerhebung	100							100
Datenauswertung	100							100
Statistische Auswertung	70				30			100
Interpretation der Datenauswertung	60				30	10		100
Revision Auswertung					50	50		100
Verfassung des Manuskripts	**100**							100
Korrektur des Manuskripts					50	50		100

Aus diesem wesentlichen Anteil ergibt sich selbstverständlich die Stellung als Erstautor.

_____
Unterschrift der Doktorandin/des Doktoranden

Als Doktorvater und / oder korresp. Autor bestätige ich die Angaben von Biney Pal Singh und in Vertretung für die nicht mehr erreichbaren und **nicht deutschsprachigen** Koautoren.

_____ _____
Unterschrift des Doktorvaters Unterschrift des korresp. Autors (falls abweichend)

Ich schließe mich der Erklärung von Herrn Prof. Dr. med Michael J. Eble als Koautor an

PD Dr. med Ahmed Allam Mohamed

Namen und Unterschriften aller deutschsprachigen Koautoren

Hiermit erkläre ich, **Biney Pal Singh,** an Eides statt, dass ich den wesentlichen Anteil an der Publikation:
Biney Pal Singh, Ahmed Ali Chughtai, Jennifer Stock, Philipp Bruners, Michael J. Eble and Ahmed Allam Mohamed**:** Comparative dosimetric analysis of deep inspiration breath-hold versus free breathing radiotherapy in lung cancer: special emphasis on cardiac and subcardiac structure protection; Technical Innovations & Patient Support in Radiation Oncology (tipsRO).; 2026, Volume 37, 100371, März 2026

Die Anteile an der Arbeit waren wie folgt:

**Bitte Namen angeben ----------> **	Doktor-and, Biney Pal Singh	Facharzt, Herr Dr. Ahmed Ali Chughtai	Medizinphysikexpertin, Frau Jennifer Heinrichs (geb. Stock)	Radiologe, Herr Prof. Dr. Philipp Bruners	Betreuender Oberarzt, Herr PD Dr. Ahmed Allam Mohamed	Doktorvater, Herr Prof. Dr. Michael J. Eble	Arbeiten aus früheren Prüfungen gelten als Fremdleistung *	Summe (%)
Studienüberwachung					30	70		100
Studiendesign/Konzeption	60	20				20		100
Erstellung Segmentation medizinisch	70	30						100
Revision Segmentation				50	40	10		100
Erstellung Bestrahlungspläne physikalisch			100					100
Datenerhebung	100							100
Datenauswertung	100							100
Statistische Auswertung	70				30			100
Interpretation der Datenauswertung	60				30	10		100
Revision Auswertung					50	50		100
Verfassung des Manuskripts	**100**							100
Korrektur des Manuskripts					50	50		100

Ich schließe mich der Erklärung von Herrn Prof. Dr. med Michael J. Eble als Koautor an

MUDr. Ahmed Ali Chughtai, PhD, MHBA

Namen und Unterschriften aller deutschsprachigen Koautoren

Jennifer Heinrichs, geb. Stock, MSc

Namen und Unterschriften aller deutschsprachigen Koautoren

Prof. Dr. med Philipp Bruners

Namen und Unterschriften aller deutschsprachigen Koautoren

BINEY PAL SINGH

PERSÖNLICHE DATEN

Geburtsdatum	27.02.1995
Geburtsort	Solingen
Familienstand	ledig
Nationalität	Deutsch

SCHULAUSBILDUNG

08/2005 – 07/2013	Abitur, *Gymnasium Schwertstraße Solingen*

STUDIUM

10/2013 – 11/2019	Studium der Humanmedizin, *RWTH Aachen*
09/2016	Ärztliche Basisprüfung (Physikumsäquivalent der RWTH Aachen)
10/2018	2. Staatsexamen
11/2019	3. Staatsexamen

FAMULATUREN

18.09. – 05.10.2015	Kardiologie, *Städtisches Klinikum Solingen*
03.03. – 01.04.2017	Nephrologie, *Städtisches Klinikum Solingen*
17.11. – 01.12.2017	Augenheilkunde, *Uniklinik RWTH Aachen*
05.02. – 06.03.2018	Allgemeinmedizin, *Hausarztpraxis Dr. Simadirakis, Solingen*
07.03. – 21.03.2018	Strahlentherapie, *Uniklinik RWTH Aachen*
22.03. – 06.04.2018	Neuroradiologie, *Uniklinik RWTH Aachen*

PRAKTISCHES JAHR

11/2018 – 03/2019	Innere Medizin, *Städtisches Klinikum Solingen*
03/2019 – 06/2019	Radioonkologie und Strahlentherapie, *Uniklinik RWTH Aachen*
07/2019 – 10/2019	Chirurgie, *Städtisches Klinikum Solingen*

BERUFLLICHE ERFAHRUNG

01/2020 – 01/2026	Assistenzarzt in der Klinik für Radioonkologie und Strahlentherapie, *Uniklinik RWTH Aachen* Direktor: Prof. Michael J. Eble
Seit 19.01.2026	Facharzt für Strahlentherapie, Klinik für Radioonkologie und Strahlentherapie, *Uniklinik RWTH Aachen* Direktor: Prof. Michael J. Eble

SONSTIGE TÄTIGKEITEN

01/2016 – 03/2017	Studentische Aushilfskraft Pflege, *Städtisches Klinikum Solingen*
Seit 10/2014	Leitung von Physik-Brückenkursen des Modellstudiengangs Aachen

ZUSATZQUALIFIKATIONEN

Seit 04/2015	Diverse Kurse und Veranstaltungen in der Radiologischen Bildgebung und Medizintechnik im Rahmen des Studiums
10/2020	Kombinierter Grund- und Kenntniskurs Strahlenschutz - Nuklearmedizin, Strahlentherapie, Röntgen
02/2021	GCP-Basiskurs, AMG-Prüferkurs, MPG-Prüferkurs, *Center for Translational & Clinical Research Aachen (CTC-A) Uniklinik RWTH Aachen*
01/2026	Fachkunde im Strahlenschutz – Gesamtgebiet der Strahlenbehandlung und In der Röntgendiagnostik für den Bereich Strahlentherapieplanung

FORTBILDUNGEN

06/2022	Artificial Intelligence for Imaging, *Maastricht University*
09/2022	ESTRO Basic Clinical Radiobiology
02/2023	ESTRO Particle Therapy
03/2023	Medizinische Physik und Technik für Radioonkolog*innen, *Universität Heidelberg*
05/2023	ESTRO Interdisciplinary - SBRT/SRS: a one-day bootcamp
09/2023	ESTRO Target Volume Determination
03/2024	ESTRO Immunotherapy
12/2025	ESTRO Pediatric radiotherapy

PUBLIKATIONEN

Mohamed AA, Goncalves M, **Singh BP**, Tometten M, Rashad A, Hölzle F, Hackenberg S, Eble M. Stereotactic radiotherapy in the management of oligometastatic and recurrent head and neck cancer: a single-center experience. Strahlenther Onkol. 2024 May;200(5):400-408. doi: 10.1007/s00066-023-02180-9. Epub 2023 Dec 8. PMID: 38063900; PMCID: PMC11039526.

SONSTIGES

Sprachkenntnisse	Panjabi, *Muttersprache*
	Deutsch, verhandlungssicher
	Englisch, *sehr gute Fremdsprachenkenntnisse*
	Spanisch, *Grundkenntnisse*
	Japanisch, Grundkenntnisse
Hobbys	Malen und Zeichnen, Fotographie
	Wandern
	Hockey
	Fußball

Aachen, 13. August 2026