σ -Galois Theory of Linear Difference Equations

Alexey Ovchinnikov^{1,2} and Michael Wibmer³

¹Department of Mathematics, CUNY Queens College, 65-30 Kissena Blvd, Queens, NY 11367-1597, USA, ²CUNY Graduate Center, Department of Mathematics, 365 Fifth Avenue, New York, NY 10016, USA, and

³RWTH Aachen, 52056 Aachen, Germany

Correspondence to be sent to: e-mail: aovchinnikov@qc.cuny.edu

We develop a Galois theory for systems of linear difference equations with an action of an endomorphism σ . This provides a technique to test whether solutions of such systems satisfy σ -polynomial equations and, if yes, then characterize those. We also show how to apply our work to study isomonodromic difference equations and difference algebraic properties of meromorphic functions.

1 Introduction

Inspired by the numerous applications of the differential algebraic independence results from [36], we develop a Galois theory with an action of an endomorphism σ for systems of linear difference equations of the form $\phi(y) = Ay$, where $A \in \mathrm{GL}_n(K)$ and K is a $\phi\sigma$ -field, that is, a field with two given commuting endomorphisms ϕ and σ , such as in Example 2.1. This provides a technique to test whether solutions of such systems satisfy σ -polynomial equations and, if yes, then characterize those. Galois groups, in this approach, are groups of invertible matrices defined by σ -polynomial equations with coefficients in the σ -field $K^\phi := \{a \in K \mid \phi(a) = a\}$. In more technical terms, such groups are functors from K^ϕ - σ -algebras to sets represented by finitely σ -generated K^ϕ - σ -Hopf

Received October 14, 2013; Revised March 6, 2014; Accepted March 18, 2014
Communicated by Prof. Dmitry Kaledin

algebras [23]. Also, our work is a highly nontrivial generalization of [5], where similar problems were considered but σ was required to be of finite order (there exists n such that $\sigma^n = \text{id}$).

Our main result is a construction of a σ -Picard–Vessiot (σ -PV) extension (see Theorem 2.28), that is, a minimal $\phi\sigma$ -extension of the base $\phi\sigma$ -field K containing solutions of $\phi(y) = Ay$. It turns out that the standard constructions and proofs in the previously existing difference Galois theories do not work in our setting. Indeed, this is mainly due to the reason that even if the field K^ϕ is σ -closed [56], consistent systems of σ -equations (such that the equation $1 = 0$ is not a σ -algebraic consequence of the system) with coefficients in K^ϕ might not have a solution with coordinates in K^ϕ (see more details in Remarks 2.19 and 2.22). However, our method avoids this issue. In our approach, a σ -PV extension is built iteratively (applying σ), by carefully choosing a suitable usual PV extension [58] at each step, and then “patching” them together. This is a difficult problem and requires several preparatory steps as described in Section 2.4. A similar approach was also taken in [62, Theorem 8] for systems of differential equations with parameters. However, our case is more subtle and, as a result, requires more work. A Tannakian approach is taken in [49] to build a foundation that will allow several endomorphisms by considering semigroup actions on Tannakian categories.

Galois theory of difference equations $\phi(y) = Ay$ without the action of σ was studied in [1–4, 13, 58, 63], with a nonlinear generalization considered in [31, 44], as well as with an action of a derivation ∂ in [17–22, 33, 34, 36]. The latter works provide algebraic methods to test whether solutions of difference equations satisfy polynomial differential equations (see also [40] for a general Tannakian approach). In particular, these methods can be used to prove Hölder’s theorem that states that the Γ -function, which satisfies the difference equation $\Gamma(x+1) = x \cdot \Gamma(x)$, satisfies no nontrivial differential equation over $\mathbb{C}(x)$. A Galois theory of differential equations $\partial(y) = Ay$ (the matrix A does not have to be invertible in this case) with an action of σ was also developed in [23].

Our work has numerous applications to studying difference and differential algebraic properties of functions. Isomonodromic q -difference equations, which lead to q -difference Painlevé equations, have been recently studied in [37–39, 45]. In Theorem 2.55, we show how this property can be detected using our σ -PV theory, which can be combined with [48, Theorem 4.1] to study difference isomonodromy with several parameters. On the other hand, Theorem 3.1 gives a general σ -algebraic independence (called difference hypertranscendence in [55]) test for first-order ϕ -difference equations. Theorem 3.5 translates this to a σ -algebraic dependence test over the field of meromorphic functions with Nevanlinna growth order < 1 (see (3.6)). It turns out that our

methods allow us to generalize a modification (Lemma 3.4) of complex-analytic results from [6], which is another interesting application. Theorem 3.6 combined with either Theorems 3.1 or 3.5 can be used as computational tool. We illustrate this in Examples 3.7 and 3.8 as well as show how our work could be used to study differential algebraic properties of functions given by power series in Example 3.9. Not only do we show practical applications of our work, we also hope that our theory will be applied in the future in diverse areas, such as described in [15, 52] and the papers on isomonodromic q -difference equations mentioned above.

The paper is organized as follows. We start with the basic definitions, notation, and review of existing results in Sections 2.1 and 2.2. We then introduce σ -PV extensions and study their basic properties in Section 2.3. The main result, existence of σ -PV extensions, is contained in Section 2.4, which starts by developing the needed technical tools. We extend the main result in Section 2.5 to include more useful situations in which σ -PV extensions exist. Uniqueness for σ -PV extensions is established in Section 2.6. We recall from the appendix of [23] what difference algebraic groups are, establish the σ -Galois correspondence, and show that the σ -dimension of the σ -Galois group coincides with the σ -dimension of the σ -PV extension in Section 2.7. The relation between isomonodromic difference equations and our Galois theory is given in Section 2.8. Applications to difference and differential algebraic properties of functions, including functions with a slow Nevanlinna growth order, and illustrative examples are given in Section 3.

2 σ -PV Extensions

2.1 Basic definitions and preliminaries

We need to introduce some terminology from difference algebra. Standard references for difference algebra are [16, 43]. All rings are assumed to be commutative. By a ϕ -ring, we mean a ring R equipped with a ring endomorphism $\phi: R \rightarrow R$. We do not require that ϕ is an automorphism. If ϕ is an automorphism, we say that R is *inversive*. By a $\phi\sigma$ -ring, we mean a ring equipped with two commuting endomorphisms ϕ and σ . A morphism of ϕ -rings (or $\phi\sigma$ -rings) is a morphism of rings that commutes with the endomorphisms. If the underlying ring is a field, we speak of ϕ -fields (or $\phi\sigma$ -fields). Here are some basic examples of $\phi\sigma$ -fields of interest to us:

Example 2.1.

- (i) The $\phi\sigma$ -field M of meromorphic functions on $\mathbb{C}$ with $\phi(f)(z) = f(z + z_\phi)$ and $\sigma(f)(z) = f(z + z_\sigma)$, $f \in M$, $z_\phi, z_\sigma \in \mathbb{C}$ and its $\phi\sigma$ -subfields $\mathbb{C}(z)$ and $M_{<1}$, the

field of meromorphic functions on $\mathbb{C}$ with Nevanlinna growth order < 1 (see Section 3.2).

- (ii) The $\phi\sigma$ -field M with $\phi(f)(z) = f(z \cdot q_\phi)$ and $\sigma(f)(z) = f(z \cdot q_\sigma)$, $f \in M$, $q_\phi, q_\sigma \in \mathbb{C}^\times$ and its subfields $\mathbb{C}(z)$ and $M_{<1}$.
- (iii) The $\phi\sigma$ -field $\mathbb{C}(z, w)$ with $\phi(f)(z, w) = f(z + z_\phi, w)$ and $\sigma(f)(z, w) = f(z, w + w_\sigma)$, $f \in \mathbb{C}(z, w)$, $z_\phi, w_\sigma \in \mathbb{C}$ and various other actions of ϕ and σ that commute. $\square$

Recall that:

- (1) A ϕ -ideal in a ϕ -ring R is an ideal $\mathfrak{a}$ of R such that $\phi(\mathfrak{a}) \subset \mathfrak{a}$. Similarly, one defines $\phi\sigma$ -ideals in $\phi\sigma$ -rings.
- (2) A ϕ -ring is called ϕ -simple if the zero ideal and the whole ring are the only ϕ -ideals.
- (3) A ϕ -ideal $\mathfrak{q}$ in a ϕ -ring R is called ϕ -prime if $\mathfrak{q}$ is a prime ideal of R and $\phi^{-1}(\mathfrak{q}) = \mathfrak{q}$.
- (4) If ϕ is an endomorphism of a ring R , then ϕ^d is also an endomorphism of R for every $d \geq 1$, and we can speak of ϕ^d -prime ideals of R .
- (5) A ϕ -ring R is called a ϕ -domain if its zero ideal is ϕ -prime. (Equivalently, R is an integral domain and $\phi: R \rightarrow R$ is injective.)
- (6) A ϕ -ideal in a ϕ -ring R is called ϕ -maximal if it is a maximal element in the set of all ϕ -ideals of R , not equal to R , ordered by inclusion.

The theory of difference fields does exhibit some pathologies. For example, two extensions of the same difference field can be incompatible, see [43, Chapter 5]. As it has been recognized in [58], the Galois theory of linear difference equations runs much smoother if one allows certain finite products of fields instead of fields. In this context, the following definition has turned out to be useful.

Definition 2.2. A ϕ -pseudo field is a ϕ -simple, Noetherian ϕ -ring K such that every nonzero divisor of K is invertible in K . $\square$

The concept of ϕ -pseudo fields (in certain variants) is also used in [2, 5, 36, 56, 57, 60, 61].

If K is a ϕ -pseudo field, then there exist orthogonal, idempotent elements $e_1, \dots, e_d$ of K such that

- (1) $K = e_1 \cdot K \oplus \dots \oplus e_d \cdot K$;

- (2) $\phi(e_1) = e_2, \phi(e_2) = e_3, \dots, \phi(e_d) = e_1$;
- (3) $e_i \cdot K$ is a field for $i = 1, \dots, d$ (so, $e_i \cdot K$ is a ϕ^d -field)

(see, e.g., [60, Proposition 1.3.2, p. 9]). The integer d is called the *period* of K and denoted by $\text{period}(K)$.

Definition 2.3. A ϕ -ideal $\mathfrak{p}$ of a ϕ -ring R is called *ϕ -pseudo prime* if it is the kernel of a morphism from R into some ϕ -pseudo field. Equivalently, $\mathfrak{p}$ is of the form

$$\mathfrak{p} = \mathfrak{q} \cap \phi^{-1}(\mathfrak{q}) \cap \dots \cap \phi^{-(d-1)}(\mathfrak{q}) \quad (2.1)$$

for some ϕ^d -prime ideal $\mathfrak{q}$ of R . The smallest number d such that (2.1) holds for some ϕ^d -prime ideal $\mathfrak{q}$ of R is called the period of $\mathfrak{p}$. $\square$

Definition 2.4. By a *ϕ -pseudo domain*, we mean a ϕ -ring whose zero ideal is ϕ -pseudo prime. If R is a ϕ -pseudo domain, the period of the zero ideal of R is also called the period of R . $\square$

Note that every ϕ -subring of a ϕ -pseudo field is a ϕ -pseudo domain. The total ring of fractions of a ϕ -pseudo domain is a ϕ -ring in a natural way, indeed it is a ϕ -pseudo field.

Definition 2.5. A $\phi\sigma$ -ring R is called a *ϕ -pseudo σ -domain* if (R, ϕ) is a ϕ -pseudo domain. $\square$

Definition 2.6. A $\phi\sigma$ -ring K is called a *ϕ -pseudo σ -field* if (K, ϕ) is a ϕ -pseudo field. $\square$

Most of the employed nomenclature is self-explanatory. For example,

- (1) A K - $\phi\sigma$ -algebra is a K -algebra R equipped with the structure of a $\phi\sigma$ -ring such that the K -algebra structure map $K \rightarrow R$ is a morphism of $\phi\sigma$ -rings.
- (2) Constants are denoted by upper indices. For example, if R is a ϕ -ring, then the ϕ -constants of R are

$$R^\phi := \{r \in R \mid \phi(r) = r\}.$$

If K is a ϕ -pseudo σ -field, then K^ϕ is a σ -field (as R^ϕ is a field for any ϕ -simple ϕ -ring R [58, (Lemma 1.7a), p. 6].)

- (3) If R is a ring, we denote the total quotient ring of R , that is, the localization of R at the multiplicatively closed subset of all nonzero divisors, by $\text{Quot}(R)$.

- (4) If K is a ring, R a K -algebra, and S a subset of R , then $K(S)$ denotes the smallest K -subalgebra of R that contains S and is closed under taking inverses. So, explicitly

$$K(S) = \{a/b \mid a \in K[S], b \in K[S] \cap R^\times\} \subset R.$$

- (5) If K is a σ -ring, R a K - σ -algebra, and S a subset of R , then $K\{S\}_\sigma$ denotes the K - σ -subalgebra of R generated by S , that is, the K -subalgebra of R generated by all elements of the form $\sigma^d(s)$, where $s \in S$ and $d \geq 0$. (By definition, σ^0 is the identity map.) If $R = K\{S\}_\sigma$ with S finite, we say that R is finitely σ -generated over K .

- (6) Let

$$K\langle S \rangle_\sigma := \{a/b \mid a \in K\{S\}_\sigma, b \in K\{S\}_\sigma \cap R^\times\} \subset R. \quad (2.2)$$

If $L|K$ is an extension of σ -pseudo fields, we say that L is finitely σ -generated over K if there exists a finite subset S of L such that $K\langle S \rangle_\sigma = L$.

- (7) Tensor products of difference rings are considered as difference rings in a natural fashion. For example, if R is a ϕ -ring and S, T are R - ϕ -algebras, then $S \otimes_R T$ becomes an R - ϕ -algebra by setting $\phi(s \otimes t) = \phi(s) \otimes \phi(t)$.

Finally, we record some simple and well-known lemmas that we use repeatedly throughout the text.

Lemma 2.7 ([60, Lemma 1.1.5, p. 4]). Let R be a ϕ -simple ϕ -ring. Then, $\text{Quot}(R)^\phi = R^\phi$. □

Lemma 2.8. Let R be a ϕ -simple ϕ -ring, and D a R^ϕ -algebra (considered as constant ϕ -ring). The map $\mathfrak{b} \mapsto R \otimes_{R^\phi} \mathfrak{b}$ defines a bijection between the set of all ideals in D and the set of all ϕ -ideals in $R \otimes_{R^\phi} D$. The inverse map is given by $\mathfrak{a} \mapsto \mathfrak{a} \cap D$. □

Proof. In [60, Proposition 1.4.15, p. 15], this is stated for the case that R is a ϕ -pseudo field. However, the proof given there only uses the assumption that R is ϕ -simple. ■

Lemma 2.9. Let R be a ϕ -simple ϕ -ring and D a (ϕ -constant) field extension of R^ϕ . Then, $R \otimes_{R^\phi} D$ is ϕ -simple. □

Proof. This is clear from Lemma 2.8. ■

Lemma 2.10 ([60, Lemma 1.1.6, p. 4]). Let K be a ϕ -simple ϕ -ring and R a K - ϕ -algebra. Then, K and R^ϕ are linearly disjoint over K^ϕ . $\square$

Lemma 2.11. Let R be a ϕ -simple $\phi\sigma$ -ring that is a ϕ -pseudo domain. Then, σ is injective on R and the zero ideal of R is the finite intersection of σ -pseudo prime ideals. Moreover, $\text{Quot}(R)$ is naturally a ϕ -pseudo σ -field. $\square$

Proof. Since ϕ and σ commute, the kernel of σ is a ϕ -ideal. Therefore, σ must be injective. Since R is a ϕ -pseudo domain, the zero ideal of R is a finite intersection of prime ideals. As σ is injective, the map $\mathfrak{q} \mapsto \sigma^{-1}(\mathfrak{q})$ is a permutation of the set of minimal prime ideals of R . Every cycle in the cycle decomposition of this permutation corresponds to a σ -pseudo prime ideal. Since R is a finite direct sum of integral domains [60, Proposition 1.1.2, p. 2], it is clear that σ and ϕ extend to $\text{Quot}(R)$. $\blacksquare$

2.2 Review of the classical PV theory

To maximize the applicability of our σ -Galois theory, we have been careful to avoid unnecessary technical conditions on the base field:

- (i) we work in arbitrary characteristic;
- (ii) we do not assume that our endomorphisms are automorphisms;
- (iii) we do not make any initial requirements on the constants.

Unfortunately, the assumptions in the standard presentations of the classical Galois theory of linear difference equations (e.g., [58]) are somewhat more restrictive. Since, at some points in the development of our σ -Galois theory, we need to use the classical Galois theory, we have to give the definitions and recall the results in our slightly more general setup. This review of the classical theory will also help the reader see the analogy between the classical Galois theory and the σ -Galois theory.

Definition 2.12. Let K be a ϕ -pseudo field and $A \in \text{GL}_n(K)$. An extension $L|K$ of ϕ -pseudo fields with $L^\phi = K^\phi$ is called a *PV extension* for $\phi(y) = Ay$ if there exists a matrix $Y \in \text{GL}_n(L)$ such that $\phi(Y) = AY$ and $L = K(Y) := K(Y_{ij} \mid 1 \leq i, j \leq n)$.

A ϕ -simple K - ϕ -algebra R is called a *PV ring* for $\phi(y) = Ay$ if there exists $Y \in \text{GL}_n(R)$ such that $\phi(Y) = AY$ and $R = K[Y, 1/\det(Y)]$. $\square$

It is easy to describe a construction of a PV ring. Indeed, let X be the $n \times n$ -matrix of indeterminates over K . We turn $K[X, 1/\det(X)]$ into a K - ϕ -algebra by setting

$\phi(X) = AX$. Then,

$$K[X, 1/\det(X)]/\mathfrak{m}$$

is a PV ring for $\phi(Y) = AY$ for every ϕ -maximal ϕ -ideal $\mathfrak{m}$ of $K[X, 1/\det(X)]$. Moreover, every PV ring for $\phi(Y) = AY$ is of this form.

The existence of PV extensions is a more delicate issue, unless we assume that K^ϕ is algebraically closed. The problem is that a PV ring might contain new constants. The following lemma guarantees that the constants of a PV ring over K are an algebraic field extension of K^ϕ .

Lemma 2.13. Let K be a ϕ -pseudo field and R a ϕ -simple K - ϕ -algebra which is finitely generated as K -algebra. Then, R^ϕ is an algebraic field extension of K^ϕ . $\square$

Proof. This is a slight generalization of [58, Lemma 1.8, p. 7]. It also follows from [61, Proposition 2.11, p. 1389]. $\blacksquare$

The following proposition explains the intimate relation between PV extensions and PV rings.

Proposition 2.14. Let K be a ϕ -pseudo field and $A \in \mathrm{GL}_n(K)$. Let R be a K - ϕ -algebra that is a ϕ -pseudo domain. Assume that $R = K[Y, 1/\det(Y)]$ for some $Y \in \mathrm{GL}_n(R)$ with $\phi(Y) = AY$. Then, R is ϕ -simple if and only if $\mathrm{Quot}(R)^\phi$ is algebraic over K^ϕ . $\square$

Proof. It is clear from Lemmas 2.13 and 2.7 that $\mathrm{Quot}(R)^\phi$ is algebraic over K^ϕ if R is ϕ -simple. So, we assume that $\mathrm{Quot}(R)^\phi$ is algebraic over K^ϕ . Indeed, we will first assume that $\mathrm{Quot}(R)^\phi = K^\phi$. Let

$$R' = K[Y', 1/\det(Y')]$$

be a PV ring for $\phi(Y) = AY$, where $Y' \in \mathrm{GL}_n(R')$ satisfies $\phi(Y') = AY'$. Note that $L := \mathrm{Quot}(R)$ is a ϕ -pseudo field. The matrix

$$Z := (Y'^{-1} \otimes 1) \cdot (1 \otimes Y') \in \mathrm{GL}_n(L \otimes_K R')$$

satisfies

$$\phi(Z) = ((AY)^{-1} \otimes 1) \cdot (1 \otimes AY') = Z.$$

It follows from Lemma 2.10 that

$$\begin{aligned}
 L \otimes_{K^\phi} K^\phi[Z, 1/\det(Z)] &= L \cdot (K[Z, 1/\det(Z)]) = L[Z, 1/\det(Z)] \\
 &= L[Y \otimes 1, Z, 1/\det(Y) \otimes 1, 1/\det(Z)] \\
 &= L[1 \otimes Y', 1 \otimes 1/\det(Y')] = L \otimes_K R'.
 \end{aligned} \tag{2.3}$$

Since $K^\phi[Z, 1/\det(Z)]$ is finitely generated as K^ϕ -algebra, there exists an algebraic field extension C of K^ϕ and a K^ϕ -morphism

$$\psi: K^\phi[Z, 1/\det(Z)] \rightarrow C.$$

Composing the inclusion $R' \rightarrow L \otimes_K R'$ with (2.3) and $\text{id} \otimes \psi$, we obtain a K - ϕ -morphism

$$R' \rightarrow L \otimes_{K^\phi} C.$$

Since R' is ϕ -simple, we can identify R' with a subring of $L \otimes_{K^\phi} C$. The two solution matrices Y and Y' in $\text{GL}_n(L \otimes_{K^\phi} C)$ only differ by multiplication by an invertible matrix with entries in C . Therefore,

$$R \otimes_{K^\phi} C = K[Y, 1/\det(Y)] \otimes_{K^\phi} C = K[Y, 1/\det(Y), C] = K[Y', 1/\det(Y'), C] = R' \otimes_{R'^\phi} C,$$

by Lemma 2.10 again. From Lemma 2.9, we know that $R' \otimes_{R'^\phi} C$ is ϕ -simple. This implies that R is ϕ -simple, because a nontrivial ϕ -ideal of R would give rise to a nontrivial ϕ -ideal of $R \otimes_{K^\phi} C$.

In the general case, we set

$$\tilde{K} = K \otimes_{K^\phi} L^\phi \subset L.$$

We claim that $\tilde{K}$ is a ϕ -pseudo field. We already know from Lemma 2.9 that $\tilde{K}$ is ϕ -simple and, since L is a ϕ -pseudo domain, $\tilde{K}$ is also a ϕ -pseudo domain. Then, $\tilde{K}$ is a finite direct sum of integral domains R_i [60, Proposition 1.1.2, p. 2]. Since L^ϕ is algebraic over K^ϕ , $\tilde{K}$ is integral over K . As K is a direct sum of fields K_j , this implies that each R_i is integral over some K_j . But, since R_i is an integral domain and K_j a field, R_i must be a field. So, $\tilde{K}$ is a finite direct sum of fields. Consequently, $\tilde{K}$ is a ϕ -pseudo field.

From the first part of the proof, it follows that $\tilde{K}[Y, 1/\det(Y)]$ is ϕ -simple. We have to show that $R = K[Y, 1/\det(Y)]$ is ϕ -simple. Suppose that $\mathfrak{a} \subset R$ is a nontrivial ϕ -ideal of R . Since L^ϕ is algebraic over K^ϕ , $\tilde{K}[Y, 1/\det(Y)]$ is integral over R . Therefore,

the ideal $\mathfrak{a}'$ of $\tilde{K}[Y, 1/\det(Y)]$ generated by $\mathfrak{a}$ does not contain 1 [24, Proposition 4.15, p. 129]. As $\mathfrak{a}'$ is a ϕ -ideal, this yields a contradiction. ■

Corollary 2.15. Let K be a ϕ -pseudo field and $A \in \mathrm{GL}_n(K)$. If $L|K$ is a PV extension for $\phi(y) = Ay$ with fundamental solution matrix $Y \in \mathrm{GL}_n(L)$, then $K[Y, 1/\det(Y)]$ is a PV ring for $\phi(y) = Ay$. Conversely, if R is a PV ring with $R^\phi = K^\phi$, then $\mathrm{Quot}(R)$ is a PV extension for $\phi(y) = Ay$. □

Proof. This is clear from Proposition 2.14 and Lemma 2.7. ■

Theorem 2.16. Let K be a ϕ -pseudo field. Let R_1 and R_2 be two PV rings for the same equation $\phi(y) = Ay$, $A \in \mathrm{GL}_n(K)$. Then, there exists a finite algebraic field extension $\tilde{k}$ of $k := K^\phi$, containing $k_1 := R_1^\phi$ and $k_2 := R_2^\phi$ and an isomorphism

$$R_1 \otimes_{k_1} \tilde{k} \simeq R_2 \otimes_{k_2} \tilde{k}$$

of $K \otimes_k \tilde{k}$ - ϕ -algebras. □

Proof. This is a straightforward generalization of [58, Proposition 1.9, p. 7]. ■

Of course, the above result immediately gives the uniqueness (up to K - ϕ -isomorphisms) of PV extensions provided that K^ϕ is algebraically closed.

2.3 σ -PV extensions and σ -PV rings

In this section, we define σ -PV extensions and σ -PV rings and clarify the relation between them. Let K be a ϕ -pseudo σ -field. We study a linear difference equation

$$\phi(y) = Ay, \quad \text{where } A \in \mathrm{GL}_n(K).$$

We are mainly interested in the case when K is a field. Typically, K will be one of the $\phi\sigma$ -fields from Example 2.1. However, for consistency reasons, we will give all definitions over a general ϕ -pseudo σ -field.

If R is a K - $\phi\sigma$ -algebra, then a matrix $Y \in \mathrm{GL}_n(R)$ is called a *fundamental solution matrix* for $\phi(y) = Ay$ if $\phi(Y) = AY$.

Remark 2.17. If $Y_1, Y_2 \in \mathrm{GL}_n(R)$ are two fundamental solution matrices for $\phi(y) = Ay$, then there exists a matrix $C \in \mathrm{GL}_n(R^\phi)$ such that $Y_2 = Y_1 C$. □

Proof. This follows from the well-known computation $\phi(Y_1^{-1}Y_2) = (AY_1)^{-1}AY_2 = Y_1^{-1}Y_2$. ■

Let L be a ϕ -pseudo σ -field extension of K and $Y \in \mathrm{GL}_n(L)$ a fundamental solution matrix for $\phi(y) = Ay$. If $L = K\langle Y \rangle_\sigma$, we say that L is σ -generated by Y .

Definition 2.18. Let K be a ϕ -pseudo σ -field and $A \in \mathrm{GL}_n(K)$. A ϕ -pseudo σ -field extension L of K is called a σ -PV extension (or σ -parameterized PV extension in case we need to be more precise) for $\phi(y) = Ay$ if $L^\phi = K^\phi$ and L is σ -generated by a fundamental solution matrix for $\phi(y) = Ay$.

A K - $\phi\sigma$ -algebra R that is a ϕ -pseudo σ -domain is called a σ -PV ring for $\phi(y) = Ay$ if R is ϕ -simple and σ -generated by a fundamental solution matrix for $\phi(y) = Ay$, that is, $R = K\{Y, 1/\det(Y)\}_\sigma$ for some fundamental solution matrix $Y \in \mathrm{GL}_n(R)$. □

Remark 2.19. A Noetherian ϕ -simple ϕ -ring is automatically a ϕ -pseudo domain [60, Proposition 1.1.2, p. 2]. This is why the condition that R should be a ϕ -pseudo domain does not appear in the definition of classical PV rings (Definition 2.12). Here, in the σ -parameterized setting, one of the more subtle steps in the existence proof of σ -PV rings (or extensions) is to verify the ϕ -pseudo domain property (cf. Corollary 2.27.) □

By a σ -PV extension $L|K$, we mean a ϕ -pseudo σ -field extension L of K that is a σ -PV extension for some linear ϕ -equation $\phi(y) = Ay$, with $A \in \mathrm{GL}_n(K)$. Similarly, for σ -PV rings. The σ -field of ϕ -constants of a σ -PV extension $L|K$ will usually be denoted by k , that is,

$$k := K^\phi = L^\phi.$$

To clarify the relation between σ -PV extensions and σ -PV rings, we will use the following important observation.

Lemma 2.20. Let $L|K$ be a σ -PV extension for $\phi(y) = Ay$ with fundamental solution matrix $Y \in \mathrm{GL}_n(L)$. Set

$$L_d = K(Y, \sigma(Y), \dots, \sigma^d(Y)) \subset L, \quad d \geq 0.$$

Then, $L_d|K$ is a PV extension for the ϕ -linear system $\phi(Y) = A_d Y$, where

$$A_d = \begin{pmatrix} A & 0 & \cdots & 0 \\ 0 & \sigma(A) & \cdots & 0 \\ \vdots & & \ddots & \vdots \\ 0 & \cdots & 0 & \sigma^d(A) \end{pmatrix} \in \mathrm{GL}_{n(d+1)}(K).$$

□

Proof. Note that $K[Y, \sigma(Y), \dots, \sigma^d(Y)]$ is a ϕ -subring of L . Therefore, $K(Y, \sigma(Y), \dots, \sigma^d(Y))$ is a ϕ -pseudo field by [60, Lemma 1.3.4, p. 9]. Applying σ^i to $\phi(Y) = AY$ for $i = 0, \dots, d$ yields $\phi(\sigma^i(Y)) = \sigma^i(A)\sigma^i(Y)$. Therefore,

$$Y_d = \begin{pmatrix} Y & 0 & \cdots & 0 \\ 0 & \sigma(Y) & \cdots & 0 \\ \vdots & & \ddots & \vdots \\ 0 & \cdots & 0 & \sigma^d(Y) \end{pmatrix} \in \mathrm{GL}_{n(d+1)}(L_d)$$

is a fundamental solution matrix for $\phi(Y) = A_d Y$. Since $L_d^\phi \subset L^\phi = K^\phi$, $L_d|K$ is a PV extension for $\phi(Y) = A_d Y$. ■

The following proposition is the σ -analog of Corollary 2.15.

Proposition 2.21. Let K be a ϕ -pseudo σ -field and $A \in \mathrm{GL}_n(K)$.

- (i) If $L|K$ is a σ -PV extension for $\phi(Y) = AY$ with fundamental solution matrix $Y \in \mathrm{GL}_n(L)$, then $R := K\{Y, 1/\det(Y)\}_\sigma \subset L$ is a σ -PV ring for $\phi(Y) = AY$.
- (ii) Conversely, if R is a σ -PV ring for $\phi(Y) = AY$ with $R^\phi = K^\phi$, then $\mathrm{Quot}(R)$ is a σ -PV extension for $\phi(Y) = AY$. □

Proof. Clearly, $R := K\{Y, 1/\det(Y)\}_\sigma$ is a ϕ -pseudo domain. So, we only have to show that R is ϕ -simple. We know from Lemma 2.20 that

$$L_d := K(Y, \sigma(Y), \dots, \sigma^d(Y)) \subset L$$

is a PV extension of (K, ϕ) for every $d \geq 0$. It, thus, follows from Corollary 2.15 that

$$R_d := K[Y, \sigma(Y), \dots, \sigma^d(Y), 1/(\det(Y) \cdot \dots \cdot \det(\sigma^d(Y)))] \subset R$$

is a PV ring over K . So, R_d is ϕ -simple for every $d \geq 0$ and $R = \bigcup_{d \geq 0} R_d$. Thus, R must be ϕ -simple.

Now assume that R is a σ -PV ring with $R^\phi = K^\phi$. From Lemma 2.11, we know that $\text{Quot}(R)$ is a ϕ -pseudo σ -field and, by Lemma 2.7, we have $\text{Quot}(R)^\phi = R^\phi = K^\phi$. ■

2.4 Existence of σ -PV extensions

In this section, we will establish the existence of σ -PV rings (Theorem 2.8) and σ -PV extensions (Corollary 2.29) for a given linear ϕ -equation $\phi(y) = Ay$ under rather mild conditions on the base $\phi\sigma$ -field K . The key idea for the existence proof is the prolongation construction from [61, Lemma 2.16, p. 1392]. The differential analog of this construction has also been recently used to establish the existence of ∂ -parameterized PV extensions for linear differential or difference equations provided that the constants are algebraically closed (see [22, 62]). A more elaborate discussion of the existence of differentially parameterized PV extensions for linear differential equations (including the case of several differential parameters) can be found in [28].

Remark 2.22. The idea of the prolongation construction is easy to explain. Indeed, let K be a $\phi\sigma$ -field and $A \in \text{GL}_n(K)$. We would like to construct a σ -PV ring or a σ -PV extension for $\phi(y) = Ay$. Let

$$S := K\{X, 1/\det(X)\}_\sigma$$

be the generic solution ring for $\phi(y) = Ay$. By this, we mean that X is the $n \times n$ -matrix of σ -indeterminates, and the action of ϕ is determined by $\phi(X) = AX$. Finding a σ -PV ring for $\phi(y) = Ay$ is equivalent to finding a $\phi\sigma$ -ideal $\mathfrak{m}$ of S that is ϕ -pseudo prime and ϕ -maximal. The existence of a ϕ -maximal ideal in S is, of course, guaranteed by Zorn's lemma, but it is unclear if we can find a ϕ -maximal ideal that is additionally a σ -ideal and ϕ -pseudo prime. □

If L is a σ -PV extension for $\phi(y) = Ay$ with fundamental solution matrix $Y \in \text{GL}_n(L)$, then R_d is a PV ring over K for $\phi(y) = A_d y$, as we have already seen in Lemma 2.20 and Proposition 2.21. Thus, we should better find a $\phi\sigma$ -ideal $\mathfrak{m}$ of S such that

$$\mathfrak{m}_d := \mathfrak{m} \cap S_d, \quad S_d := K[X, \dots, \sigma^d(X), 1/\det(X \cdot \dots \cdot \sigma^d(X))] \subset S$$

is ϕ -maximal in S_d for every $d \geq 0$. Note that not every ϕ -maximal ϕ -ideal of S_d is of the form $\mathfrak{m}_d$ for some ϕ -maximal $\phi\sigma$ -ideal $\mathfrak{m}$ of S . A necessary condition is given by

$$\sigma(\mathfrak{m}_d \cap S_{d-1}) \subset \mathfrak{m}_d.$$

However, if we assume that we have already constructed a ϕ -maximal ϕ -ideal $\mathfrak{m}_d$ of S_d that satisfies this condition, we can try to construct $\mathfrak{m}_{d+1}$ by choosing a ϕ -maximal ϕ -ideal of S_{d+1} that contains $\mathfrak{m}_d$ and $\sigma(\mathfrak{m}_d)$. Then, we could define $\mathfrak{m}$ as the union of all the $\mathfrak{m}_d$'s.

There are two obstructions to this procedure that we will have to overcome:

- (i) The ideal of S_{d+1} generated by $\mathfrak{m}_d$ and $\sigma(\mathfrak{m}_d)$ might contain 1. In this case, the construction would not apply.
- (ii) The union $\bigcup \mathfrak{m}_d$ is a ϕ -maximal $\phi\sigma$ -ideal, but, *a priori*, it is unclear why it should be a ϕ -pseudo prime ideal.

Lemma 2.24 is the crucial ingredient to overcome the first difficulty. The second difficulty will be resolved in Lemma 2.26, which will eventually provide a bound for the period of $\mathfrak{m}_d$.

To prove Lemma 2.24, we need an algebraic version of Chevalley's theorem on constructible sets (cf. [32, Theorem 1.8.4, p. 239]).

Lemma 2.23. Let K be a field and $R \subset S$ an inclusion of finitely generated K -algebras. Then there exists an element $r \in R$ that is not contained in any minimal prime ideal of R and has the following property: for every prime ideal $\mathfrak{q}$ of R with $r \notin \mathfrak{q}$, there exists a prime ideal $\mathfrak{q}'$ of S with $\mathfrak{q}' \cap R = \mathfrak{q}$. $\square$

Proof. If R is an integral domain, this follows from [9, Corollaire 3, Chapitre V, Section 3.1, p. 58]. The general case can be reduced to the case in which R is an integral domain as follows. Let $\mathfrak{p}_1, \dots, \mathfrak{p}_n$ denote the minimal prime ideals of R . By [8, Proposition 16, Chapitre II, Section 2.6, p. 96], there exist minimal prime ideals $\mathfrak{p}'_1, \dots, \mathfrak{p}'_n$ of S with $\mathfrak{p}'_i \cap R = \mathfrak{p}_i$ for $i = 1, \dots, n$. For $i = 1, \dots, n$, consider the inclusion of integral domains $R/\mathfrak{p}_i \hookrightarrow S/\mathfrak{p}'_i$, and let $r_i \in R$ be such that the image $\overline{r}_i$ of r_i in $R/\mathfrak{p}_i$ is nonzero and has the property that, for every prime ideal $\mathfrak{q}$ of $R/\mathfrak{p}_i$ with $\overline{r}_i \notin \mathfrak{q}$, there exists a prime ideal $\mathfrak{q}'$ of $S/\mathfrak{p}'_i$ with

$$\mathfrak{q}' \cap (R/\mathfrak{p}_i) = \mathfrak{q}.$$

For $i = 1, \dots, n$, let

$$e_i \in (\mathfrak{p}_1 \cap \dots \cap \mathfrak{p}_{i-1} \cap \mathfrak{p}_{i+1} \cap \dots \cap \mathfrak{p}_n) \setminus \mathfrak{p}_i$$

and set

$$r = e_1 r_1 + \dots + e_n r_n.$$

Since $\bar{r} = \overline{e_i r_i} \in R/\mathfrak{p}_i$, we see that r does not belong to any minimal prime ideal of R . Let $\mathfrak{q}$ be a prime ideal of R with $r \notin \mathfrak{q}$. Then, there exists an $i \in \{1, \dots, n\}$ such that $\mathfrak{p}_i \subset \mathfrak{q}$. Since the image of $\mathfrak{q}$ in $R/\mathfrak{p}_i$ does not contain $\bar{r}_i \in R/\mathfrak{p}_i$, it follows from the construction of r_i that there exists a prime ideal $\mathfrak{q}'$ of S with $\mathfrak{q}' \cap S = R$. $\blacksquare$

Lemma 2.24. Let K be a field and let R be a finitely generated K -algebra. For $d \geq 0$, let $R_0, \dots, R_{d+1}$ denote isomorphic copies of R . Let $\mathfrak{a} \subset R_0 \otimes \dots \otimes R_d$ and $\mathfrak{b} \subset R_1 \otimes \dots \otimes R_{d+1}$ be ideals not containing 1. (The tensors are understood to be over K .) Assume that

$$\mathfrak{a} \cap (R_1 \otimes \dots \otimes R_d) = \mathfrak{b} \cap (R_1 \otimes \dots \otimes R_d). \quad (2.4)$$

Then, the ideal of $R_0 \otimes \dots \otimes R_{d+1}$ generated by $\mathfrak{a}$ and $\mathfrak{b}$ does not contain 1. $\square$

Proof. Let $\mathfrak{c} = \mathfrak{a} \cap (R_1 \otimes \dots \otimes R_d) = \mathfrak{b} \cap (R_1 \otimes \dots \otimes R_d) \subset R_1 \otimes \dots \otimes R_d$ and consider the inclusions

$$\begin{aligned} B &:= (R_1 \otimes \dots \otimes R_d)/\mathfrak{c} \hookrightarrow (R_0 \otimes \dots \otimes R_d)/\mathfrak{a} \hookrightarrow ((R_0 \otimes \dots \otimes R_d)/\mathfrak{a}) \otimes R_{d+1} \\ &= (R_0 \otimes \dots \otimes R_{d+1})/(\mathfrak{a}). \end{aligned}$$

By Lemma 2.23, there exists an element $r_a \in B$ not contained in any minimal prime ideal of B and such that, for every prime ideal $\mathfrak{q}$ of B with $r \notin \mathfrak{q}$, there exists a prime ideal $\mathfrak{q}'$ of $(R_0 \otimes \dots \otimes R_d)/(\mathfrak{a})$ with $\mathfrak{q}' \cap B = \mathfrak{q}$.

Let $r_b \in B$ be defined similarly. Since $r_a r_b$ does not belong to any minimal prime ideal of B , there exists a prime ideal $\mathfrak{q}$ of B with $r_a, r_b \notin \mathfrak{q}$. Then, by construction of r_a and r_b , there exist prime ideals

$$\mathfrak{q}'_a \subset (R_0 \otimes \dots \otimes R_{d+1})/(\mathfrak{a}) \quad \text{and} \quad \mathfrak{q}'_b \subset (R_0 \otimes \dots \otimes R_{d+1})/(\mathfrak{b})$$

such that

$$\mathfrak{q}'_a \cap B = \mathfrak{q} = \mathfrak{q}'_b \cap B.$$

Then, $\mathfrak{q}'_a$ and $\mathfrak{q}'_b$ correspond to prime ideals $\mathfrak{q}_a$ and $\mathfrak{q}_b$ of $R_0 \otimes \cdots \otimes R_{d+1}$ with

$$\mathfrak{q}_a \cap (R_1 \otimes \cdots \otimes R_d) = \mathfrak{q}_b \cap (R_1 \otimes \cdots \otimes R_d) =: \mathfrak{d}.$$

Denoting the residue field of a prime ideal $\mathfrak{p}$ of a ring S by $k(\mathfrak{p}) = \text{Quot}(S/\mathfrak{p})$, we have $k(\mathfrak{d}) \subset k(\mathfrak{q}_a)$ and $k(\mathfrak{d}) \subset k(\mathfrak{q}_b)$. Let L be a field extension of $k(\mathfrak{d})$ containing $k(\mathfrak{q}_a)$ and $k(\mathfrak{q}_b)$ and let

$$\psi: R_0 \otimes \cdots \otimes R_{d+1} \rightarrow L$$

be the morphism of K -algebras that extends the canonical map

$$R_1 \otimes \cdots \otimes R_d \rightarrow k(\mathfrak{d}) \subset L$$

by sending R_0 and R_{d+1} to their canonical images in $k(\mathfrak{q}_a)$ and $k(\mathfrak{q}_b)$, respectively. Since $\mathfrak{a} \subset \mathfrak{q}_a$ and $\mathfrak{b} \subset \mathfrak{q}_b$, the kernel of ψ is a prime ideal of $R_0 \otimes \cdots \otimes R_{d+1}$, which contains $\mathfrak{a}$ and $\mathfrak{b}$. Therefore, the ideal generated by $\mathfrak{a}$ and $\mathfrak{b}$ does not contain 1. ■

For the convenience of the readers who prefer the geometric language, we have included a geometric proof of Lemma 2.24. This proof is more intuitive than the algebraic proof given above, but the algebraic proof is more accessible.

Proof. We set $X_i := \text{Spec}(R_i)$ for $i = 0, \dots, d+1$. Let Y and Z denote the closed subschemes of $X_0 \times \cdots \times X_d$ and $X_1 \times \cdots \times X_{d+1}$ defined by $\mathfrak{a}$ and $\mathfrak{b}$, respectively. Then, the ideal $\mathfrak{a}'$ generated by $\mathfrak{a}$ in $R_0 \otimes \cdots \otimes R_{d+1}$ defines the closed subscheme $Y \times X_{d+1} \subset X_0 \times \cdots \times X_{d+1}$. Similarly, the ideal $\mathfrak{b}'$ generated by $\mathfrak{b}$ in $R_0 \otimes \cdots \otimes R_{d+1}$ defines the closed subscheme $X_0 \times Z \subset X_0 \times \cdots \times X_{d+1}$. Since the sum of the ideals $\mathfrak{a}'$ and $\mathfrak{b}'$ corresponds to the intersection of the closed subschemes $Y \times X_{d+1}$ and $X_0 \times Z$, the statement of the lemma is equivalent to

$$(Y \times X_{d+1}) \cap (X_0 \times Z) \subset X_0 \times \cdots \times X_{d+1}$$

being nonempty. Let

$$\pi_{1d}: X_0 \times \cdots \times X_{d+1} \rightarrow X_1 \times \cdots \times X_d, \quad (x_0, \dots, x_{d+1}) \mapsto (x_1, \dots, x_d)$$

denote the projection onto the factors “in the middle”. The ideal $\mathfrak{a} \cap (R_1 \otimes \cdots \otimes R_d)$ of $R_1 \otimes \cdots \otimes R_d$ corresponds to the Zariski closure $\overline{\pi_{1d}(Y \times X_{d+1})} \subset X_1 \times \cdots \times X_d$; similarly

for b. Assumption (2.4), thus, means that

$$\overline{\pi_{1d}(Y \times X_{d+1})} = \overline{\pi_{1d}(X_0 \times Z)} =: W.$$

By Chevalley's theorem, the image of a morphism of schemes of finite type over a field contains a dense open subset of its closure. Thus, there exist open dense subsets $U, V \subset W$ with

$$U \subset \pi_{1d}(Y \times X_{d+1}) \quad \text{and} \quad V \subset \pi_{1d}(X_0 \times Z).$$

Then, $U \cap V$ is also dense and open in W . In particular,

$$U \cap V \subset \pi_{1d}(Y \times X_{d+1}) \cap \pi_{1d}(X_0 \times Z)$$

is nonempty. But, if $x = (x_0, \dots, x_{d+1}) \in Y \times X_{d+1}$ and $x' = (x'_0, \dots, x'_{d+1}) \in X_0 \times Z$ satisfy

$$(x_1, \dots, x_d) = \pi_{1d}(x) = \pi_{1d}(x') = (x'_1, \dots, x'_d),$$

then $(x_0, x_1, \dots, x_d, x'_{d+1}) \in (Y \times X_{d+1}) \cap (X_0 \times Z)$. ■

If R is a ϕ -ring, we denote the ring of its ϕ -periodic elements by

$$R^{\phi^\infty} = \{r \in R \mid \exists m \geq 1 \text{ such that } \phi^m(r) = r\}.$$

It is a ϕ -subring of R .

Remark 2.25. If K is a ϕ -field, then K^{ϕ^∞} is the relative algebraic closure of K^ϕ in K [43, Theorem 2.1.12, p. 114]. In particular, if K^ϕ is algebraically closed, then $K^{\phi^\infty} = K^\phi$. □

Analogues of the generic solution field U in the following lemma appear in [13, Section 4] and [50]. The relation between the periodic elements in a universal solution field and the period of a PV ring, which we shall eventually use to bound the period of $\mathfrak{m}_d$, has been found in [13]. In the language of [13], the following lemma essentially says that the m -invariant of the systems $\phi(y) = A_d y$ is bounded (as a function of $d \geq 0$).

Lemma 2.26. Let K be a $\phi\sigma$ -field such that $K^{\phi^\infty} = K^\phi$. Let $A \in \text{GL}_n(K)$ and let X denote the $n \times n$ -matrix of σ -indeterminates over K . Set

$$U = K\langle X \rangle_\sigma (= \text{Quot}(K\{X_{ij} \mid 1 \leq i, j \leq n\}_\sigma))$$

and define a $\phi\sigma$ -structure on U by $\phi(\sigma^i(X)) = \sigma^i(A)\sigma^i(X)$, $i \geq 0$. Then, U^{ϕ^∞} is a finite field extension of U^ϕ . $\square$

Proof. We have a tower of $\phi\sigma$ -fields $KU^\phi \subset KU^{\phi^\infty} \subset U$. By construction, U is a finitely σ -generated σ -field extension of KU^ϕ . Since an intermediate σ -field of a finitely σ -generated σ -field extension is itself finitely σ -generated [43, Theorem 4.4.1, p. 292], it follows that KU^{ϕ^∞} is finitely σ -generated over KU^ϕ . Hence, we can find

$$a_1, \dots, a_m \in U^{\phi^\infty}$$

that σ -generate KU^{ϕ^∞} as a σ -field extension of KU^ϕ . We claim that

$$U^{\phi^\infty} = U^\phi \langle a_1, \dots, a_m \rangle_\sigma.$$

The inclusion " $\supset$ " is clear. So, let $a \in U^{\phi^\infty}$. Let $(b_i)_{i \in I}$ be a K^ϕ -basis of $U^\phi \langle a_1, \dots, a_m \rangle_\sigma$. As

$$a \in KU^{\phi^\infty} = KU^\phi \langle a_1, \dots, a_m \rangle_\sigma,$$

we can write

$$a = \frac{\sum \lambda_i \cdot b_i}{\sum \mu_i \cdot b_i}$$

with $\lambda_i, \mu_i \in K$. Multiplying by the denominator yields

$$\sum \mu_i \cdot a \cdot b_i = \sum \lambda_i \cdot b_i. \quad (2.5)$$

We can choose an integer $e \geq 1$ such that $a, b_i \in U^{\phi^e}$ whenever λ_i or μ_i is nonzero. Then, (2.5) signifies that the family

$$(a \cdot b_i, b_j)_{i,j \in I} \quad \text{in } U^{\phi^e}$$

is K -linearly dependent. Since K is linearly disjoint from U^{ϕ^e} over $K^{\phi^e} = K^\phi$ (Lemma 2.10), we can find a nontrivial relation

$$\sum \mu'_i \cdot a \cdot b_i = \sum \lambda'_i \cdot b_i \quad (2.6)$$

with $\mu'_i, \lambda'_i \in K^\phi$. Suppose that

$$\sum \mu'_i b_i = 0.$$

Then, also $\sum \lambda'_i b_i = 0$. Since the b_i 's are K^ϕ -linearly independent, this is only possible if relation (2.6) is trivial. Therefore, we can divide by the denominator to find that

$$a = \frac{\sum \lambda'_i \cdot b_i}{\sum \mu'_i \cdot b_i} \in U^\phi \langle a_1, \dots, a_m \rangle_\sigma$$

as desired. Now let $e \geq 1$ be such that $a_1, \dots, a_m \in U^{\phi^e}$. Then, it follows from

$$U^{\phi^\infty} = U^\phi \langle a_1, \dots, a_m \rangle_\sigma$$

that $U^{\phi^\infty} = U^{\phi^e} =: F$. Let $g = \phi|_F$ and $G = \{g^0, g, \dots, g^{e-1}\} \subset \text{Aut}(F|U^\phi)$. Since $F^G = U^\phi$, we have $[F : U^\phi] = |G|$ [42, Chapter VI, Theorem 1.8]. Since $|G| \leq e$, we, therefore, obtain $[U^{\phi^\infty} : U^\phi] \leq e$. $\blacksquare$

Corollary 2.27. Let K be a $\phi\sigma$ -field such that $K^{\phi^\infty} = K^\phi$. Let $A \in \text{GL}_n(K)$. For $d \geq 0$, let R_d be a PV ring for $\phi(y) = A_d y$, where

$$A_d = \begin{pmatrix} A & 0 & \cdots & 0 \\ 0 & \sigma(A) & \cdots & 0 \\ \vdots & & \ddots & \vdots \\ 0 & \cdots & 0 & \sigma^d(A) \end{pmatrix} \in \text{GL}_{n(d+1)}(K).$$

Then, the sequence $(\text{period}(R_d))_{d \geq 0}$ is bounded. $\square$

Proof. Let $U = K \langle X \rangle_\sigma$ as in Lemma 2.26. We will show that, for $d \geq 0$,

$$\text{period}(R_d) \leq [U^{\phi^\infty} : U^\phi].$$

Let $\overline{U^\phi}$ denote an algebraic closure of U^ϕ , considered as a constant ϕ -ring. We know that K is a regular field extension of K^ϕ . (By assumption, $K^\phi = K^{\phi^\infty}$ is relatively algebraically closed in K (see Remark 2.25) and K is always separable over K^ϕ [60, Corollary 1.4.16, p. 16]). Therefore, $K \otimes_{K^\phi} \overline{U^\phi}$ is an integral domain. Moreover, $K \otimes_{K^\phi} \overline{U^\phi}$ is ϕ -simple by

Lemma 2.9. It follows that

$$K' := \text{Quot}(K \otimes_{K^\phi} \overline{U^\phi})$$

is a ϕ -field with $K'^\phi = (K \otimes_{K^\phi} \overline{U^\phi})^\phi = \overline{U^\phi}$ (by Lemma 2.7) algebraically closed. Indeed, let $\{b_j\}$ be a basis of $\overline{U^\phi}$ over K^ϕ and, for some m ,

$$c = \sum_{i=1}^m a_i \otimes b_i \in (K \otimes_{K^\phi} \overline{U^\phi})^\phi.$$

Then,

$$0 = \phi(c) - c = \sum_{i=1}^m (\phi(a_i) - a_i) \otimes b_i,$$

which implies that, for all i , $1 \leq i \leq m$, $a_i \in K^\phi$. It is clear from the definition of U that

$$KU^\phi(X, \dots, \sigma^d(X)) \subset U$$

is a PV extension of KU^ϕ for the linear ϕ -equation $\phi(y) = A_d y$. It follows from Corollary 2.15 that

$$S_d := KU^\phi[X, \dots, \sigma^d(X), 1/\det(X \cdot \dots \cdot \sigma^d(X))]$$

is a PV ring over KU^ϕ . Then, $S'_d := S_d \otimes_{U^\phi} \overline{U^\phi}$ is a PV ring over

$$KU^\phi \otimes_{U^\phi} \overline{U^\phi} = \text{Quot}(K \otimes_{K^\phi} U^\phi) \otimes_{U^\phi} \overline{U^\phi} = \text{Quot}(K \otimes_{K^\phi} \overline{U^\phi}) = K'$$

by Lemma 2.9. Note that $S_d \subset U$ is an integral domain and that

$$\text{period}(S'_d) \leq [U^{\phi^\infty} : U^\phi]$$

as U^{ϕ^∞} is the relative algebraic closure of U^ϕ in U . As R_d is a PV ring for $\phi(y) = A_d y$ over K , $R_d \otimes_{R_d^\phi} \overline{U^\phi}$ is ϕ -simple by Lemma 2.9. (Note that R_d^ϕ can be embedded in $\overline{U^\phi}$ by Proposition 2.14.) The canonical map

$$K \otimes_{K^\phi} \overline{U^\phi} \rightarrow R_d \otimes_{R_d^\phi} \overline{U^\phi}$$

is injective, because $K \otimes_{K^\phi} \overline{U^\phi}$ is ϕ -simple. Localizing this inclusion at the nonzero divisors of $K \otimes_{K^\phi} \overline{U^\phi}$, we obtain a PV ring R'_d over K' . Since $K'^\phi = \overline{U^\phi}$ is algebraically closed,

R'_d and S'_d are isomorphic. It follows that

$$\text{period}(R_d) \leq \text{period}(R'_d) = \text{period}(S'_d) \leq [U^{\phi^\infty} : U^\phi]. \quad \blacksquare$$

We are now prepared to establish the main existence theorem.

Theorem 2.28. Let K be a $\phi\sigma$ -field such that $K^{\phi^\infty} = K^\phi$, $\sigma: K^\phi \rightarrow K^\phi$ is an automorphism, and $A \in \text{GL}_n(K)$. Then, there exists a σ -PV ring R for $\phi(y) = Ay$ such that R^ϕ is an algebraic field extension of K^ϕ . $\square$

Proof. We first assume that $\sigma: K \rightarrow K$ is an automorphism. Let X be the $n \times n$ -matrix of σ -indeterminates over K . We denote the localization of the σ -polynomial ring $K\{X_{ij} \mid 1 \leq i, j \leq n\}_\sigma$ at the multiplicatively closed subset generated by $\det(X), \sigma(\det(X)), \dots$ by S . This is naturally a K - σ -algebra. We define a $\phi\sigma$ -structure on S by setting

$$\phi(X) = AX, \quad \phi(\sigma(X)) = \sigma(A)\sigma(X), \quad \phi(\sigma^2(X)) = \sigma^2(A)\sigma^2(X), \quad \dots$$

For $0 \leq i \leq j$, we also define the following K - ϕ -subalgebras of S :

$$\begin{aligned} S_{i,j} &= K\left[\sigma^i(X), \frac{1}{\sigma^i(\det(X))}, \dots, \sigma^j(X), \frac{1}{\sigma^j(\det(X))}\right] \\ &= K\left[\sigma^i(X), \dots, \sigma^j(X), \frac{1}{\det(\sigma^i(X) \dots \sigma^j(X))}\right] \subset S, \quad S_j := S_{0,j}. \end{aligned}$$

We will show by induction on $d \geq 0$ that there exists a sequence $(\mathfrak{m}_d)_{d \geq 0}$ with the following properties:

- (i) $\mathfrak{m}_d$ is a ϕ -maximal ϕ -ideal of S_d ;
- (ii) $\mathfrak{m}_d \cap S_{d-1} = \mathfrak{m}_{d-1}$;
- (iii) $\sigma^{-1}(\mathfrak{m}_d) = \mathfrak{m}_{d-1}$, where $\sigma: S_{d-1} \rightarrow S_d$.

For $d=0$, we can choose $\mathfrak{m}_0$ to be any ϕ -maximal ϕ -ideal of $S_0 = K[X, 1/\det(X)]$. Assume that a sequence $\mathfrak{m}_0, \dots, \mathfrak{m}_d$ with the desired properties has been already constructed. We will construct $\mathfrak{m}_{d+1}$. Let $\mathfrak{a}$ denote the ideal of S_{d+1} generated by $\mathfrak{m}_d$ and $\sigma(\mathfrak{m}_d)$. The crucial step now is to show that $1 \notin \mathfrak{a}$. For this, we would like to apply Lemma 2.24. Note that S_{d+1} is the $d+2$ -fold tensor product of S_0 with itself. Since σ is an automorphism on K ,

$$\sigma: S_d \rightarrow S_{1,d+1}$$

is an isomorphism and so $\sigma(\mathfrak{m}_d)$ is an ideal of $S_{1,d+1}$. We need to verify that

$$\mathfrak{m}_d \cap S_{1,d} = \sigma(\mathfrak{m}_d) \cap S_{1,d}.$$

Let $f \in \mathfrak{m}_d \cap S_{1,d}$. Then, f is of the form $f = \sigma(g)$ for some $g \in S_{d-1}$. Since $f \in \mathfrak{m}_d$, we have

$$g \in \sigma^{-1}(\mathfrak{m}_d) = \mathfrak{m}_{d-1} \subset \mathfrak{m}_d.$$

Thus, $f \in \sigma(\mathfrak{m}_d)$. Now let $f \in \sigma(\mathfrak{m}_d) \cap S_{1,d}$. Then, f is of the form $f = \sigma(g)$ with

$$g \in \mathfrak{m}_d \cap S_{d-1} = \mathfrak{m}_{d-1}.$$

So $f = \sigma(g) \in \mathfrak{m}_d$. We can thus apply Lemma 2.24 to conclude that $1 \notin \mathfrak{a}$. By construction, $\mathfrak{a}$ is a ϕ -ideal of S_{d+1} . Let $\mathfrak{m}_{d+1}$ be a ϕ -maximal ϕ -ideal of S_{d+1} containing $\mathfrak{a}$. Then,

$$\mathfrak{m}_{d+1} \cap S_d \quad \text{and} \quad \sigma^{-1}(\mathfrak{m}_{d+1})$$

are ϕ -ideals of S_d containing $\mathfrak{m}_d$. As $\mathfrak{m}_d$ is ϕ -maximal in S_d , it follows that

$$\mathfrak{m}_{d+1} \cap S_d = \mathfrak{m}_d \quad \text{and} \quad \sigma^{-1}(\mathfrak{m}_{d+1}) = \mathfrak{m}_d.$$

This concludes the inductive step. Now that we have constructed the sequence $(\mathfrak{m}_d)_{d \geq 0}$, we can define

$$\mathfrak{m} := \bigcup_{d \geq 0} \mathfrak{m}_d.$$

This is a $\phi\sigma$ -ideal of $S = \bigcup_{d \geq 0} S_d$. Since the $\mathfrak{m}_d$'s are ϕ -maximal, it follows that $\mathfrak{m}$ is also ϕ -maximal. The next crucial step is to show that $\mathfrak{m}$ is ϕ -pseudo prime.

In general, a ϕ -maximal ϕ -ideal need not be ϕ -pseudo prime. However, a ϕ -maximal ϕ -ideal that has only finitely many minimal prime ideals is ϕ -pseudo prime [60, Proposition 1.1.2, p. 2]. In particular, in a Noetherian ϕ -ring, every ϕ -maximal ϕ -ideal is ϕ -pseudo prime. So the $\mathfrak{m}_d$'s are ϕ -pseudo prime ideals.

For any prime ideal $\mathfrak{q} \subset S_d$ that is minimal above $\mathfrak{m}_d$, there exists a prime ideal $\mathfrak{q}' \subset S_{d+1}$ that is minimal above $\mathfrak{m}_{d+1}$ such that $\mathfrak{q}' \cap S_d = \mathfrak{q}$ by [8, Proposition 16, Section 2, Chapter II]. Therefore, the sequence $(\text{period}(\mathfrak{m}_d))_{d \geq 0}$ is nondecreasing. Since $R_d := S_d/\mathfrak{m}_d$ is a PV ring for $\phi(y) = A_d y$, it follows from Corollary 2.27 that there exists an integer $b \geq 1$ such that $\text{period}(\mathfrak{m}_d) = b$ for all sufficiently large d . This shows that there are precisely b prime ideals minimal above $\mathfrak{m}$.

So, $\mathfrak{m}$ is ϕ -pseudo prime and $R := S/\mathfrak{m}$ is a ϕ -pseudo domain. It is clear from the construction that R is a σ -PV ring for $\phi(y) = Ay$ over K . It remains to see that R^ϕ is algebraic over K^ϕ . But R is the union of the R_d 's and the R_d 's are PV rings over K , so R_d^ϕ is algebraic over K^ϕ (Lemma 2.13) and, consequently, R^ϕ is algebraic over K^ϕ . This concludes the proof for the case that $\sigma: K \rightarrow K$ is surjective.

Now let $\sigma: K \rightarrow K$ be arbitrary. We consider the inversive closure K^* of K with respect to σ (see [43, Definition 2.1.6, p. 109].) For every $a \in K^*$, there exists an integer $l \geq 1$ such that $\sigma^l(a) \in K$. We naturally extend ϕ from K to K^* by

$$\phi(a) = \sigma^{-l}(\phi(\sigma^l(a))).$$

Suppose that $a \in K^{*\phi^d}$. Then,

$$a = \phi^d(a) = \sigma^{-l}(\phi^d(\sigma^l(a)))$$

and so

$$\sigma^l(a) = \phi^d(\sigma^l(a)),$$

that is, $\sigma^l(a) \in K^{\phi^d} = K^\phi$. By the hypothesis, K^ϕ is σ -inversive. Therefore, $a \in K^\phi$. It follows that

$$K^{*\phi^\infty} = K^\phi = K^{*\phi}.$$

By the first part of the proof, there exists a σ -PV ring R^* over K^* for $\phi(y) = Ay$ with $R^{*\phi}$ algebraic over K^ϕ . Let $Y \in \mathrm{GL}_n(R^*)$ denote a fundamental matrix. We claim that

$$R := K\{Y, 1/\det(Y)\}_\sigma \subset R^*$$

is a σ -PV ring for $\phi(y) = Ay$ over K with R^ϕ algebraic over K^ϕ . As $R^{*\phi}$ is algebraic over $K^{*\phi} = K^\phi$, R^ϕ is algebraic over K^ϕ . So it only remains to show that R is ϕ -simple. For this, it suffices to show that

$$R_d := K[Y, 1/\det(Y), \dots, \sigma^d(Y), 1/\det(\sigma^d(Y))]$$

is ϕ -simple for every $d \geq 0$. Let L^* denote the total quotient ring of R^* and L_d the total quotient ring of R_d . Since R^* is ϕ -simple, we have $L^{*\phi} = R^{*\phi}$ by Lemma 2.7. As $L_d \subset L^*$, it follows that L_d^ϕ is algebraic over K^ϕ . By Proposition 2.14, this implies that R_d is ϕ -simple. ■

Corollary 2.29 (Existence of σ -PV extensions). Let K be a $\phi\sigma$ -field and $A \in \text{GL}_n(K)$. Assume that K^ϕ is an algebraically closed inversive σ -field. Then, there exists a σ -PV extension for $\phi(y) = Ay$. $\square$

Proof. By Remark 2.25, if K^ϕ is algebraically closed, then $K^{\phi^\infty} = K^\phi$. The statement now follows from Theorem 2.28 and Proposition 2.21. $\blacksquare$

2.5 Existence of σ -PV extensions for some specific base fields

The purpose of this section is to establish the existence of σ -PV extensions over important $\phi\sigma$ -fields like $K = \mathbb{C}(t, z)$, where

$$\phi(f(t, z)) = f(t, z + 1) \quad \text{and} \quad \sigma(f(t, z)) = f(qt, z) \quad \text{or} \quad \sigma(f(t, z)) = f(t + \alpha, z)$$

for some $q, \alpha \in \mathbb{C}^\times$. Note that the general existence result for σ -PV extensions (Corollary 2.29) does not apply because $K^\phi = \mathbb{C}(t)$ is not algebraically closed.

We will show quite generally that, for every linear ϕ -equation $\phi(y) = Ay$ over $K = k(z)$, there exists a σ -PV extension, where k is an arbitrary σ -field of characteristic zero. Moreover, we give a very concrete recipe how σ -PV rings over such K can be constructed inside rings of sequences; cf. [58, Proposition 4.1, p. 45].

Let k be a field. The ring Seq_k of sequences in k (cf. [58, Example 1.3, p. 4]) consists of all sequences

$$a = (a(0), a(1), \dots), \quad a(0), a(1), \dots \in k,$$

and two sequences are identified if they agree starting from some index. The ring structure of Seq_k is given by the componentwise addition and multiplication. By setting

$$\phi((a(0), a(1), a(2), \dots)) = (a(1), a(2), \dots),$$

we turn Seq_k into an inversive ϕ -ring. If k is a σ -field, then Seq_k naturally becomes a $\phi\sigma$ -ring by setting

$$\sigma((a(0), a(1), \dots)) = (\sigma(a(0)), \sigma(a(1)), \dots).$$

Note that $\text{Seq}_k^\phi = k$. We consider $k(z)$, the field of rational function in one variable over k , as $\phi\sigma$ -field by setting

$$\phi(f(z)) = f(z + 1), \quad f \in k(z), \quad \text{and} \quad \sigma(z) = z.$$

If $\text{char } k = 0$, we can define a $\phi\sigma$ -embedding

$$k(z) \rightarrow \text{Seq}_k \quad \text{by} \quad f \mapsto (f(0), f(1), \dots).$$

The expression $f(i)$ is well-defined for $i \gg 0$, as the denominator of $f \in k(z)$ has only finitely many zeros.

Proposition 2.30. Let k be a σ -field of characteristic zero and consider $K = k(z)$ as a $\phi\sigma$ -field via

$$\phi(f(z)) = f(z+1) \quad \text{and} \quad \sigma(z) = z.$$

Let $A \in \text{GL}_n(K)$ and $i_0 \geq 0$ be an integer such that $A(i)$ is well-defined and $\det(A(i)) \neq 0$ for all $i \geq i_0$. Define $Y \in \text{GL}_n(\text{Seq}_k)$ by

$$Y(i_0) = \text{id} \quad \text{and} \quad Y(i) = A(i-1)Y(i-1), \quad i > i_0.$$

Then, Y is a fundamental solution matrix for $\phi(y) = Ay$ and

$$K\{Y, 1/\det(Y)\}_\sigma \subset \text{Seq}_k$$

is a σ -PV ring for $\phi(y) = Ay$. Moreover, there exists a σ -PV extension for $\phi(y) = Ay$. $\square$

Proof. It is clear that Y is a fundamental solution matrix and that

$$R := K\{Y, 1/\det(Y)\}_\sigma$$

is a $\phi\sigma$ -ring. It remains to see that R is a ϕ -simple ϕ -pseudo domain. To see that R is ϕ -simple, it suffices to show that

$$R_d := K[Y, \dots, \sigma^d(Y), 1/\det(Y \cdots \sigma^d(Y))]$$

is ϕ -simple for every $d \geq 0$. Note that

$$Y_d = \begin{pmatrix} Y & 0 & \cdots & 0 \\ 0 & \sigma(Y) & \cdots & 0 \\ \vdots & & \ddots & \vdots \\ 0 & \cdots & 0 & \sigma^d(Y) \end{pmatrix} \in \text{GL}_{n(d+1)}(R)$$

is a fundamental solution matrix for $\phi(y) = A_d y$ (cf. Lemma 2.20). By [64, Proposition 2.4, p. 4], there exists a PV ring S_d for $\phi(y) = A_d y$ over K inside Seq_k . As $\text{Seq}_k^\phi = k$ and two fundamental solution matrices for the same equation only differ by multiplication by a matrix with constant entries, it follows that $R_d = S_d$. In particular, R_d is ϕ -simple. As in the proof of Theorem 2.28, it follows from Corollary 2.27 that R is a ϕ -pseudo domain.

As $\text{Seq}_k^\phi = k = K^\phi$, Proposition 2.21 implies that $\text{Quot}(R)$ is a σ -PV extension for $\phi(y) = Ay$. ■

Remark 2.31. Let $Y \in \text{GL}_n(\text{Seq}_k)$ be defined as in Proposition 2.30. It is unclear whether or not $K\langle Y \rangle_\sigma \subset \text{Seq}_k$ (see (2.2)) is a σ -PV extension for $\phi(y) = Ay$. The difficulty here is to know that a nonzero divisor of $K\langle Y, 1/\det(Y) \rangle_\sigma \subset \text{Seq}_k$ is a unit in Seq_k . This problem is closely related to the generalization of the Skolem–Mahler–Lech theorem to rational function coefficients (see [64]). It follows from [64, Corollary 3.4, p. 8] that $K\langle Y \rangle_\sigma \subset \text{Seq}_k$ is a σ -PV extension for $\phi(y) = Ay$ if $A \in \text{GL}_n(k[z])$. □

2.6 Uniqueness

In this section, we will establish the uniqueness of σ -PV rings and σ -PV extensions (for a given equation $\phi(y) = Ay$). In other words, we prove a result analogous to the classical uniqueness theorem (Theorem 2.16). The main difficulty is to understand what the σ -analog of the algebraic closure in the classical case is. There is a notion of a difference-closed difference field that has been used and studied extensively by model theorists (see, e.g., [11, 12]).

Definition 2.32. A σ -field k is called *σ -closed* if for every finitely σ -generated k - σ -algebra R which is a σ -domain, there exists a k - σ -morphism $R \rightarrow k$. □

In contrast to differential algebra, there appears to be no satisfactory notion of a σ -closure of a σ -field. Kolchin preferred the term “constrainedly closed” to “differentially closed” because a differentially closed differential field can have proper differential algebraic extensions. The following definition can be seen as an adaptation of Kolchin’s notion of constrained extensions of differential fields ([41]) to difference algebra.

Definition 2.33. Let $L|K$ be an extension of σ -pseudo fields. We say that L is *constrained* over K if, for every finite tuple a from L , there exists a nonzero divisor $b \in L$ such that (0) is the only σ -pseudo prime ideal of $K\{a, 1/b\}_\sigma$. □

The basic properties of constrained extensions of σ -pseudo fields have been established in [61, Section 2.1]. The relation to σ -closed σ -fields is given by the fact that a σ -closed σ -field does not have proper constrained σ -field extensions. More generally, every finitely σ -generated σ -pseudo field extension of a σ -closed σ -field k is of the form $k \oplus \cdots \oplus k$ (see [61, Example 2.8, p. 1388]).

The following theorem is the crucial tool from difference algebra for proving our uniqueness result. It can be seen as a difference analog of a theorem of Chevalley. For a prime ideal $\mathfrak{q}$ in a σ -ring R and $r \in R$, we write

$$r \notin_{\sigma} \mathfrak{q}$$

if $\sigma^d(r) \notin \mathfrak{q}$ for every $d \geq 0$.

Theorem 2.34. Let $R \subset S$ be an inclusion of σ -rings such that S is finitely σ -generated over R . Assume that R is a σ -domain and $(0) \subset S$ is a finite intersection of σ -pseudo prime ideals. Then, there exist $0 \neq r \in R$ and an integer $l \geq 1$ such that, for every $d \geq 1$ and σ^d -prime ideal $\mathfrak{q}$ of R with $r \notin_{\sigma} \mathfrak{q}$, there exists a σ^{ld} -prime ideal $\mathfrak{q}'$ of S with $\mathfrak{q}' \cap R = \mathfrak{q}$. $\square$

Proof. This is a slight generalization of [61, Theorem 1.15, p. 1384], where it is assumed that S is a σ -domain. There exists a minimal prime ideal $\hat{\mathfrak{q}}$ of S with $\hat{\mathfrak{q}} \cap R = (0)$ [8, Chapter II, Section 2, Section 6, Proposition 16, p. 74]. By assumption, $\hat{\mathfrak{q}}$ is a $\sigma^{\hat{d}}$ -prime ideal for some $\hat{d} \geq 1$. We can now apply [61, Theorem 1.15, p. 1384] to the inclusion $R \subset S/\hat{\mathfrak{q}}$ of $\sigma^{\hat{d}}$ -domains to obtain $0 \neq r \in R$ and an integer $\hat{l} \geq 1$ such that, for every $\sigma^{\hat{l}\hat{d}}$ -prime ideal $\mathfrak{q}$ of R with $r \notin_{\sigma^{\hat{d}}} \mathfrak{q}$, there exists a $\sigma^{\hat{l}\hat{d}}$ -prime ideal $\mathfrak{q}'$ of $S/\hat{\mathfrak{q}}$ with $\mathfrak{q}' \cap R = \mathfrak{q}$. Set $l := \hat{d}\hat{l}$. Observing that a σ^d -prime ideal is a $\sigma^{\hat{d}d}$ -prime ideal and that $r \notin_{\sigma} \mathfrak{q}$ implies $r \notin_{\sigma^{\hat{d}}} \mathfrak{q}$ yields the claim of the theorem. $\blacksquare$

We will need a few more preparatory results.

Lemma 2.35. Let k be an inversive σ -field and R a k - σ -algebra with $\sigma : R \rightarrow R$ injective. If (λ_i) is a family of k -linearly independent elements from R , then the family $(\sigma(\lambda_i))$ is k -linearly independent as well. $\square$

Proof. If $\sum a_i \sigma(\lambda_i) = 0$ with $a_i \in k$, then, as k is inversive, we can find $b_i \in k$ with $\sigma(b_i) = a_i$. We have $\sigma(\sum b_i \lambda_i) = 0$, and this implies $\sum b_i \lambda_i = 0$. Therefore, the b_i 's and also the a_i 's are all zeroes. $\blacksquare$

Lemma 2.36. Let k be an inversive σ -field and R a k - σ -algebra with $\sigma: R \rightarrow R$ injective. Then, $\sigma: R \otimes_k K \rightarrow R \otimes_k K$ is injective for every σ -field extension K of k . Moreover, if $\mathfrak{a}$ is a reflexive σ -ideal of R (i.e., $\sigma^{-1}(\mathfrak{a}) = \mathfrak{a}$), then $\mathfrak{a} \otimes_k K$ is a reflexive σ -ideal of $R \otimes_k K$. $\square$

Proof. Let (λ_i) be a k -basis of K and $s = \sum r_i \otimes \lambda_i \in R \otimes_k K$ with $\sigma(s) = 0$. Then, $\sum \sigma(r_i) \otimes \sigma(\lambda_i) = 0$ implies $\sigma(r_i) = 0$, because the family $(\sigma(\lambda_i))$ is k -linearly independent by Lemma 2.35. Since σ is injective on R , $s = 0$. The latter claim of the lemma follows by applying the above result to $R/\mathfrak{a}$. $\blacksquare$

Proposition 2.37. Let K be a $\phi\sigma$ -field such that $K^{\phi^\infty} = K^\phi$ and $\sigma: K^\phi \rightarrow K^\phi$ is surjective. Let R be a ϕ -simple K - $\phi\sigma$ -algebra that is a ϕ -pseudo domain and finitely σ -generated over K . Then, R^ϕ is a finitely σ -generated constrained σ -field extension of K^ϕ . $\square$

Proof. We set $k = K^\phi$. The assumption $K^{\phi^\infty} = K^\phi$ means that k is relatively algebraically closed in K . We also know that K is separable over k [60, Corollary 1.4.16, p. 16]. Thus, K is a regular field extension of k . Let c be a finite tuple with coordinates in R^ϕ . Then,

$$K\{c\}_\sigma = K \otimes_k k\{c\}_\sigma$$

is an integral domain, because $k\{c\}_\sigma$ is contained in the field R^ϕ and K is regular over k . Moreover, $(0) \subset R$ is a finite intersection of σ -pseudo prime ideals of R by Lemma 2.11. We can thus apply Theorem 2.34 to the inclusion $K\{c\}_\sigma \subset R$ to find $0 \neq r \in K\{c\}_\sigma$ and an integer $l \geq 1$ such that every σ^d -prime ideal $\mathfrak{q}'$ of $K\{c\}_\sigma$ with $r \notin_\sigma \mathfrak{q}'$ lifts to a σ^{ld} -prime ideal of R . We may write

$$r = \lambda_1 \otimes a_1 + \cdots + \lambda_m \otimes a_m \in K \otimes_k k\{c\}_\sigma = K\{c\}_\sigma$$

with the λ_i 's linearly independent over k . Let $b \in k\{c\}_\sigma$ denote one of the nonzero a_i 's. We will show that $k\{c, 1/b\}_\sigma$ has no σ -pseudo prime ideals other than (0) . Let $\mathfrak{q}$ be a σ^d -prime ideal of $k\{c\}_\sigma$ with $b \notin_\sigma \mathfrak{q}$ (for some $d \geq 1$). We have to show that $\mathfrak{q} = (0)$.

Since K is a regular field extension of k , $\mathfrak{q}' := K \otimes \mathfrak{q}$ is a prime ideal of $K \otimes_k k\{c\}_\sigma$. It follows from Lemma 2.36 that $\mathfrak{q}'$ is a σ^d -prime ideal of $K \otimes_k k\{c\}_\sigma$. We claim that $r \notin_\sigma \mathfrak{q}'$. Suppose the contrary. Then, $\sigma^n(r) \in \mathfrak{q}'$ for some $n \geq 1$. By Lemma 2.35, the family $(\sigma^n(\lambda_i))$ is linearly independent over k . By considering the image of $\sigma^n(r)$ in

$$(K \otimes_k k\{c\}_\sigma)/\mathfrak{q}' = K \otimes_k (k\{c\}_\sigma/\mathfrak{q}),$$

we see that this implies $\sigma^n(b) \in \mathfrak{q}$. This contradicts $b \notin_\sigma \mathfrak{q}$. Therefore, $r \notin_\sigma \mathfrak{q}'$. By the construction of r , this implies the existence of a σ^{ld} -prime ideal $\mathfrak{q}''$ of R with

$$\mathfrak{q}'' \cap K\{c\}_\sigma = \mathfrak{q}'.$$

In particular, $\mathfrak{q}'' \supset \mathfrak{q}R$. But, since the elements of $\mathfrak{q}$ are ϕ -constants, $\mathfrak{q}R$ is a ϕ -ideal. Since R is ϕ -simple, we must have $\mathfrak{q}R = (0)$. So, also $\mathfrak{q} = (0)$ as desired.

It remains to see that R^ϕ is finitely generated as a σ -field extension of $k = K^\phi$. Let $\mathfrak{q}$ be a minimal prime ideal of R . Then, there exists $d \geq 1$ such that $\mathfrak{q}$ is ϕ^d -prime and σ^d -prime (Lemma 2.11). Since R is finitely generated as K - σ -algebra, we see that $R/\mathfrak{q}$ is finitely generated as K - σ^d -algebra. So, $\text{Quot}(R/\mathfrak{q})$ is finitely generated as σ^d -field extension of K . As $k = K^{\phi^d}$ by assumption, it follows from Lemma 2.9 that $K \otimes_k R^\phi$ is ϕ^d -simple. Therefore, the canonical map

$$K \otimes_k R^\phi = K \cdot R^\phi \rightarrow \text{Quot}(R/\mathfrak{q})$$

is injective, and we can think of $K R^\phi = \text{Quot}(K \cdot R^\phi)$ as a σ^d -subfield of $\text{Quot}(R/\mathfrak{q})$. By [43, Theorem 4.4.1, p. 292], every intermediate difference field of a finitely generated difference field extension is finitely generated. Therefore, $K R^\phi$ is finitely generated as a σ^d -field extension of K . A fortiori, $K R^\phi$ is finitely generated as σ -field extension of K . We can, therefore, find $a_1, \dots, a_m \in R^\phi$ such that

$$K R^\phi = K \langle a_1, \dots, a_m \rangle_\sigma.$$

So,

$$\text{Quot}(K \otimes_k R^\phi) = \text{Quot}(K \otimes_k k \langle a_1, \dots, a_m \rangle_\sigma).$$

As $K \otimes_k R^\phi$ and $K \otimes_k K^\phi \langle a_1, \dots, a_m \rangle_\sigma$ are ϕ -simple (Lemma 2.9), it follows from Lemma 2.7 that

$$R^\phi = \text{Quot}(K \otimes_k R^\phi)^\phi = \text{Quot}(K \otimes_k k \langle a_1, \dots, a_m \rangle_\sigma)^\phi = k \langle a_1, \dots, a_m \rangle_\sigma. \quad \blacksquare$$

Corollary 2.38. Let K be a $\phi\sigma$ -field and R a σ -PV ring over K with K^ϕ being a σ -closed σ -field. Then $R^\phi = K^\phi$. $\square$

Proof. Since a σ -closed σ -field is algebraically closed and inversive, the hypotheses of Proposition 2.37 are met, and it follows that R^ϕ is a constrained σ -field extension of k .

By [61, Example 2.8, p. 1388], a σ -closed σ -field cannot have a proper constrained σ -field extension. ■

Lemma 2.39. Let K be a $\phi\sigma$ -field and R, R' σ -PV rings over K . Then, there exists a σ -pseudo prime ideal in $R \otimes_K R'$. □

Proof. We begin the proof with a general observation on ϕ -pseudo σ -fields. Let L be a ϕ -pseudo σ -field. Then, L need not be a σ -pseudo field. However, if we write

$$L = e_1 \cdot L \oplus \cdots \oplus e_t \cdot L$$

as after Definition 2.2, then σ -permutes the e_i 's and it follows that L is a finite direct sum (or product) of σ -pseudo fields (cf. Lemma 2.11.) In other words, there are idempotent elements $f_1, \dots, f_m \in L$ such that

$$L = f_1 \cdot L \oplus \cdots \oplus f_m \cdot L,$$

with the $f_i \cdot L$'s σ -pseudo fields. Set $L = \text{Quot}(R)$ and $L' := \text{Quot}(R')$. It suffices to show that there exists a σ -pseudo prime ideal in $L \otimes_K L'$, because a σ -pseudo prime ideal of $L \otimes_K L'$ contracts to a σ -pseudo prime ideal of $R \otimes_K R'$. As above, we can write

$$L = f_1 \cdot L \oplus \cdots \oplus f_m \cdot L \quad \text{and} \quad L' = f'_1 \cdot L' \oplus \cdots \oplus f'_{m'} \cdot L'$$

with the $f_i \cdot L$'s and $f'_j \cdot L'$'s σ -pseudo fields. Then,

$$L \otimes_K L' = \bigoplus_{i,j} f_i \cdot L \otimes_K f'_j \cdot L'.$$

Note that the $f_i \cdot L$'s are finitely σ -generated as σ -pseudo field extensions of K . Indeed, if $Y \in \text{GL}_n(L)$ is a suitable fundamental solution matrix, then $f_i \cdot L = K\langle f_i \cdot Y \rangle_\sigma$. Since $f_1 \cdot L$ is finitely σ -generated over K , it follows from [61, Theorem 1.2, p. 1375] that there exists a σ -pseudo prime ideal $\mathfrak{p}$ in $f_1 \cdot L \otimes_K f'_1 \cdot L'$. Then,

$$\tilde{\mathfrak{p}} := \mathfrak{p} \bigoplus_{\substack{i,j \\ (i,j) \neq (1,1)}} f_i \cdot L \otimes_K f'_j \cdot L'$$

is a σ -pseudo prime ideal of $L \otimes_K L'$. ■

Finally, we are prepared to prove our main uniqueness theorem.

Theorem 2.40 (Uniqueness of σ -PV rings). Let K be a $\phi\sigma$ -field such that $K^\phi = K^{\phi^\infty}$ and $\sigma: K^\phi \rightarrow K^\phi$ is surjective. Let R_1 and R_2 be two σ -PV rings over K for the same equation $\phi(y) = Ay$, $A \in \mathrm{GL}_n(K)$. Then, there exists a finitely σ -generated constrained σ -pseudo field extension k' of $k := K^\phi$ containing $k_1 := R_1^\phi$ and $k_2 := R_2^\phi$ and an isomorphism of $K \otimes_k k'$ - $\phi\sigma$ -algebras between $R_1 \otimes_{k_1} k'$ and $R_2 \otimes_{k_2} k'$. $\square$

Proof. We know from Proposition 2.37 that k_1 and k_2 are finitely σ -generated constrained σ -field extensions of k . Let $Y_1 \in \mathrm{GL}_n(R_1)$ and $Y_2 \in \mathrm{GL}_n(R_2)$ be fundamental solution matrices for $\phi(y) = Ay$. Set

$$Z = (Y_1 \otimes 1)^{-1} (1 \otimes Y_2) \in \mathrm{GL}_n(R_1 \otimes_K R_2).$$

As noted in Remark 2.17, we have

$$Z \in \mathrm{GL}_n((R_1 \otimes_K R_2)^\phi).$$

Since $1 \otimes Y_2 = (Y_1 \otimes 1) \cdot Z$, the entries of $1 \otimes Y_2$ lie in

$$R_1 \cdot S_Z, \quad S_Z := k_1 \{Z, 1/\det(Z)\}_\sigma \subset R_1 \otimes_K R_2.$$

Using Lemma 2.10, it follows that

$$R_1 \otimes_K R_2 = R_1 \cdot S_Z = R_1 \otimes_{k_1} S_Z.$$

Our next goal is to find a k_1 - σ -morphism $\psi: S_Z \rightarrow k'$ for some finitely σ -generated constrained σ -pseudo field extension k' of k_1 . We know from Lemma 2.39 that there exists a σ -pseudo prime ideal in $R_1 \otimes_K R_2$. This σ -pseudo prime ideal contracts to a σ -pseudo prime ideal of S_Z . We can thus apply [61, Proposition 2.12, p. 1390] to find a maximal element $\mathfrak{p}$ in the set of all σ -pseudo prime ideals of S_Z ordered by inclusion. By [61, Proposition 2.9, p. 1389], the residue σ -pseudo field

$$k' := \mathrm{Quot}(S_Z/\mathfrak{p})$$

is a constrained σ -pseudo field extension of k . Moreover, we have a natural k_1 - σ -morphism $\psi: S_Z \rightarrow k'$. Then,

$$\varphi: R_2 \rightarrow R_1 \otimes_K R_2 = R_1 \otimes_{k_1} S_Z \xrightarrow{\mathrm{id} \otimes \psi} R_1 \otimes_{k_1} k'$$

is a morphism of K - $\phi\sigma$ -algebras. Since $(R_1 \otimes_{k_1} k')^\phi = k'$, this yields an embedding of k_2 into k' , and we can extend φ to a $K \otimes_k k'$ - $\phi\sigma$ -morphism

$$\varphi: R_2 \otimes_{k_2} k' \rightarrow R_1 \otimes_{k_1} k'.$$

As $\varphi(Y_2)$ and Y_1 are fundamental solution matrices in $R_1 \otimes_{k_1} k'$ for $\phi(y) = Ay$, there exists $C \in \text{GL}_n(k')$ such that

$$Y_1 = \varphi(Y_2)C = \varphi(Y_2C).$$

Since R_1 is σ -generated by Y_1 , this shows that φ is surjective. Now $R_2 \otimes_{k_2} k'$ need not be ϕ -simple. However, by Lemma 2.8, every ϕ -ideal of $R_2 \otimes_{k_2} k'$ is of the form $R_2 \otimes_{k_2} \mathfrak{b}$ for some ideal $\mathfrak{b}$ of k' . Since the kernel of φ is a ϕ -ideal, this implies that φ is injective. ■

Lemma 2.41. Let K be a ϕ -pseudo σ -field and R a σ -PV ring over K with $R^\phi = K^\phi =: k$. Then,

$$R \otimes_K R = R \otimes_k (R \otimes_K R)^\phi. \quad \square$$

Proof. This follows as in the beginning of the proof of Theorem 2.40 (with $R_1 = R_2 = R$). ■

Corollary 2.42 (Uniqueness of σ -PV extensions). Let K be a $\phi\sigma$ -field and let L_1, L_2 be two σ -PV extensions for the same equation $\phi(y) = Ay$, $A \in \text{GL}_n(K)$. Assume that K^ϕ is σ -closed. Then, there exists an integer $l \geq 1$ and an isomorphism of K - $\phi\sigma^l$ -algebras between L_1 and L_2 . □

Proof. Let $R_1 \subset L_1$ and $R_2 \subset L_2$ denote the corresponding σ -PV rings. As usual, we set $k := K^\phi$. We have $R_1^\phi = k$ and $R_2^\phi = k$. By Theorem 2.40, there exists a finitely σ -generated constrained σ -pseudo field extension k' of k and an isomorphism

$$\varphi: R_1 \otimes_k k' \rightarrow R_2 \otimes_k k'$$

of $K \otimes_k k'$ - $\phi\sigma$ -algebras. But, by [61, Example 2.8, p. 1388], every finitely σ -generated constrained σ -pseudo field extension of a σ -closed σ -field is trivial. This means that there

exists an integer $l \geq 1$ such that k' is of the form

$$k' = k \oplus \cdots \oplus k$$

with σ given by

$$\sigma(a_1 \oplus \cdots \oplus a_l) = \sigma(a_l) \oplus \sigma(a_1) \oplus \cdots \oplus \sigma(a_{l-1}).$$

Let $\mathfrak{a}$ be a maximal ideal of k' . Then, $\mathfrak{a}$ is a σ^l -ideal with $k'/\mathfrak{a} = k$ as σ^l -rings. For $i = 1, 2$, the ideal $R_i \otimes_k \mathfrak{a}$ is a $\phi\sigma^l$ -ideal of $R_i \otimes_k k'$, and φ is mapping $R_1 \otimes_k \mathfrak{a}$ bijectively onto $R_2 \otimes_k \mathfrak{a}$. Passing to the quotient, we obtain an isomorphism

$$\bar{\varphi}: (R_1 \otimes_k k')/(R_1 \otimes_k \mathfrak{a}) \rightarrow (R_2 \otimes_k k')/(R_2 \otimes_k \mathfrak{a})$$

of $\phi\sigma^l$ -rings. But

$$R_i \rightarrow R_i \otimes_k k' \rightarrow (R_i \otimes_k k')/(R_i \otimes_k \mathfrak{a}) = R_i \otimes_k (k'/\mathfrak{a}) = R_i$$

identifies R_i with $(R_i \otimes_k k')/(R_i \otimes_k \mathfrak{a})$ as $\phi\sigma^l$ -ring. So, we have constructed a K - $\phi\sigma^l$ -isomorphism between R_1 and R_2 . Finally, this isomorphism extends to the total quotient rings, that is, to the σ -PV extensions. ■

Let K be a $\phi\sigma$ -field and $A \in \mathrm{GL}_n(K)$. Even if, in all generality, a σ -PV extension for $\phi(y) = Ay$ need not be unique, the following remark shows that, in some situations, it is possible to make a more or less canonical choice. For example, if $K = k(z)$ as in Section 2.5, then the σ -PV ring for $\phi(y) = Ay$ inside Seq_k is unique (as a subring of Seq_k).

Remark 2.43. Let K be a $\phi\sigma$ -field and $A \in \mathrm{GL}_n(K)$. Let S be a K - $\phi\sigma$ -algebra with $S^\phi = K^\phi$. If there exists a σ -PV ring R for $\phi(y) = Ay$ in S , then R is unique in the sense that any other σ -PV ring for $\phi(y) = Ay$ in S equals R . □

Proof. Let R' be another σ -PV ring for $\phi(y) = Ay$ inside S . As R and R' are σ -generated by appropriate fundamental solution matrices, it follows from Remark 2.17 and the fact that $S^\phi \subset K$ that $R' = R$. ■

2.7 σ -Galois group and Galois correspondence

In this section, we will define the σ -Galois group of $\phi(y) = Ay$ (Definition 2.50), show that it is a σ -algebraic group (Lemma 2.51), establish the Galois correspondence

(Theorem 2.52), and finish by showing that the σ -dimension, introduced in [23], of the σ -Galois group coincides with the σ -dimension of a σ -PV ring of the equation (Lemma 2.53), which we will further use in our applications, Theorems 3.1 and 3.5. For this, we first recall what a σ -algebraic group is using the language of σ -Hopf algebras (and representable functors). See the appendix of [23] for a brief introduction to σ -algebraic groups.

Throughout Sections 2.7 and 2.8, we will make the following assumptions. Let K be a ϕ -pseudo σ -field and $k := K^\phi$ its σ -field of ϕ -constants. Assume that there exists a σ -PV ring R for the linear ϕ -equation $\phi(y) = Ay$, $A \in \mathrm{GL}_n(K)$, with $R^\phi = k$, and let $L = \mathrm{Quot}(R)$ be the corresponding σ -PV extension (cf. Proposition 2.21). The category of k - σ -algebras is denoted by $\mathrm{Alg}_{k-\sigma}$.

Definition 2.44. A k - σ -Hopf algebra is a Hopf algebra over k in which the comultiplication Δ , antipode S , and counit ε are k - σ -algebra homomorphisms. $\square$

Definition 2.45. A k - σ -algebraic group is a functor $G : \mathrm{Alg}_{k-\sigma} \rightarrow \mathrm{Sets}$ represented by a k - σ -Hopf algebra H , which is finitely σ -generated over k . That is, for every $B \in \mathrm{Alg}_{k-\sigma}$,

$$G(B) = \mathrm{Hom}_{k-\sigma}(H, B).$$

For simplicity, we say that H represents G . $\square$

In other words, a k - σ -algebraic group is a group object in the category of σ -algebraic k - σ -schemes (in the sense of [23, Definition A.1]).

Definition 2.46 ([23, Definition A.37]). A k - σ -algebraic group G' is called a k - σ -subgroup of a k - σ -algebraic group G if $G'(B)$ is a subgroup of $G(B)$ for every k - σ -algebra B . $\square$

Proposition 2.47 ([23, Remark A.38]). For every k - σ -algebraic subgroup G' of a k - σ -algebraic group G represented by H , there exists a σ -Hopf ideal I in H such that G' is represented by H/I and vice versa. $\square$

The multiplicative k - σ -algebraic group $\mathbf{G}_m$ is the k - σ -algebraic group represented by $k\{x, 1/x\}_\sigma$ with $\Delta(x) = x \otimes x$, $S(x) = 1/x$, and $\varepsilon(x) = 1$.

Proposition 2.48 ([23, Lemma A.40]). For every σ -Hopf ideal I of $H := k\{x, 1/x\}_\sigma$ with the above Hopf algebra structure, there exists a multiplicative function $\varphi = x^{n_0} \cdot \sigma(x)^{n_1} \cdot \dots \cdot \sigma^t(x)^{n_t} \in H$ such that I contains $\varphi - 1$. $\square$

Lemma 2.49. The k - σ -algebra

$$H := (R \otimes_K R)^\phi$$

is a k - σ -Hopf algebra via the $\phi\sigma$ - R -bimodule structure on $C := R \otimes_K R$ (see [3, (1.5, 1.6)]):

$$\begin{aligned} \Delta : C &\rightarrow C \otimes_R C, & \Delta(a \otimes b) &= a \otimes 1 \otimes b \in R \otimes_K R \otimes_K R \cong R \otimes_K R \otimes_R R \otimes_K R, \\ \varepsilon : C &\rightarrow R, & \varepsilon(a \otimes b) &= ab, \end{aligned}$$

and the K - $\phi\sigma$ -linear flip homomorphism $\tau : C \rightarrow C$, $\tau(a \otimes b) = b \otimes a$. Moreover,

$$\mu : R \otimes_k H \rightarrow R \otimes_K R, \quad r \otimes h \mapsto (r \otimes 1) \cdot h \quad (2.7)$$

is an isomorphism of K - $\phi\sigma$ -algebras. $\square$

Proof. The proof is a modification of the proof of [3, Proposition 1.7] and [2, Proposition 3.4]. We already noted in Lemma 2.41 that (2.7) is an isomorphism. It follows that the K - $\phi\sigma$ -algebra homomorphism

$$R \otimes_k H \otimes_k H \xrightarrow{\mu \otimes \text{id}} R \otimes_K R \otimes_k H \xrightarrow{\text{id} \otimes \mu} R \otimes_K R \otimes_K R$$

is an isomorphism. By taking ϕ -constants, we, therefore, obtain a k - σ -algebra isomorphism

$$H \otimes_k H \rightarrow (R \otimes_K R \otimes_K R)^\phi. \quad (2.8)$$

To show that, given the above, H becomes a k - σ -Hopf algebra, one proceeds as in the proof of [3, Proposition 1.7]. $\blacksquare$

Definition 2.50. Let R and L be as above. Then, the σ -Galois group of L over K is defined as the functor

$$\text{Gal}^\sigma(L|K) : \text{Alg}_{k-\sigma} \rightarrow \text{Sets}, \quad B \mapsto \text{Gal}^\sigma(L|K)(B) := \text{Aut}_{\phi\sigma}(R \otimes_k B | K \otimes_k B),$$

where ϕ acts as the identity on B . $\square$

Lemma 2.51. Let R , L , and H be as above. Then, $G := \text{Gal}^\sigma(L|K)$ is a k - σ -algebraic group represented by H . $\square$

Proof. As in the proof of [3, Lemma 1.9], R is an H -comodule via

$$\theta : R \rightarrow R \otimes_k H, \quad r \mapsto \mu^{-1}(1 \otimes r),$$

which is a K - $\phi\sigma$ -algebra homomorphism, where μ is defined in (2.7). For every k - σ -algebra B and $g \in \text{Hom}_{k-\sigma}(H, B)$, we have a K - $\phi\sigma$ -algebra homomorphism

$$\Phi_g : R \otimes_k B \xrightarrow{\theta \otimes \text{id}_B} R \otimes_k H \otimes_k B \xrightarrow{\text{id}_R \otimes g \otimes \text{id}_B} R \otimes_k B \otimes_k B \xrightarrow{\text{id}_R \otimes m} R \otimes_k B,$$

which is an automorphism by [59, Theorem 3.2]. Moreover, by [59, Theorem 3.2] as well, the map $g \mapsto \Phi_g$ is a group homomorphism. For the reverse direction, let $Y \in \text{GL}_n(R)$ be a fundamental solution matrix of $\phi(Y) = AY$ and $Z = (Y \otimes 1)^{-1}(1 \otimes Y) \in \text{GL}_n(R \otimes_K R)$. Then, $H = k\{Z, 1/\det(Z)\}_\sigma$ and it follows from Remark 2.17 that, for any

$$\varphi \in \text{Aut}_{\phi\sigma}(R \otimes_k B | K \otimes_k B),$$

there exists $C_\varphi \in \text{GL}_n(B)$ such that $\varphi(Y) = Y \cdot C_\varphi$. We define a k - σ -algebra homomorphism $H \rightarrow B$ by sending Z to C_φ . ■

Theorem 2.52. There is a one-to-one correspondence between k - σ -algebraic subgroups in G and intermediate ϕ -pseudo σ -fields in $L|K$ given by

$$M = L^{G'} := \{a/b \in L \mid \theta'(a) \cdot b = a \cdot \theta'(b), \quad a, b \in R\} \longleftrightarrow G' := \text{Gal}^\sigma(L|M), \quad (2.9)$$

or, alternatively,

$$M = L^{G'} := \{x \in L \mid \text{for all } B \in \text{Alg}_{k-\sigma}, \quad g \in G'(B), \quad g(x \otimes 1) = x \otimes 1\} \longleftrightarrow G' := \text{Gal}^\sigma(L|M), \quad (2.10)$$

where $\theta' : R \rightarrow R \otimes_k H'$, and H' represents G' . □

Proof. We will show that there is a one-to-one correspondence between the σ -Hopf ideals in H and intermediate ϕ -pseudo σ -fields in $L|K$ given by

$$M = \{x \in L \mid 1 \otimes x - x \otimes 1 \in I \cdot (L \otimes_K L)\} \longleftrightarrow I = H \cap \ker(L \otimes_K L \rightarrow L \otimes_M L).$$

The proof below is partly an adaptation of [3, Proposition 2.3]. It follows from [60, Theorem 3.1.17] that there is a one-to-one correspondence between the intermediate

ϕ -pseudo σ -fields in $L|K$ and $\phi\sigma$ -coideals of $L \otimes_K L$ given by

$$M = \{x \in L \mid 1 \otimes x - x \otimes 1 \in J \subset L \otimes_K L\} \longleftrightarrow J = \ker(L \otimes_K L \rightarrow L \otimes_M L).$$

By Lemma 2.8, there is a one-to-one correspondence between $\phi\sigma$ -ideals of $L \otimes_k H$ and σ -ideals of H given by

$$\mathfrak{b} = \mathfrak{a} \cap H \longleftrightarrow \mathfrak{a} = L \otimes_k \mathfrak{b}. \quad (2.11)$$

By localizing (2.7), we obtain K - $\phi\sigma$ -algebra isomorphisms

$$\varphi_1 : L \otimes_k H \rightarrow L \otimes_K R \quad \text{and} \quad \varphi_2 : L \otimes_k H \rightarrow R \otimes_K L. \quad (2.12)$$

Therefore, we have a one-to-one correspondence between σ -ideals of H and $\phi\sigma$ -ideals of $L \otimes_K L$ given by composing (2.12) and (2.11) and using the fact that the set of ideals of the localization $L \otimes_K L$ consists of the intersection of the set of ideals in the smaller localizations $L \otimes_K R$ and $R \otimes_K L$ inside the set of ideals in $R \otimes_K R$.

We will now show that, under the above correspondence and in the above notation, $(L \otimes_K L) \cdot \mathfrak{a}$ is a $\phi\sigma$ -coideal of $L \otimes_K L$ if and only if $\mathfrak{b}$ is a σ -Hopf ideal of H . For this, note that, similarly to the above, we have a one-to-one correspondence between ideals in $H \otimes_k H$ and ϕ -ideals in $L \otimes_K L \otimes_K L$. Indeed, by Lemma 2.8 and isomorphisms (2.8) and (2.12), there is a one-to-one correspondence between $\phi\sigma$ -ideals of $L \otimes_K R \otimes_K R$ (as well as those in $R \otimes_K L \otimes_K R$ and $R \otimes_K R \otimes_K L$) and σ -ideals of $H \otimes_k H$ with

$$H \otimes_k \mathfrak{b} \longleftrightarrow \mathfrak{a} = L \otimes_K \mathfrak{b} \quad \text{and} \quad \mathfrak{b} \otimes_k H \longleftrightarrow \mathfrak{a} = \mathfrak{b} \otimes_K L,$$

therefore,

$$\mathfrak{b}_1 := H \otimes_k \mathfrak{b} + \mathfrak{b} \otimes_k H \longleftrightarrow ida_1 := L \otimes_K \mathfrak{a} + \mathfrak{a} \otimes_K L$$

under the correspondence $\mathfrak{a} \subset L \otimes_K L \leftrightarrow \mathfrak{b} \subset H$ from the preceding paragraph. Therefore,

$$\Delta(\mathfrak{a}) \subset \mathfrak{a}_1, \quad \varepsilon(\mathfrak{a}) = 0 \iff \Delta(\mathfrak{b}) \subset \mathfrak{b}_1, \quad \varepsilon(\mathfrak{b}) = 0.$$

By [47, Theorem 1(iv)], $\mathfrak{b}$ is a Hopf ideal of H if and only if $\mathfrak{b}$ is a coideal of H , which finishes the proof. To show correspondence (2.9), note that, by Lemma 2.51, $\text{Gal}^\sigma(L|M)$ is

represented by

$$H/H \cap \ker(L \otimes_K L \rightarrow L \otimes_M L).$$

Therefore, it remains to show that

$$\begin{aligned} L_1 &:= \{x \in L \mid 1 \otimes x - x \otimes 1 \in \ker(L \otimes_K L \rightarrow L \otimes_M L)\} \\ &= L_2 := \{a/b \in L \mid \theta'(a) \cdot b = a \cdot \theta'(b), \ a, b \in R\} = \\ &= L_3 := \{x \in L \mid \text{for all } B \in \text{Alg}_{k-\sigma}, \ g \in G'(B), \ g(x \otimes 1) = x \otimes 1\}. \end{aligned}$$

For every $x = a/b \in L_2$, $B \in \text{Alg}_{k-\sigma}$, and $g \in G'(B)$, we have

$$g(a/b \otimes 1) = (\theta'(a) \cdot b \otimes 1)(g) / (\theta'(b) \cdot b \otimes 1)(g) = (\theta'(b) \cdot a \otimes 1)(g) / (\theta'(b) \cdot b \otimes 1)(g) = a/b \otimes 1.$$

Hence, $x \in L_3$. Now, for all $x = a/b \in L_3$, we have $\theta'(a) \cdot b = a \cdot \theta'(b)$ by taking $B := H$ and $g := \text{id}_H$. Therefore, $L_2 = L_3$. For $L_1 = L_3$, see the proof of [60, Lemma 3.1.11]. ■

For σ -dimension, see [23, Section A.7]. Let K be a $\phi\sigma$ -field and R , L , and H be as above.

Lemma 2.53. We have

$$\sigma\text{-dim}_K R = \sigma\text{-dim}_k H. \quad \square$$

Proof. Let $Y \in \text{GL}_n(R)$ be a fundamental solution matrix of $\phi(y) = Ay$ and $Z = (Y \otimes 1)^{-1} (1 \otimes Y) \in \text{GL}_n(R \otimes_K R)$. Then,

$$R = K\{Y, 1/\det(Y)\}_\sigma \quad \text{and} \quad H = k\{Z, 1/\det(Z)\}_\sigma.$$

The claim now follows from [23, Definition A.25], Lemma 2.20, and [58, Theorem 1.13]. ■

2.8 Isomonodromic difference equations

In this section, we develop a σ -Galois treatment for isomonodromic difference equations. In particular, in Theorem 2.55, not assuming that the field $k = K^\phi$ is difference closed, we give a criterion, which says that $\phi(y) = Ay$ is isomonodromic if and only if the matrices in its σ -Galois group all satisfy an equation of a special form (2.14). This

result is a difference analog of the corresponding results for isomonodromic differential equations, [10, Proposition 3.9; 30, Theorem 6.6], and can be combined with [48, Theorem 4.1] to study difference isomonodromy of linear difference equations with several parameters. We further illustrate this by considering a q -hypergeometric equation in Example 2.56.

Definition 2.54. The system $\phi(y) = Ay$ is called *isomonodromic* if there exists $B \in \mathrm{GL}_n(K)$ such that

$$\phi(B)AB^{-1} = \sigma(A). \quad (2.13)$$

□

Theorem 2.55. The equation $\phi(y) = Ay$ is isomonodromic if and only if there exists $D \in \mathrm{GL}_n(k)$ such that the following equation is in the defining ideal of the σ -Galois group G :

$$\sigma(x_{ij}) = D^{-1}(x_{ij})D. \quad (2.14)$$

Moreover, if (2.14) is in the defining ideal of G , then there exists a finitely generated σ -field extension F of k and $C \in \mathrm{GL}_n(F)$ such that

$$\sigma(C^{-1}(x_{ij})C) = C^{-1}(x_{ij})C \quad (2.15)$$

is in the defining ideal of G , that is, G is conjugate over F to a group of matrices with σ -constant entries. □

Proof. Let $Y \in \mathrm{GL}_n(R)$ be a fundamental solution matrix. Let $B \in \mathrm{GL}_n(K)$ be such that (2.13) is satisfied. We have

$$\phi(\sigma(Y)^{-1}BY) = \sigma(\phi(Y))^{-1}\phi(B)\phi(Y) = \sigma(AY)^{-1}\sigma(A)BA^{-1}AY = \sigma(Y)^{-1}BY.$$

Therefore, there exists $D \in \mathrm{GL}_n(k)$ such that $\sigma(Y) = BYD$. For every k - σ -algebra S and $g \in G(S)$, let $C_g \in \mathrm{GL}_n(S)$ be such that $g(Y) = YC_g$. Then, on the one hand,

$$g(\sigma(Y)) = g(BYD) = BYC_gD.$$

On the other hand,

$$g(\sigma(Y)) = \sigma(g(Y)) = \sigma(YC_g) = \sigma(Y)\sigma(C_g) = BYD\sigma(C_g).$$

Therefore, for all $g \in G(S)$, we have

$$\sigma(C_g) = D^{-1}C_gD,$$

showing (2.14). To show (2.15), let F be a σ -field generated over k by the entries of an invertible matrix C satisfying $\sigma(C) = D^{-1}C$. Then,

$$\sigma(C^{-1}C_gC) = \sigma(C)^{-1}\sigma(C_g)\sigma(C) = C^{-1}DD^{-1}C_gDD^{-1}C = C^{-1}C_gC.$$

Suppose now that, for all k - σ -algebras S and $g \in G(S)$, we have $\sigma(C_g) = D^{-1}C_gD$, where $C_g := Y^{-1}g(Y)$. Let $B := \sigma(Y)D^{-1}Y^{-1}$. Then, for all $g \in G$,

$$g(B) = \sigma(YC_g)D^{-1}(YC_g)^{-1} = \sigma(Y)D^{-1}C_gDD^{-1}C_g^{-1}Y^{-1} = B.$$

By Theorem 2.52, $B \in \text{GL}_n(K)$. We, moreover, have

$$\phi(B) = \phi(\sigma(Y))D^{-1}\phi(Y)^{-1} = \sigma(AY)D^{-1}(AY)^{-1} = \sigma(A)\sigma(Y)D^{-1}Y^{-1}A^{-1} = \sigma(A)BA^{-1},$$

showing (2.13). ■

Example 2.56. Consider a q -hypergeometric equation

$$y(q^2x) - \frac{2ax - 2}{a^2x - 1}y(qx) + \frac{x - 1}{a^2x - 1}y(x) = 0. \quad (2.16)$$

It is shown in [51] that, over $\mathbb{C}(x)$, if $a \notin q^{\mathbb{Z}}$, then, if $a^2 \notin q^{\mathbb{Z}}$, then the difference Galois group of (2.16) is $\text{GL}_2(\mathbb{C})$, otherwise it is $\text{SL}_2(\mathbb{C})$. Equation (2.16) has been also studied from the differential-parametric viewpoint in [36, Example 3.14]. Let now $\mathcal{C}$ be any field such that (2.16) has a σ -PV extension over $\mathcal{C}(x, a)$, with a being transcendental over $\mathcal{C}(x, a)$, ϕ and σ acting as id on $\mathcal{C}$, and

$$\phi(x) = qx, \quad \phi(a) = a, \quad \sigma(x) = x, \quad \sigma(a) = qa.$$

The existence can be shown as in Proposition 2.30. A calculation in MAPLE, similar to the one given in [35], but using the procedure `RationalSolution` in the `QDifferenceEquations` package, shows that (2.16), once transformed into the matrix

form, is isomonodromic over $\mathcal{C}(x, a)$ with

$$B = \begin{pmatrix} \frac{1}{a^2x - 1} & -\frac{2a}{(a+1)(a^2x - 1)} \\ \frac{2a(x-1)}{(a+1)(a^2x - 1)(a^2qx - 1)} & \frac{3a - 1 + (a^3 - 3a^2)x}{(a+1)(a^2x - 1)(a^2qx - 1)} \end{pmatrix}.$$

Therefore, (2.13) is in the defining ideal of the σ -PV group G of (2.16) by Theorem 2.55. It follows from [4, Corollary 3.3.2.1; 51, Theorem 10] that the (non- σ -parametric) PV group of (2.16) over $\mathcal{C}(x, a)$ is GL_2 . Similarly to [36, Proposition 6.21], it follows from Theorem 2.52 that G is Zariski dense in GL_2 . It follows from Theorem 2.55 that, G is conjugate to $\mathrm{GL}_2(\mathcal{C})$ over a (proper, as `RationalSolution` shows) finitely generated σ -field extension of $\mathcal{C}(a)$, where $\mathrm{GL}_2(\mathcal{C})$ is defined by $\mathrm{GL}_2(\mathcal{C})(B) = \{g \in \mathrm{GL}_2(B) \mid \sigma(g) = g\}$ for every $\mathcal{C}(a)$ - σ -algebra B . $\square$

3 Applications and examples

In this section, we will illustrate how our Galois theory can be used to study difference and differential algebraic properties of functions. We start by showing a general σ -independence criterion in Theorem 3.1 (see also [5, Theorem 4.1]). In Section 3.1, we show a σ -independence criterion over the field of meromorphic function with Nevanlinna growth order < 1 (Theorem 3.5). For this, we need some preparatory work, Lemmas 3.2 and 3.4, which are interesting on their own, as they generalize a natural modification of a classical result in complex analysis [6]. We then show how to apply our results in practice in Theorem 3.6, which is followed by illustrative examples in Section 3.4.

3.1 General result

Theorem 3.1. Let F be a $\phi\sigma$ -field containing the field $\mathbb{C}(z)$ with

$$\begin{aligned} \phi(z) &= a_1z + a_2, \quad \sigma(z) = b_1z + b_2, \quad a_1, a_2, b_1, b_2 \in \mathbb{C}, \quad a_1b_1 \neq 0, \\ \phi\sigma &= \sigma\phi, \quad \phi^n \neq \mathrm{id}, \quad n \in \mathbb{N}, \end{aligned} \tag{3.1}$$

and $k := F^\phi$. Let $0 \neq f \in F$ and $0 \neq a \in \mathbb{C}(z)$ be such that f is a solution of

$$\phi(y) = ay. \tag{3.2}$$

Then, f is σ -algebraically dependent over the field $\mathbf{k}(z)$ if and only if

$$\varphi(a) = \phi(b)/b \quad (3.3)$$

for some $0 \neq b \in \mathbb{C}(z)$ and $\varphi(x) \in \mathbb{Q}\{x, 1/x\}_\sigma$, $1 \neq \varphi(x) = x^{n_0} \sigma(x)^{n_1} \cdot \dots \cdot \sigma^{t-1}(x)^{n_{t-1}}$. $\square$

Proof. If (3.3) holds, then

$$\phi(\varphi(f)/b) = \varphi(\phi(f))/\phi(b) = \varphi(af)/\phi(b) = \varphi(a)\varphi(f)/\phi(b) = \varphi(f)/b.$$

Therefore, $\varphi(f)/b = c \in F^\phi = \mathbf{k}$. Thus, $\varphi(f) = c \cdot b \in \mathbf{k}(z)$, which gives a σ -algebraic dependence for f over $\mathbf{k}(z)$.

Assume now that f is σ -algebraically dependent over $\mathbf{k}(z)$. Let L be the smallest $\phi\sigma$ -subfield in F containing $\mathbf{k}(z)$ and f . Since $\mathbf{k} \subset L^\phi \subset F^\phi = \mathbf{k}$, the $\phi\sigma$ -field L is a σ -PV extension over $\mathbf{k}(z)$ for Equation (3.2). It follows from Lemma 2.53 that f is σ -algebraically dependent over $\mathbf{k}(z)$ if and only if the σ -Galois group G of $L|K$ is a proper σ -algebraic subgroup of $\mathbf{G}_m$. Then, by Proposition 2.48, there exists a multiplicative $\varphi \in \mathbf{k}\{x, 1/x\}_\sigma$ such that the ideal of G contains the equation $\varphi(x) = 1$. Therefore, for every $\mathbf{k}$ - σ -algebra B and $g \in G(B)$, we have

$$g(\varphi(f)) = \varphi(g(f)) = \varphi(c_g \cdot f) = \varphi(c_g) \cdot \varphi(f) = 1 \cdot \varphi(f) = \varphi(f).$$

Hence, by Theorem 2.52, we have $b := \varphi(f) \in \mathbf{k}(z)$. Since $f \neq 0$ and φ is multiplicative, $\varphi(f) \neq 0$. Therefore,

$$\varphi(a) = \varphi(\phi(f)/f) = \phi(\varphi(f))/\varphi(f) = \phi(b)/b. \quad (3.4)$$

We will show now that b can be chosen from $\mathbb{C}(z)$ satisfying (3.3). For this, first note that z is transcendental over $\mathbf{k}$. Indeed, for all $n \in \mathbb{N}$ and $a_0, \dots, a_n \in \mathbf{k}$,

$$a_n z^n + \dots + a_1 z + a_0 = 0$$

implies that, for all $q \in \mathbb{N}$,

$$a_n(\phi^q(z))^n + \dots + a_1(\phi^q(z)) + a_0 = 0.$$

This implies that there exists $r \in \mathbb{N}$ such that $z = \phi^r(z)$, which contradicts (3.1). Now, we have the equalities $a = \bar{a}/c$ and $b = \bar{b}/d$, where $\bar{a}, c \in \mathbb{C}[z]$ and $\bar{b}, d \in \mathbf{k}[z]$. Consider the

coefficients of $\bar{b}$ and d with respect to z as new indeterminates. Equation (3.4) is equivalent to

$$\varphi(\bar{a}/c) = \phi(\bar{b}/d)/(\bar{b}/d).$$

So, we have

$$\varphi(\bar{a}) \cdot \phi(d) \cdot \bar{b} - \varphi(c) \cdot \phi(\bar{b}) \cdot d = 0. \quad (3.5)$$

The left-hand side of Equation (3.5) is a polynomial in z . Hence, Equation (3.4) can be considered as a system of polynomial equations given by the equalities for all coefficients. Since the field $\mathbb{C}$ is algebraically closed, existence of $\bar{b}$ and d with coefficients in $\mathbf{k}$ implies existence of $\bar{b}$ and d with coefficients in $\mathbb{C}$. ■

3.2 Meromorphic functions and Nevanlinna property

Let M be the $\phi\sigma$ -field of meromorphic functions on the plane with

$$\phi(f)(z) := f(z+1), \quad \sigma(f)(z) := f(z+a_\sigma), \quad f \in M, \quad z, a_\sigma \in \mathbb{C}.$$

Also, let $\mathbf{k} := M^\phi$, which is the field of 1-periodic meromorphic functions. For $f \in M$, the standard Nevanlinna characteristics $m(r, f)$, $N(r, f)$, and $T(r, f)$ were introduced in [46, pp. 6, 12] (see also [6, 14, 29]). Let

$$M_{<1} := \{g \in M \mid T(r, g) = o(r), \ r \rightarrow +\infty\}, \quad (3.6)$$

which is a $\phi\sigma$ -field as well [6, 8. Proposition]. Note that

$$\mathbb{C}(z) \subsetneq M_{<1}. \quad (3.7)$$

The proof of the following result, which we need to prove Theorem 3.5, was suggested by D. Drasin and S. Merenkov, to whom the authors are highly grateful, as a modification of [6, 7. Lemma (c)].

Lemma 3.2. Let $f \in M$ and there exist $R \in \mathbb{C}(z)$ such that, for all $z \in \mathbb{C}$,

$$f(z+1) = R(z) \cdot f(z). \quad (3.8)$$

If $f \in M_{<1}$, then $f \in \mathbb{C}(z)$. □

Proof. Let $L > 0$ be a real number such that all finite poles and zeroes of R lie in

$$D(L) := \{c \in \mathbb{C} \mid |c| < L\}.$$

Similarly to the proof of [6, 7. Lemma (c)], one shows that (3.8) and (3.6) imply that all finite poles and zeroes of f lie in $D(L)$. This implies that there exists a rational function h such that $g := hf$ is an entire function with no zeroes. Since $M_{<1}$ is a field and $h \in M_{<1}$, we have $g = hf \in M_{<1}$. Hence, it follows from [29, Lemma I.6.2] that g is constant. Therefore, $f = g/h$ is rational. ■

Corollary 3.3. We have

$$\mathbf{k} \cap M_{<1} = \mathbb{C}. \quad \square$$

We will need one more complex-analytic result (which has an algebraic proof) to prove Theorem 3.5 as well.

Lemma 3.4. Let $a \in \mathbb{C}(z) \setminus \{0\}$. Assume that there exists a nonzero $b \in \mathbf{k} M_{<1}$ such that $\phi(b) = ab$. Then,

- (i) there also exists a nonzero $b' \in M_{<1}$ with $\phi(b') = ab'$;
- (ii) $b \in \mathbf{k}(z)$. □

Proof. We know from Corollary 3.3 that $M_{<1}^\phi = \mathbb{C}$, and it follows from Lemma 2.10 that $M_{<1}$ is linearly disjoint from $\mathbf{k}$ over $\mathbb{C}$. Hence,

$$\mathbf{k} M_{<1} = \text{Quot}(M_{<1} \otimes_{\mathbb{C}} \mathbf{k}). \quad (3.9)$$

Moreover, $M_{<1} \otimes_{\mathbb{C}} \mathbf{k}$ is ϕ -simple by Lemma 2.9. We will first show that b must lie in $M_{<1} \otimes_{\mathbb{C}} \mathbf{k}$. Set

$$\mathfrak{a} = \{f \in M_{<1} \otimes_{\mathbb{C}} \mathbf{k} \mid f \cdot b \in M_{<1} \otimes_{\mathbb{C}} \mathbf{k}\}.$$

It follows from (3.9) that $\mathfrak{a}$ is a nonzero ideal of $M_{<1} \otimes_{\mathbb{C}} \mathbf{k}$. For all $f \in \mathfrak{a}$, we have $\phi(fb) \in M_{<1} \otimes_{\mathbb{C}} \mathbf{k}$ and, therefore,

$$\phi(fb) = \phi(f) \cdot ab \in M_{<1} \otimes_{\mathbb{C}} \mathbf{k}.$$

Since $a \in \mathbb{C}(z) \subset M_{<1}$, this implies

$$\phi(f) \cdot b \in M_{<1} \otimes_{\mathbb{C}} \mathbf{k},$$

that is, $\phi(f) \in \mathfrak{a}$. So, $\mathfrak{a}$ is a ϕ -ideal. Since $M_{<1} \otimes_{\mathbb{C}} \mathbf{k}$ is ϕ -simple, we must have $1 \in \mathfrak{a}$. So,

$$b \in M_{<1} \otimes_{\mathbb{C}} \mathbf{k}.$$

Choose a $\mathbb{C}$ -basis (c_i) of $\mathbf{k}$ and write $b = \sum_i b_i \otimes c_i$ with $b_i \in M_{<1}$. Then,

$$\sum_i \phi(b_i) \otimes c_i = \phi(b) = ab = \sum_i ab_i \otimes c_i.$$

Hence, for all i , we have $\phi(b_i) = ab_i$. By Lemma 3.2, we conclude that, for all i , $b_i \in \mathbb{C}(z)$, which implies that $b \in \mathbf{k}(z)$, showing (ii). Moreover, since $b \neq 0$, there exists i such that $b_i \neq 0$, showing (i). $\blacksquare$

Theorem 3.5. Let $f \in M$ and $0 \neq a \in \mathbb{C}(z)$ be such that f is a nonzero solution of

$$\phi(y) = ay. \quad (3.10)$$

Then, f is σ -algebraically dependent over $M_{<1}$ if and only if

$$\varphi(a) = \phi(b)/b \quad (3.11)$$

for some $0 \neq b \in \mathbb{C}(z)$ and $1 \neq \varphi(x) = x^{n_0} \sigma(x)^{n_1} \cdots \sigma^{t-1}(x)^{n_{t-1}}$. $\square$

Proof. The converse follows as in Theorem 3.1, noting (3.7), and Corollary 3.1. Let now f be σ -algebraically dependent over $M_{<1}$. As in the proof of Theorem 3.1, we will show that there exists $b \in M_{<1}$ and multiplicative φ such that

$$\varphi(a) = \phi(b)/b.$$

Lemma 3.2 implies that $b \in \mathbb{C}(z)$. To do the above, let L be the smallest $\phi\sigma$ -subfield in M containing $\mathbf{k}$, $M_{<1}$, and f . Since $\mathbf{k} \subset L^\phi \subset M^\phi = \mathbf{k}$, the $\phi\sigma$ -field L is a σ -PV extension over $\mathbf{k} M_{<1}$ for Equation (3.2). It follows from Lemma 2.53 and Proposition 2.48 that f is σ -algebraically dependent over $\mathbf{k} M_{<1}$ if and only if the σ -Galois group G of Equation (3.2) is a proper σ -algebraic subgroup of $\mathbf{G}_m$. Then, by Proposition 2.48, there exists a multiplicative $\varphi \in \mathbf{k}\{x, 1/x\}_\sigma$ such that the ideal of G contains the equation $\varphi(x) = 1$. Therefore, for every $\mathbf{k}$ - σ -algebra B and $g \in G(B)$, we have

$$g(\varphi(f)) = \varphi(g(f)) = \varphi(c_g \cdot f) = \varphi(c_g) \cdot \varphi(f) = 1 \cdot \varphi(f) = \varphi(f).$$

Hence, by Theorem 2.52, we have $b := \varphi(f) \in \mathbf{k} M_{<1}$. Since $f \neq 0$ and φ is multiplicative, $\varphi(f) \neq 0$. Therefore,

$$\varphi(a) = \varphi(\phi(f)/f) = \phi(\varphi(f))/\varphi(f) = \phi(b)/b.$$

By Lemma 3.4, there exists $b' \in M_{<1}$ such that $\varphi(a) = \phi(b')/b'$, which finishes the proof. ■

3.3 How to use the above results in practice

Let $a \in \mathbb{C}(z)^\times$ and $w_0, z_0 \in \mathbb{C}^\times$ and ϕ and σ act on $\mathbb{C}(z)$ as follows:

$$\phi(f)(z) = f(z + w_0) \quad \text{and} \quad \sigma(f)(z) = f(z + z_0), \quad f \in \mathbb{C}(z).$$

Then, for some $N \geq 0$, a can be represented as follows:

$$a = \lambda \cdot \prod_{k=0}^{t-1} \prod_{d=-N-1}^N \prod_{i=1}^R (z - k \cdot z_0 - d \cdot w_0 - r_i)^{s_{k,d,i}},$$

where $\lambda, r_i \in \mathbb{C}$ and the r_i 's are distinct in $\mathbb{C}/w_0 \cdot \mathbb{Z} + z_0 \cdot \mathbb{Z}$. For all i and k , $1 \leq i \leq R$, $0 \leq k \leq t-1$, let

$$a_{i,k} = \sum_{d=-N-1}^N s_{k,d,i}. \quad (3.12)$$

The following result combined with Theorems 3.1 and 3.5 provides a complete characterization of all equations (3.2) whose solutions are σ -algebraically independent.

Theorem 3.6. Let $a \in \mathbb{C}(z)$ be as above and $z_0/w_0 \notin \mathbb{Q}$. Then,

- (i) If λ is a root of unity, then there exist $b \in \mathbb{C}(z)$ and a multiplicative function

$$\varphi(x) = x^{n_0} \cdot (\sigma(x))^{n_1} \cdot \dots \cdot (\sigma^A(x))^{n_A} \neq 1$$

such that $\varphi(a) = \phi(b)/b$ if and only if, for all i , $1 \leq i \leq R$,

$$a_{i,0} = \dots = a_{i,t-1} = 0.$$

- (ii) If λ is not a root of unity, then there exist $b \in \mathbb{C}(z)$ and a multiplicative function

$$\varphi(x) = x^{n_0} \cdot (\sigma(x))^{n_1} \cdot \dots \cdot (\sigma^A(x))^{n_A} \neq 1$$

such that $\varphi(a) = \phi(b)/b$ if and only if, for all i , $1 \leq i \leq R$,

$$a_{i,0} = \dots = a_{i,t-1} = 0 \quad \text{and} \quad t \geq 2.$$

□

Proof. We will write φ and b with undetermined coefficients and exponents. Suppose that

$$b = \mu \cdot \prod_{k=-B}^B \prod_{d=-N}^N \prod_{i=1}^R (z - k \cdot z_0 - d \cdot w_0 - r_i)^{l_{k,d,i}} \quad \text{and} \quad \varphi(x) = x^{n_0} \cdot (\sigma(x))^{n_1} \cdot \dots \cdot (\sigma^A(x))^{n_A}$$

are such that $\varphi(a) = \phi(b)/b$ and $A, B \geq 0$. Let us calculate the right- and left-hand sides of this equality. We see that

$$\phi(b) = \mu \cdot \prod_{k=-B}^B \prod_{d=-N}^N \prod_{i=1}^R (z - k \cdot z_0 - (d-1) \cdot w_0 - r_i)^{l_{k,d,i}}.$$

Hence,

$$\begin{aligned} \frac{\phi(b)}{b} &= \prod_{k=-B}^B \prod_{d=-N-1}^{N-1} \prod_{i=1}^R (z - k \cdot z_0 - d \cdot w_0 - r_i)^{l_{k,d+1,i}} \cdot \prod_{k=-B}^B \prod_{d=-N}^N \prod_{i=1}^R (z - k \cdot z_0 - d \cdot w_0 - r_i)^{-l_{k,d,i}} \\ &= \prod_{k=-B}^B \prod_{i=1}^R \left[(z - k \cdot z_0 + (N+1) \cdot w_0 - r_i)^{l_{k,-N,i}} \right. \\ &\quad \left. \prod_{d=-N}^{N-1} (z - k \cdot z_0 - d \cdot w_0 - r_i)^{l_{k,d+1,i} - l_{k,d,i}} (z - k \cdot z_0 - N \cdot w_0 - r_i)^{-l_{k,N,i}} \right]. \end{aligned}$$

Now, we calculate the left-hand side. We see that, for all $r \geq 0$,

$$\begin{aligned} \sigma^r(a)^{n_r} &= \lambda^{n_r} \cdot \prod_{k=0}^{t-1} \prod_{d=-N-1}^N \prod_{i=1}^R (z - (k-r) \cdot z_0 - d \cdot w_0 - r_i)^{n_r s_{k,d,i}} \\ &= \lambda^{n_r} \cdot \prod_{k=-r}^{t-1-r} \prod_{d=-N-1}^N \prod_{i=1}^R (z - k \cdot z_0 - d \cdot w_0 - r_i)^{n_r s_{r+k,d,i}}. \end{aligned}$$

Hence,

$$\varphi(a) = \lambda^{\sum_{r=0}^A n_r} \cdot \prod_{k=-A}^{t-1} \prod_{d=-N-1}^N \prod_{i=1}^R (z - k \cdot z_0 - d \cdot w_0 - r_i)^{\sum_{\substack{0 \leq r \leq A \\ 0 \leq r+k \leq t-1}} n_r s_{r+k,d,i}}.$$

Now, the equation $\varphi(a) = \phi(b)/b$ gives $A = B = t - 1$ and the following system of linear equations:

$$\left\{ \begin{array}{l} \sum_{0 \leq r, r+k \leq t-1} s_{r+k, -N-1, i} \cdot n_r = l_{k, -N, i}, \\ \sum_{0 \leq r, r+k \leq t-1} s_{r+k, d, i} \cdot n_r = l_{k, d+1, i} - l_{k, d, i}, \quad -N \leq d \leq N-1, \quad 1 \leq i \leq R, \quad 1-t \leq k \leq t-1, \\ \sum_{0 \leq r, r+k \leq t-1} s_{r+k, N, i} \cdot n_r = -l_{k, N, i}, \\ \lambda \sum_{r=0}^{t-1} n_r = 1. \end{array} \right.$$

The first subsystem, for all i and k , $1 \leq i \leq R$, can be rewritten as follows:

$$\left\{ \begin{array}{l} \begin{pmatrix} s_{0, -N-1, i} \\ s_{0, -N, i} \\ \vdots \\ s_{0, N, i} \end{pmatrix} (n_{t-1}) = \begin{pmatrix} l_{k, -N, i} \\ l_{k, -N+1, i} - l_{k, -N, i} \\ \vdots \\ -l_{k, N, i} \end{pmatrix}, \quad k = 1 - t, \\ \begin{pmatrix} s_{0, -N-1, i} & s_{1, -N-1, i} \\ s_{0, -N, i} & s_{1, -N, i} \\ \vdots & \vdots \\ s_{0, N, i} & s_{1, N, i} \end{pmatrix} \begin{pmatrix} n_{t-2} \\ n_{t-1} \end{pmatrix} = \begin{pmatrix} l_{k, -N, i} \\ l_{k, -N+1, i} - l_{k, -N, i} \\ \vdots \\ -l_{k, N, i} \end{pmatrix}, \quad k = 2 - t, \\ \vdots \\ \begin{pmatrix} s_{0, -N-1, i} & s_{1, -N-1, i} & \dots & s_{t-1, -N-1, i} \\ s_{0, -N, i} & s_{1, -N, i} & \dots & s_{t-1, -N, i} \\ \vdots & \vdots & \ddots & \vdots \\ s_{0, N, i} & s_{1, N, i} & \dots & s_{t-1, N, i} \end{pmatrix} \begin{pmatrix} n_0 \\ n_1 \\ \vdots \\ n_{t-1} \end{pmatrix} = \begin{pmatrix} l_{k, -N, i} \\ l_{k, -N+1, i} - l_{k, -N, i} \\ \vdots \\ -l_{k, N, i} \end{pmatrix}, \quad k = 0, \\ \vdots \\ \begin{pmatrix} s_{t-1, -N-1, i} \\ s_{t-1, -N, i} \\ \vdots \\ s_{t-1, N, i} \end{pmatrix} (n_0) = \begin{pmatrix} l_{k, -N, i} \\ l_{k, -N+1, i} - l_{k, -N, i} \\ \vdots \\ -l_{k, N, i} \end{pmatrix}, \quad k = t - 1. \end{array} \right.$$

Each subsystem has a solution in $l_{k,d,i}$ if and only if the sum of all equations is zero. Thus, we can replace this system with the following:

$$\left\{ \begin{array}{l} n_{t-1} \cdot \sum_{d=-N-1}^N s_{0,d,i} = 0, \\ n_{t-2} \cdot \sum_{d=-N-1}^N s_{0,d,i} + n_{t-1} \cdot \sum_{d=-N-1}^N s_{1,d,i} = 0, \\ \vdots \\ n_0 \cdot \sum_{d=-N-1}^N s_{0,d,i} + n_1 \cdot \sum_{d=-N-1}^N s_{1,d,i} + \cdots + n_{t-1} \cdot \sum_{d=-N-1}^N s_{t-1,d,i} = 0, \\ \vdots \\ n_0 \cdot \sum_{d=-N-1}^N s_{t-1,d,i} = 0. \end{array} \right.$$

Using (3.12), we obtain the following system:

$$\begin{pmatrix} 0 & 0 & \cdots & 0 & a_{i,0} \\ 0 & 0 & \cdots & a_{i,0} & a_{i,1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{i,0} & a_{i,1} & \cdots & a_{i,t-2} & a_{i,t-1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{i,t-2} & a_{i,t-1} & \cdots & 0 & 0 \\ a_{i,t-1} & 0 & \cdots & 0 & 0 \end{pmatrix} \begin{pmatrix} n_0 \\ n_1 \\ \vdots \\ n_{t-1} \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}. \quad (3.13)$$

Thus, for some integers $\gamma_{k,d,i,j}$, we have

$$\left\{ \begin{array}{l} \begin{pmatrix} 0 & 0 & \cdots & 0 & a_{i,0} \\ 0 & 0 & \cdots & a_{i,0} & a_{i,1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{i,0} & a_{i,1} & \cdots & a_{i,t-2} & a_{i,t-1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{i,t-2} & a_{i,t-1} & \cdots & 0 & 0 \\ a_{i,t-1} & 0 & \cdots & 0 & 0 \end{pmatrix} \begin{pmatrix} n_0 \\ n_1 \\ \vdots \\ n_{t-1} \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}, \\ \lambda^{\sum_{r=0}^{t-1} n_r} = 1, \\ l_{k,d,i} = \sum_{r=0}^{t-1} \gamma_{k,d,i,r} \cdot n_r. \end{array} \right.$$

Consider the *first case*: λ is a root of unity. Then, for some $u \in \mathbb{Z} \setminus \{0\}$, we have $\lambda^u = 1$. In this situation, if $n_r, l_{k,d,i}$ is a solution of all equations except for the second one, then

$$u \cdot n_r, \quad u \cdot l_{k,d,i}$$

is a solution of the whole system. Therefore, in this case, the existence of φ and b is equivalent to (3.13) having a nontrivial common solution.

Consider the *second case*: λ is not a root of unity. Then, the second equation gives $\sum_{r=0}^{t-1} n_r = 0$. Thus, in this case, we need to show the existence of a nontrivial solution of the system

$$\begin{pmatrix} 0 & 0 & \dots & 0 & a_{i,0} \\ 0 & 0 & \dots & a_{i,0} & a_{i,1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{i,0} & a_{i,1} & \dots & a_{i,t-2} & a_{i,t-1} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ a_{i,t-2} & a_{i,t-1} & \dots & 0 & 0 \\ a_{i,t-1} & 0 & \dots & 0 & 0 \\ 1 & 1 & \dots & 1 & 1 \end{pmatrix} \begin{pmatrix} n_0 \\ n_1 \\ \vdots \\ n_{t-1} \end{pmatrix} = \begin{pmatrix} 0 \\ \vdots \\ 0 \end{pmatrix}. \quad (3.14)$$

Since all the coefficients in (3.13) and (3.14) are integers, there is a nontrivial solution with integral coefficients if and only if there is a nontrivial solution with complex coefficients.

In the *first case*, the rank is $< t$ if and only if

$$a_{i,0} = \dots = a_{i,t-1} = 0.$$

In the *second case*, the rank is $< t$ if and only if

$$a_{i,0} = \dots = a_{i,t-1} = 0 \quad \text{and } t \geq 2. \quad \blacksquare$$

3.4 Examples

We will now illustrate Theorems 3.1, 3.5, and 3.6.

Example 3.7. The gamma function Γ satisfying

$$\Gamma(z+1) = z \cdot \Gamma(z)$$

does not satisfy any polynomial difference equation over $M_{<1}$ (see (3.6)) for any shift by $z_0 \notin \mathbb{Q}$ as, in the notation of Theorem 3.6, $N = 0$, $t = 1$, $R = 1$, and $1 = s_{0,0,1} = a_{1,0} \neq 0$. A differential algebraic independence statement over $M_{<1}$ for Γ was shown in [6] using analytic techniques. Also, [14, Theorem 1] gives difference algebraic independence of the Riemann zeta function ζ over $M_{<1}$. Note the following relation between ζ and Γ :

$$\zeta(1-s) = 2^{1-s} \cdot \pi^{-s} \cdot \cos(\pi \cdot s/2) \cdot \Gamma(s) \cdot \zeta(s). \quad \square$$

Example 3.8. For $f \in K := \mathbb{C}(z, \alpha)$, let

$$\phi(f)(z, \alpha) = f(z, \alpha + 1) \quad \text{and} \quad \sigma(f)(z, \alpha) = f(z + z_0, \alpha).$$

Let F be a $\phi\sigma$ -field over K that contains a nonzero solution of

$$\phi(y) = z \cdot y,$$

which we denote by z^α . Let φ be as in the statement of Theorem 3.1. If z^α were σ -algebraic dependent over $\mathbb{C}(z, \alpha)$, then, by the proof of Theorem 3.1, there would exist $0 \neq b \in F^\phi$ (note that $\mathbb{C}(z) \subset F^\phi$ in our case) such that

$$1 = \phi(b)/b = \varphi(z).$$

Since σ is a shift, $\varphi = 1$, which is a contradiction. This proves the difference algebraic independence of z^α over $\mathbb{C}(z, \alpha)$ with respect to shifts of z (see [7, 27] for a related statement, in which α takes values in $\mathbb{Q}$). $\square$

Example 3.9. Let K be a field. Consider Seq_K as a σ -ring with σ acting as the shift. Let L be a σ -subfield of Seq_K . Consider Seq_L as a $\phi\sigma$ -ring with ϕ acting as the shift and σ acting coordinate-wise. Let F be a $\phi\sigma$ -subfield of Seq_L and $\{S(m, \alpha)\} \in F$ satisfy a first-order ϕ -difference equation

$$S(m, \alpha + 1) = f(m, \alpha) \cdot S(m, \alpha), \quad \{f(m, \alpha)\} \in M,$$

where M is a $\phi\sigma$ -subfield of F , which contains L . Then, it follows from the proof of Theorem 3.1, [23, Lemma A.40; 25, Proposition 1.1] that, if $\{S(m, \alpha)\}$ satisfies a linear

σ -difference equation, then there exists $\{b(m, \alpha)\} \in M$ and $n \geq 1$ such that, for all m and α ,

$$S(m+n, \alpha) = b(m, \alpha) \cdot S(m, \alpha). \quad (3.15)$$

In particular, we can take M to be the image of $L(z)$ in Seq_L , as in Section 2.5. Let

$$y(x, \alpha) := \sum_{m \geq 0} S(m, \alpha) \cdot x^m.$$

By [54, Theorem 1.5], the function $y(x, \alpha)$ satisfies a linear differential equation in x if and only if $S(m, \alpha)$ satisfies a homogeneous linear difference equation in m (see also [26, App. B.4] and the reference given there). Suppose it is known that $S(m, \alpha)$ satisfies a first-order homogeneous linear difference equation with respect to α , and one wants to know whether $y(x, \alpha)$ satisfies a linear differential equation in x . The above method helps find difference equations in m if they are hard to find otherwise, as such equations are all of the form (3.15). Just to illustrate the process (but not the difficulty), consider the Bessel functions of the first kind, which are given by

$$J_\alpha(x) = \sum_{m \geq 0} \frac{(-1)^m}{m! \cdot \Gamma(m + \alpha + 1)} (x/2)^{2m + \alpha},$$

where α is an integer. It is a solution of the following differential equation:

$$x^2 y'' + x y' + (x^2 - \alpha^2) \cdot y = 0, \quad (3.16)$$

where $'$ stands for $\frac{d}{dx}$. Let

$$S(m, \alpha) = \frac{(-1)^m}{m! \cdot \Gamma(m + \alpha + 1)} \quad \text{and} \quad I_\alpha(x) = \sum_{m \geq 0} S(m, \alpha) x^m.$$

Then, $J_\alpha(x) = (x/2)^\alpha \cdot I_\alpha(x^2/4)$. We have

$$(m + \alpha + 1) \cdot S(m, \alpha + 1) = S(m, \alpha).$$

Moreover, we have

$$(m + 1)(m + \alpha + 1) \cdot S(m + 1, \alpha) + S(m, \alpha) = 0.$$

Therefore, by a calculation using the Gfun package in MAPLE [53], $I_\alpha(x)$ satisfies the second-order linear differential equation

$$xy'' + y' + y = 0 \quad (3.17)$$

(see also the proof of [54, Theorem 1.5]). One now obtains (3.16) by substituting the expression of I_α in terms of J_α into (3.17). $\square$

Acknowledgments

We are grateful to S. Abramov, Y. André, D. Drasin, D. Khmelnov, S. Merenkov, J. Roques, D. Trushin, and the referee for their helpful suggestions.

Funding

A.O. was supported by the grants: NSF CCF-0952591 and PSC-CUNY No. 60001-40 41.

References

- [1] Amano, K. "Liouville extensions of Artinian simple module algebras." *Communications in Algebra* 34, no. 5 (2006): 1811–23. URL <http://dx.doi.org/10.1080/00927870500542796>.
- [2] Amano, K. and A. Masuoka. "Picard–Vessiot extensions of Artinian simple module algebras." *Journal of Algebra* 285, no. 2 (2005): 743–67. ISSN 0021-8693. URL <http://dx.doi.org/10.1016/j.jalgebra.2004.12.006>.
- [3] Amano, K., A. Masuoka, and M. Takeuchi. "Hopf Algebraic Approach to Picard–Vessiot Theory." In *Handbook of Algebra*, vol. 6, 127–71. Amsterdam: Elsevier/North-Holland, 2009. URL [http://dx.doi.org/10.1016/S1570-7954\(08\)00204-0](http://dx.doi.org/10.1016/S1570-7954(08)00204-0).
- [4] André, Y. "Différentielles non-commutatives et théorie de Galois différentielle ou aux différences." *Annales Scientifiques de l'École Normale Supérieure. Quatrième Série* 34, no. 5 (2001): 685–739. URL [http://dx.doi.org/10.1016/S0012-9593\(01\)01074-6](http://dx.doi.org/10.1016/S0012-9593(01)01074-6).
- [5] Antieau, B., A. Ovchinnikov, and D. Trushin. "Galois theory of difference equations with periodic parameters." *Communications in Algebra* 42, no. 9 (2014): 3902–43. URL <http://dx.doi.org/10.1080/00927872.2013.797991>.
- [6] Bank, S. and R. Kaufman. "An extension of Hölder's theorem concerning the gamma function." *Funkcialaj Ekvacioj. Serio Internacia* 19, no. 1 (1976): 53–63. URL http://fe.math.kobe-u.ac.jp/FE/FE_pdf_with_bookmark/FE11-20-en_KML/fe19%-053-063/fe19-053-063.pdf.
- [7] Bell, J., S. Gerhold, M. Klazar, and F. Luca. "Non-holonomicity of sequences defined via elementary functions." *Annals of Combinatorics* 12, no. 1 (2008): 1–16. URL <http://dx.doi.org/10.1007/s00026-008-0333-6>.
- [8] Bourbaki, N. *Algèbre Commutative. Chapitres 1 à 4*. "Éléments de Mathématique. Berlin: Springer, 2006. URL <http://dx.doi.org/10.1007/978-3-540-33976-2>.

- [9] Bourbaki, N. *Algèbre Commutative. Chapitres 5 à 7. Éléments de Mathématique*. Berlin: Springer, 2006. URL <http://dx.doi.org/10.1007/978-3-540-33979-3>.
- [10] Cassidy, P. and M. Singer. "Galois theory of parametrized differential equations and linear differential algebraic groups." *IRMA Lectures in Mathematics and Theoretical Physics* 9 (2007) 113–57. URL <http://dx.doi.org/10.4171/020-1/7>.
- [11] Chatzidakis, Z. and E. Hrushovski. "Model theory of difference fields." *Transactions of the American Mathematical Society* 351, no. 8 (1999): 2997–3071. ISSN 0002-9947. URL <http://dx.doi.org/10.1090/S0002-9947-99-02498-8>.
- [12] Chatzidakis, Z., E. Hrushovski, and Y. Peterzil. "Model Theory of Difference Fields. II. Periodic ideals and the Trichotomy in all Characteristics." *Proceedings of the London Mathematical Society. Third Series*, 85, no. 2 (2002): 257–311. ISSN 0024-6115. URL <http://dx.doi.org/10.1112/S0024611502013576>.
- [13] Chatzidakis, Z. C. Hardouin, and M. F. Singer. "On the Definitions of Difference Galois Groups." In *Model Theory with Applications to Algebra and Analysis*, Vol. 1, 73–109. London Mathematical Society Lecture Note Series 349. Cambridge: Cambridge University Press, 2008. URL <http://dx.doi.org/10.1017/CBO9780511735226.006>.
- [14] Chiang, Y.-M. and S.-J. Feng. "Difference independence of the Riemann zeta function." *Acta Arithmetica* 125, no. 4 (2006): 317–29. URL <http://dx.doi.org/10.4064/aa125-4-2>.
- [15] Chyzak, F. and B. Salvy. "Non-commutative elimination in Ore algebras proves multivariate identities." *Journal of Symbolic Computation* 26, no. 2 (1998): 187–227. URL <http://dx.doi.org/10.1006/jSCO.1998.0207>.
- [16] Cohn, R. M. *Difference Algebra*. New York–London–Sydney: Interscience Publishers, John Wiley & Sons, 1965.
- [17] Di Vizio, L. "Approche galoisienne de la transcendance différentielle." *Journée annuelle 2012 de la SMF*, 2012. URL <http://divizio.perso.math.cnrs.fr/PREPRINTS/16-JourneeAnnuelleSMF/difftransc.pdf>.
- [18] Di Vizio, L. and C. Hardouin. "Courbures, groupes de Galois génériques et D -groupoïdes de Galois d'un système aux q -différences." *Comptes Rendus Mathématique de l'Académie des sciences. Paris* 348, no. 17–18 (2010): 951–4. URL <http://dx.doi.org/10.1016/j.crma.2010.08.001>.
- [19] Di Vizio, L. and C. Hardouin. "Galois theories for q -difference equations: comparison theorems." (2011): preprint. URL <http://arxiv.org/abs/1002.4839>.
- [20] Di Vizio, L. and C. Hardouin. "On the Grothendieck conjecture on p -curvatures for q -difference equations." (2011): preprint. URL <http://arxiv.org/abs/1002.4839>.
- [21] Di Vizio, L. and C. Hardouin. "Parameterized generic Galois groups for q -difference equations, followed by the appendix 'The Galois D -groupoid of a q -difference system' by Anne Granier." (2011): preprint. URL <http://arxiv.org/abs/1002.4839>.
- [22] Di Vizio, L. and C. Hardouin. "Descent for differential Galois theory of difference equations. Confluence and q -dependency." *Pacific Journal of Mathematics* 256, no. 1 (2012): 79–104. URL <http://dx.doi.org/10.2140/pjm.2012.256.79>.

- [23] Di Vizio, L., C. Hardouin, and M. Wibmer. "Difference Galois theory of linear differential equations." to appear in *Advances in Mathematics* (2014). doi: 10.1016/j.aim.2014.04.005. URL <http://arxiv.org/abs/1302.7198>.
- [24] Eisenbud, D. *Commutative Algebra with a View Toward Algebraic Geometry*. Graduate Texts in Mathematics 150. Berlin: Springer, 1995.
- [25] Eisenbud, D. and B. Sturmfels. "Binomial ideals." *Duke Mathematical Journal* 84, no. 1 (1996). URL <http://dx.doi.org/10.1215/S0012-7094-96-08401-X>.
- [26] Flajolet, P. and R. Sedgewick. *Analytic Combinatorics*. Cambridge: Cambridge University Press, 2009. URL <http://dx.doi.org/10.1017/CBO9780511801655>.
- [27] Flajolet, P., S. Gerhold, and B. Salvy. "On the non-holonomic character of logarithms, powers, and the n th prime function." *Electronic Journal of Combinatorics* 11, no. 2 (2005): #A2. URL http://www.combinatorics.org/Volume_11/PDF/v11i2a2.pdf.
- [28] Gillet, H., S. Gorchinskiy, and A. Ovchinnikov. "Parameterized Picard–Vessiot extensions and Atiyah extensions." *Advances in Mathematics* 238 (2013): 322–411. URL <http://dx.doi.org/10.1016/j.aim.2013.02.006>.
- [29] Goldberg, A. and I. Ostrovskii. *Value Distribution of Meromorphic Functions*. Translations of Mathematical Monographs 236. Providence, RI: American Mathematical Society, 2008.
- [30] Gorchinskiy, S. and A. Ovchinnikov. "Isomonodromic differential equations and differential categories." *Journal de Mathématiques Pures et Appliquées*, 2014. URL <http://dx.doi.org/10.1016/j.matpur.2013.11.001>.
- [31] Granier, A. "A Galois D -groupoid for q -difference equations." *Annales de l'Institut Fourier* 61, no. 4 (2011): 1493–516. URL <http://dx.doi.org/10.5802/aif.2648>.
- [32] Grothendieck, A. "Éléments de géométrie algébrique (rédigés avec la collaboration de Jean Dieudonné): IV. Étude locale des schémas et des morphismes de schémas." *Publications Mathématiques de l'IHÉS* 20 (1964) 5–259. URL <http://www.numdam.org/item?id=PMIHES.1964.20.5.0>.
- [33] Hardouin, C. "Hypertranscendence et groupes de Galois aux différences." 2006. URL <http://arxiv.org/abs/math/0609646>.
- [34] Hardouin, C. "Hypertranscendence des systèmes aux différences diagonaux." *Compositio Mathematica* 144, no. 3 (2008) 565–81. URL <http://dx.doi.org/10.1112/S0010437X07003430>.
- [35] Hardouin, C. and M. Singer. "Differential independence of solutions of a class of q -hypergeometric difference equations." Maple worksheet, 2007. URL <http://www4.ncsu.edu/~singer/ms.papers.html>.
- [36] Hardouin, C. and M. F. Singer. "Differential Galois theory of linear difference equations." *Mathematische Annalen* 342, no. 2 (2008): 333–77. URL <http://dx.doi.org/10.1007/s00208-008-0238-z>.
- [37] Hay, M., J. Hietarinta, N. Joshi, and F. Nijhoff. "A Lax pair for a lattice modified KdV equation, reductions to q -Painlevé equations and associated Lax pairs." *Journal of Physics A: Mathematical and Theoretical* 40 (2007) F61–F73. URL <http://dx.doi.org/10.1088/1751-8113/40/2/F02>.

- [38] Joshi, N. and Y. Shi. "Exact solutions of a q -discrete second Painlevé equation from its isomonodromy deformation problem: I. Rational solutions." *Proceedings of the Royal Society A: Mathematical, Physical, and Engineering Sciences* 467, no. 2136 (2011): 3443–68. URL <http://dx.doi.org/10.1098/rspa.2011.0167>.
- [39] Joshi, N. and Y. Shi. "Exact solutions of a q -discrete second Painlevé equation from its isomonodromy deformation problem. II. Hypergeometric solutions." *Proceedings of the Royal Society A: Mathematical, Physical, and Engineering Sciences* 468, no. 2146 (2012): 3247–64. URL <http://dx.doi.org/10.1098/rspa.2012.0224>.
- [40] Kamensky, M. "Tannakian formalism over fields with operators." *International Mathematics Research Notices* 2013, no. 24 (2013): 5571–622. URL <http://dx.doi.org/10.1093/imrn/rns190>.
- [41] Kolchin, E. R. "Constrained extensions of differential fields." *Advances in Mathematics* 12 (1974) 141–70. URL [http://dx.doi.org/10.1016/S0001-8708\(74\)80001-0](http://dx.doi.org/10.1016/S0001-8708(74)80001-0).
- [42] Lang, S. *Algebra*, rev. 3rd ed. Graduate Texts in Mathematics 211. New York: Springer, 2002. URL <http://dx.doi.org/10.1007/978-1-4613-0041-0>.
- [43] Levin, A. *Difference Algebra*. Algebra and Applications 8. New York: Springer, 2008. ISBN 978-1-4020-6946-8. URL <http://dx.doi.org/10.1007/978-1-4020-6947-5>.
- [44] Morikawa, S. "On a general difference Galois theory. I." *Annales de l'Institut Fourier* 59, no. 7 (2009): 2709–32. URL <http://dx.doi.org/10.5802/aif.2505>.
- [45] Murata, M. "Lax forms of the q -Painlevé equations." *Journal of Physics A: Mathematical and Theoretical* 42, no. 11 (2009): 115201. URL <http://dx.doi.org/10.1088/1751-8113/42/11/115201>.
- [46] Nevanlinna, R. *La théorème de Picard–Borel et la théorie des fonctions méromorphes*, 2nd ed. Bronx, New York: Chelsea Publishing Company, 1974.
- [47] Nichols, W. "Quotients of Hopf algebras." *Communications in Algebra* 6, no. 17 (1978) 1789–800. URL <http://dx.doi.org/10.1080/00927877808822321>.
- [48] Ovchinnikov, A. "Difference integrability conditions for parameterized linear difference and differential equations." *Advances in Applied Mathematics* 53 (2014): 61–71. URL <http://dx.doi.org/10.1016/j.aam.2013.09.007>.
- [49] Ovchinnikov, A. and M. Wibmer. "Tannakian categories with semigroup actions." (2014) preprint. URL <http://arxiv.org/abs/1403.3850>.
- [50] Peón Nieto, A. "On $\sigma\delta$ -Picard–Vessiot extensions." *Communications in Algebra* 39, no. 4 (2011): 1242–9. URL <http://dx.doi.org/10.1080/00927871003645391>.
- [51] Roques, J. "Galois groups of basic hypergeometric equations." *Pacific Journal of Mathematics* 235, no. 2 (2008): 303–22. URL <http://dx.doi.org/10.2140/pjm.2008.235.303>.
- [52] Salvy, B. " D -Finiteness: Algorithms and Applications." In *Proceedings of the International Symposium on Symbolic and Algebraic Computation (ISSAC) 2005*, 2–3. New York, NY: ACM Press, 2005. URL <http://dx.doi.org/10.1145/1073884.1073886>.
- [53] Salvy, B. and P. Zimmermann. "Gfun: a Maple package for the manipulation of generating and holonomic functions in one variable." *ACM Transactions on Mathematical Software* 20, no. 2 (1994): 163–77. URL <http://dx.doi.org/10.1145/178365.178368>.
- [54] Stanley, R. "Differentiably finite power series." *European Journal of Combinatorics* 1, no. 2 (1980): 175–88. URL <http://www-math.mit.edu/~rstan/pubs/pubfiles/45.pdf>.

- [55] Takano, K. "On the hypertranscendence of solutions of a difference equation of Kimura." *Funkcialaj Ekvacioj. Serio Internacia* 16 (1973): 241–54. URL http://fe.math.kobe-u.ac.jp/FE/FE_pdf_with_bookmark/FE11-20-en_KML/fe16-241-254/fe16-241-254.pdf.
- [56] Trushin, D. "Difference Nullstellensatz." (2009). URL <http://arxiv.org/abs/0908.3865>.
- [57] Trushin, D. "Difference Nullstellensatz in the case of finite group." (2009). URL <http://arxiv.org/abs/0908.3863>.
- [58] van der Put, M. and M. F. Singer. *Galois Theory of Difference Equations*. Lecture Notes in Mathematics 1666. Berlin: Springer, 1997. ISBN 3-540-63243-3. URL <http://dx.doi.org/10.1007/BFb0096118>.
- [59] Waterhouse, W. *Introduction to Affine Group Schemes*. New York: Springer, 1979. URL <http://dx.doi.org/10.1007/978-1-4612-6217-6>.
- [60] Wibmer, M. "Geometric difference Galois theory." PhD thesis, Heidelberg, 2010. URL <http://www.ub.uni-heidelberg.de/archiv/10685>.
- [61] Wibmer, M. "A Chevalley theorem for difference equations." *Mathematische Annalen* 354, no. 4 (2012): 1369–96, ISSN 0025-5831. URL <http://dx.doi.org/10.1007/s00208-011-0770-0>.
- [62] Wibmer, M. "Existence of ∂ -parameterized Picard–Vessiot extensions over fields with algebraically closed constants." *Journal of Algebra* 361 (2012): 163–71. URL <http://dx.doi.org/10.1016/j.jalgebra.2012.03.035>.
- [63] Wibmer, M. "On the Galois Theory of Strongly Normal Differential and Difference Extensions." In *Geometric and differential Galois theories*, edited by D. Bertrand, P. Boalch, J.-M. Couveignes, and P. Dèbes, 221–40. Séminaires et Congrès 27, 2012. URL http://smf4.emath.fr/en/Publications/SeminairesCongres/2012/27/pdf/smf_sem-cong_27_221-240.pdf.
- [64] Wibmer, M. "Skolem–Mahler–Lech type theorems and Picard–Vessiot theory." *Journal of the European Mathematical Society* (2014). URL <http://arxiv.org/abs/1203.1449>.